

# Fungal population on seeds of *Arachis hypogea* L.

Jogdand S.K.\* and Talekar S.M.

P.G. Department of Botany, Mrs.K.S.K. College Beed- 431122 (M.S.), India

\*Corresponding author, Email: ram\_mango@yahoo.com

## Keywords

Ground nut  
Seed  
Mycoflora  
Agar plate  
Standard blotter paper  
method

## Abstract

The seed of four groundnut varieties viz. Gujrat, Western, Ghungroo and Local were collected from different market places of Beed, (M.S.) and seed mycoflora was isolated by standard blotter paper method and agar plate method. In all the four varieties seeds exhibited maximum number of fungi with higher percentage of incidence. *Aspergillus flavus*, *A. niger*, *Fusarium oxysporum*, *Macrophomina phaseolina* *Penicillium* sp. were found predominant. Higher numbers of fungi were isolated on agar plate method used as compared to standard blotter paper method. Surface sterilization with HgCl<sub>2</sub> reduces the incidence of *Aspergillus flavus* and *Aspergillus niger*.

## 1. Introduction

Groundnut (*Arachis hypogea* L.) a valuable legume crop is also known as peanut. It is annual, wet season plant grown in many tropical and temperate countries of the world. Groundnut seed contain 50% edible oil. Seeds are rich in fats, protein, vitamin, B1, B2, B6 and nicotinic acid. It is also good source of lecithin present to the extent of 0.5-0.7% in decorticated nuts. Groundnut flour is suitable for supplementing white flour (Sastri, 1948). Various diseases caused by organisms *Fusarium solani*, *F.oxysporium* cause damping off of groundnut seedlings (Reddy and Rao, 1980). *Aspergillus* attacks germinating groundnut seed (Clinton, 1960) *Aspergillus niger* caused crown rot disease of peanut (Gibson, 1953). Many workers have detected mold fungi and their toxin production ability in stored grains, which deteriorate the stored products (Afzal *et al.*, 1979; Vedahayagam *et al.*, 189). Therefore experiments were carried out to determine the composition of the mycoflora of groundnut seeds which is presented here.

## 2. Materials and Methods

The seed of four groundnut varieties viz. Gujrat, Western, Ghungroo and Local were collected from the different market places of Beed (M.S). For the isolation of seed mycoflora associated with seed samples, the method recommended by ISTA (1966) was adopted. For the standard blotter paper method technique, untreated seeds and treated seeds with 1% HgCl<sub>2</sub> were placed on three layers of moistened slandered blotter paper, ten seeds per petri dish. For agar plate method, the treated and untreated seeds were placed on potato dextrose agar (PDA), ten seeds

per petri dish and were incubated at 24° C for 7 days. Fungi were identified by standard literature.

## 3. Results and Discussion

It is clear from table 1 and 2 that total numbers of 17 fungi were isolated namely. *Macrophomina phaseolina*, *Penicillium* sp., *Alternaria alternate*, *Alternaria tenuis*, *A. carthami*, *Aspergillus candidus*, *A. flavus*, *A. niger*, *A. ustus*, *A. terreus*, *A. fumigatus*, *Curvularia lunata*, *Fusarium oxysporum*, *F. moniliformae*, *F. equiseti*, *Macrophomina phaseolina*, *Penicillium* sp., *Rhizopus nigricans*, *Trichoderma viride* and *T. harzianum* were isolated from four varieties of groundnut seeds by agar plate and blotter paper method. It is interesting to note that their percent incidence is more on agar plate than on blotter papers. Among these fungi, five species of *Aspergillus*, four species of *Alternaria*, three species of *Fusarium* and two species of *Trichoderma* were dominant. Surface sterilization with 1% Hg Cl<sub>2</sub> significantly reduced the incidence of *A.flavus* and *A. niger*.

Gupta and Chauhan (1970) detected *Aspergillus niger*, *A.fumigatus*, *M. Phaseolina*, *Fusarium oxysporum*, *Rhizopus arrhizus*, *Neocosmospora Vasinpect*, *Peacilomyces varioti*, *Alternaria tenuis*, *Penicillium* sp. and *Curvularia sp.* are the maximum count. Species of *Aspergillus*, *Penicillium* and *Rhizopus* have also been reported on groundnut seed (Lumpungu *et al.*, 1989). These species reduces the germination of seeds and damaged the seeds in storage (Christemen, 1973). *Fusarium solani* and *F. oxysporum* cause damping off of groundnut seedling (Reddy and Rao, 1980). Therefore there is need for reducing the fungal growth and mycotoxin production in groundnut seeds by improving the storage condition.

Table 1 Incidence of seed mycoflora on different groundnut varieties on PDA medium

Fungi	Groundnut varieties							
	Local		Ghungroo		Western		Gujrat	
	T	UT	T	UT	T	UT	T	UT
<i>Alternaria alternata</i>	15	25	12	20	-	10	-	07
<i>A. tenuis</i>	-	16	-	18	06	14	-	-
<i>A. carthami</i>	10	13	15	40	12	25	-	-
<i>Aspergillus niger</i>	20	50	16	60	07	30	12	05
<i>A. flavus</i>	25	80	10	26	-	08	10	10
<i>A. ustus</i>	07	12	-	25	-	10	-	01
<i>A. terreus</i>	02	39	08	10	-	-	-	03
<i>A. fumigatus</i>	-	40	-	-	-	-	-	-
<i>Curvularia lunata</i>	06	-	-	12	02	08	-	01
<i>Fusarium oxysporum</i>	06	12	02	10	-	02	02	05
<i>F. moniliforme</i>	-	15	-	17	-	-	-	02
<i>F. equiseti</i>	09	10	-	20	-	-	-	03
<i>Macrophomina Phaseolina</i>	02	32	02	22	02	02	-	04
<i>Penicillium sp.</i>	06	35	07	16	-	01	-	02
<i>Rhizopus nigricans</i>	04	30	-	02	-	-	-	01
<i>Trichoderma viride</i>	-	19	-	01	-	-	-	-
	-	07	-	01	-	-	-	-

T-Treated seeds, UT – Untreated seeds

Table 2. Incidence of fungi on different varieties of groundnut on standard blotter paper

Fungi	Groundnut varieties							
	Local		Ghungroo		Western		Gujrat	
	T	UT	T	UT	T	UT	T	UT
<i>Alternaria alteranata</i>	12	56	02	25	02	22	-	15
<i>A. tenuis</i>	10	47	08	45	06	30	-	20
<i>A. carthami</i>	-	22	-	20	-	15	-	10
<i>Aspergillus candidus</i>	-	12	-	10	-	10	-	09
<i>A. niger</i>	10	47	05	66	10	60	01	40
<i>A. flavus</i>	20	77	10	80	10	80	04	35
<i>A. ustus</i>	03	12	-	10	-	12	-	10
<i>A. terreus</i>	-	22	-	30	-	14	-	02
<i>A. fumigatus</i>	-	40	-	32	-	35	-	08
<i>Curvularia lunata</i>	09	39	03	25	01	13	-	-
<i>Fusarium oxysporium</i>	10	50	04	14	01	18	-	04
<i>F. moniliformi</i>	12	47	-	15	-	22	-	-
<i>F. equiseti</i>	-	12	-	20	-	22	-	04
<i>Macrophomina phaseolina</i>	-	17	10	30	02	09	-	-
<i>Penicillium sp.</i>	05	22	03	32	-	10	-	10
<i>Rhizopus nigricans</i>	02	32	7	30	-	02	-	-
<i>Trichoderma viride</i>	-	32	02	-	-	02	-	-

T-Treated seeds, UT – Untreated seeds

## References

Afzal, M.R., Cheena A. and Chaudhary. (1979).  
Incidence of aflatoxin producing fungi in

animal feed stuff. Mycopathologia, 69 (3) : 149-51.

- Christensen, C.M. (1973). Loss of viability in storage microflora. *Seed sci. and Technol.*, 1 : 547-562.
- Clinton, R.S. (1960). Seed bed pathogen of groundnut in Sudan, and an attempt at control with an artificial test a. *Empire J. Exptl. Agr.*, 28 : 211-222.
- Gibson, I.A.S. (1953). Crown rot, a seedling disease of groundnut caused by *Aspergillus niger*. *Trans. Brit. Mycol. Soc.*, 36 : 198-209.
- Gupta, V.K. and J.S. Chavhan (1970). Seed borne fungi and seed health testing in relation to seedling disease of groundnut. *Indian Phytopath* 23 (4): 622-625.
- ISTA (1966). Proc. International Seed Testing Association, 31:1-52.
- Lumpungu, K., Baelenge, B. and Bitijula, M. (1989). The effect of groundnut seed coat on the development of pathogenic fungi. *Tropicultura*, 17(4): 128-131.
- Neergaard, Pavl (1977). Seed Pathology, Vol.1 John Willey and Sons, W.Y.
- Reddy, M.N. and Rao, A.S. (1980). Dampinoff of groundnut caused by 2 *Fusarium spp.*, *F.Solani* and *F.oxysporum*. *Indian J. Myco. and pl. Patho*, 10 (1) : 65-66.
- Sastri, B. N. (1948). *The wealth of India*. A dictionary of raw materials and industrial products. Vol.I. Council of scientific and Industrial Research. New Delhi. pp. 253.
- Vedahayagam, H.S. Indulkar A. and Rao, S. (1989). The occurrence of *Aspergillus flavus* and aflatoxin in India cotton seed. *J. Food, Sci. Tech*, 23(2):73-76.

---

**Please Cite This Article As:**

S.K. Jogdand and S.M. Talekar. 2010. Fungal population on seeds of *Arachis hypogea* L. J. Ecobiotechnol. 2(6):11-13.