

# Extracellular Hydrolytic Enzymes Action of *Alternaria* Species under the Influence of Different Nutritional Sources

**Sulochana R. Rathod\* and Ashok M. Chavan**

Seed Pathology and Fungal Biotechnology Laboratory, Department of Botany, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad- 431004(M.S.) India

\*Corresponding author, Email: [ram\\_mango@yahoo.com](mailto:ram_mango@yahoo.com)

## Keywords

Amylase  
Protease  
Lipase  
*Alternaria*

## Abstract

Influence of nutritional sources like carbohydrates, nitrogen, phosphorous and sulphur on amylase, lipase and protease action of six *Alternaria* species viz. *A. alternata*, *A. citri*, *A. crassa*, *A. macrospora*, *A. dianthicola* and *A. tenuissima* were studied. It was reported that Carboxy methyl cellulose (CMC), ferrous sulphate, calcium sulphate, sodium sulphate, copper sulphate and sodium dihydrogen orthophosphate retarded the enzyme action of some *Alternaria* species.

## 1. Introduction

Seed is the basic and most vital input in crop production. About 90 % of world food crops are produced by using seeds. The seeds are also found to be responsible for transmission of diseases. More than 3000 diseases are known to be transmitted through seeds. This transmission takes place either in the field or in storage conditions. It is accepted that over all losses due to diseases can vary from 10 – 25 % annually through out the world. In India even it will take conservative estimates of around 15 % losses we are losing an average 30 metric tonnes of food grains, 4 metric tonnes of oilseed, 36 metric tonnes of cane, 23 metric tonnes of fruit and vegetables. The incidence of seed-borne pathogens mainly depends on climatic or physical conditions under which the seed crops are grown in the field. Similarly, storage conditions also enhance to develop various types of seed damages due to associated seed-borne pathogens such seeds show great loss in their chemical content. Neergaard (1977) reported several types of such abnormalities which mainly include seed abortions, sunken seeds, reduction in seed size, seed rots sclerotisation of seed, seed necrosis, loss in germinability, seed discolouration, toxification and other physiological disorders. Such seeds are considered poor in quality for seed industry and also for consumption. Fungi secrete hydrolytic enzymes and spoil the seed contents. Considering the fact attempt were made to study the impact of nutritional sources on hydrolytic enzyme production of *Alternaria* species.

## 2. Materials and Methods

### Production of amylase

Production of amylase(s) was studied by growing the fungi in liquid medium containing 1%, KNO<sub>3</sub> 0.25%, KH<sub>2</sub>PO<sub>4</sub> 0.1.% and MgSO<sub>4</sub>.7H<sub>2</sub>O 0.05%, pH of the medium was adjusted at 5.5. twenty five ml of the medium was poured in 100ml conical flasks autoclaved and inoculated separately with 01 ml spore suspension of the fungi which were grown for 7 days on PDA slants. Unless otherwise stated, the flasks were incubated for 6 days at 25 ± 1°C with diurnal periodicity of light. On 7<sup>th</sup> day, the flasks were harvested by filtering the contents through Whatman filter No.1. The filtrates were collected in presterilized bottles and termed as crude enzyme preparation.

### Assay method for amylase enzymes (Cup-plate method)

Determination of amylase activity was done with the help of cup-plate method which was adopted by Singh and Saxena (1982), where 20ml of starch agar assay medium (soluble starch – 10gm, Na<sub>2</sub>HPO<sub>4</sub> – 2.84gm, NaCl – 0.35gm, Agar agar 20gm, distilled water 1000ml and pH 6.9) was poured in each petriplate. On solidification of the medium, a cavity (08 mm diameter) was made in the centre with the help of a cork borer (No.4) and was filled with 1ml culture filtrates (crude enzyme preparation) of the test fungi. The plates were incubated at 28°C for 24 hours, then they were flooded with Lugol's iodine solution as an indicator. A clear, non blue, circular zone obtained surrounding the central cavity; diameter of the zone was measured (mm) as the amylase activity zone. Similar

procedure followed for the control except pouring of culture filtrates in the central cavity instead of the activity enzyme.

### Production of lipase

Lipase activity was studied by growing the fungi in liquid medium at pH 5.6 containing oil-10ml,  $\text{KNO}_3$  -2.5g,  $\text{KH}_2\text{PO}_4$  -1.0g,  $\text{MgSO}_2$  - 0.5g and distilled water 1000ml. Different sources of carbon, nitrogen, phosphorus and sulphur sources were added separately by replacing corresponding compounds in the above basal medium. 25ml of the medium was poured in 100ml conical flasks and autoclaved at 15 lbs pressure for 30 minutes, then on cooling the flasks were inoculated separately with 1.0ml spore suspension of the fungi which were incubated for 7 days at  $25 \pm 1^\circ\text{C}$  with diurnal periodicity of light. On 7<sup>th</sup> day, the flasks were harvested by filtering the contents through Whatman filter paper no.1. The filtrates were collected in presterilized culture filtrate bottles and termed as crude lipase.

### Assay Method (Cup-plate method)

Determination of lipase activity was done with the help of cup-plate method (Sierra, 1957.) The medium contains Difco peptone-10g, NaCl-5g,  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ -1.0g, agar 20g and 10ml lipid substrate Serbitan mono laurate (Tween-20) (Pre-sterilized), distilled water- 1000ml was added to it. The pH of the medium was adjusted to 6.00. The medium was poured in each Petri plate. On solidifying the medium with the help of a cork borer (No.4) of 8mm diameter well was made in the centre of the plate and was filled with 0.1ml culture filtrate. The plates were incubated at  $28^\circ\text{C}$ . After 24 hours, a clear circular zone was measured (mm) as lipase activity.

### Production of protease

Production of protease(s) was made by growing the fungi on liquid medium containing glucose 10g, gelatine 10g, dipotassium hydrogen phosphate 1.0g,  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  500mg and distilled water 1000ml pH of the medium was adjusted at 5.5. Twenty five ml of medium was poured in 100ml Erlenmeyer conical flasks and autoclaved at 15lbs pressure for 20 minutes. The flasks on cooling were inoculated separately with 01 ml standard spore/mycelial suspension of test fungi prepared from 7 days old cultures grown on PDA slants. The flasks were incubated for 6 days at  $25 \pm 1^\circ\text{C}$  with diurnal periodicity of light. On 7<sup>th</sup> day, the flasks were harvested by filtering the contents through Whatman's filter No.1. The filtrates were collected in the

presterilised bottles and termed as crude enzyme preparation.

### Assay method (Cup-plate method)

Determination of protease(s) activity was done with the help of cup plate method, adopted by Hislop *et.al.* (1982) and Rajamani (1990). A basal medium was prepared by adding 2 % (W/V) agar and one percent (W/V) gelatin. pH of the medium was adjusted at 5.6 with McIlvaine's buffer. Then it was sterilized at 15lbs pressure for 15 minutes. About 15ml of the medium was poured in presterilized petriplates under aseptic conditions. On solidification 6mm diameter cups/cavities were made in the centre of each of the agar plate with a sterilized cork borer (No.4). The cups/cavities were filled carefully with about 0.5ml of culture filtrate (crude enzyme preparation). The plates were incubated at  $25^\circ\text{C}$  for 24 hours. Then the plates were flooded with 15 percent mercuric chloride in 7NHCl. After 10 minutes of standing, a clear transparent zone indicated the hydrolysis of gelatin by extra cellular proteolytic enzymes, whereas the rest of the region of the petriplates became opaque due to the coagulation of gelatin (protein) by mercuric chloride. Diameter of the clear zone was used as measure (mm) of protease activity, while non appearance of clear zone considered absence of protease(s) in the culture filtrates.

## 3. Results and Discussion

It is observed from the table 1 that all six species of *Alternaria* produced amylase. However CMC proved highly inhibitory for amylase production where as glucose and fructose stimulated the amylase activity in *Alternaria alternata*, *A. crassa*, *A. dianthicola* and *A. tenuissima*. Lipase production was found to be stimulated in the presence of disaccharides and polysaccharides in all the species of *Alternaria*. Protease activity was favored in the presence of fructose & sucrose as compared with glucose. However *Alternaria citri*, *A. crassa* and *A. dianthicola* retarded the activity in the presence of CMC.

Amylase production of *A. citri* totally inhibited by sodium nitrites whereas amylase action of most of the *Alternaria* species stimulated in presence of peptone, gelatin, casein, urea and sodium nitrate. It was observed that all the species of *Alternaria* stimulated lipase and protease production in the presence of different nitrogen sources (Table 2). It is observed from table 3 that amylase production of *Alternaria*, *A. crassa*, *A. macrospora* and *A. tenuissima* was totally inhibited in the presence of sodium

dihydrogen orthophosphate where as disodium hydrogenorthophosphate dihydrate and diammonium phosphate stimulated the amylase production in all species of *Alternaria*. All the sources of phosphorus stimulated lipase and protease production in all *Alternaria* species.

Amylase production of *A. tenuissima* inhibited in the presence of ferrous sulphate and sodium sulphate. Calcium sulphate inhibited lipase production of *A. crassa*, Lipase activity of *A. dianthicola* inhibited in the presence of ferrous sulphate, sodium sulphate and copper sulphate where as lipase production of *A. tenuissima* was inhibited in ferrous sulphate and zinc sulphate only. Protease activity of *Alternaria alternata*, *A. citri* and *A. tenuissima* also inhibited in different sources sulphur (Table 4).

There are several reports that nitrogen and carbon sources behaves differently against lipase enzyme activity of oilseeds. Such type of work was earlier carried out by Sandikar. and Mukadam (1992), reported that stimulatory effect of different nitrogen sources for lipase production in seed-borne fungi. Rathod ,(2007) observed that disaccharides and polysaccharides stimulates lipase enzyme activity and nitrogen sources as like Calcium nitrate, caesin, gelatin and peptone also

increases lipase enzyme activity. Kesare (2009) found that nitrogen sources as like sodium nitrate, sodium nitrate, ammonium phosphate, ammonium sulphate, urea, gelatin and peptone inhibit lipase enzyme activity whereas, casein stimulates lipase enzyme activity of *Aspergillus glaucus*, *Fusarium roseum* and *Spicaria violacea* while sodium nitrate stimulates lipase enzyme activity of *Curvularia lunata*. Kakde and Chavan (2009) observed that fructose and sucrose stimulates lipase activity while lactose, carboxyl methyl cellulose and starch inhibited lipase activity. Sharma and Satyanarayana (1980) found that carbohydrate sources affects protease enzyme activity of *Helminthosporium*, *Curvularia* and *Alternaria sp.* Patil and Shastri (1982) reported that fructose and sucrose stimulated protease production in *Alternaria alternata* but glucose was found to no effect on protease activity.

It can be concluded that Carboxy methyl cellulose (CMC), ferrous sulphate, calcium sulphate, sodium sulphate, copper sulphate and sodium dihydrogen orthophosphate retarded the production of lipase, amylase and protease of some *Alternaria* species, such nature of inhibition of these may be useful to control the spoilage of seeds by fungi.

Table 1 Effect of carbohydrates on enzyme production of *Alternaria* species

Species of <i>Alternaria</i>	Carbohydrates					
	Glucose ©	Fructose	Maltose	Sucrose	CMC	Starch
Amylase Production						
<i>A. alternata</i>	15	22	18	11	10	12
<i>A. citri</i>	12	14	13	15	10	13
<i>A. crassa</i>	17	20	19	19	14	12
<i>A. dianthicola</i>	13	20	21	16	11	12
<i>A. macrospora</i>	13	12	10	18	10	12 ↓
<i>A. tenuissima</i>	16	22	14	12	10	14 ↓
Lipase Production						
<i>A. alternata</i>	18	22	24	26	25	20
<i>A. citri</i>	21	24	17	21	20	15
<i>A. crassa</i>	18	20	21	25	21	22
<i>A. dianthicola</i>	28	25	20	29	30	24
<i>A. macrospora</i>	14	27	20	23	17	21

<i>A. tenuissima</i>	21	24	27	27	26	20
Protease Production						
<i>A. alternata</i>	18	22	24	26	25	20
<i>A. citri</i>	21	24	17	21	15	20
<i>A. crassa</i>	22	20	18	16	20	19
<i>A. dianthicola</i>	18	17	15	24	20	17
<i>A. macrospora</i>	20	12	20	21	25	22
<i>A. tenuissima</i>	20	22	17	19	21	20

Activity zone in mm

Table 2 Effect of nitrogen sources on enzyme production of *Alternaria* species

Species of <i>Alternaria</i>	Sources of nitrogen							
	KNO <sub>3</sub> Control (C)	Sod Nitrate	Calcium Nitrate	Sod Nitrate	Urea	Casein	Gelatine	Peptone
Amylase Production								
<i>A. alternata</i>	10	14	14	18	17	17	18	18
<i>A. citri</i>	10	-	12	15	15	12	13	13
<i>A. crassa</i>	12	13	05	16	11	15	12	16
<i>A. dianthicola</i>	11	13	10	17	15	20	17	20
<i>A. macrospora</i>	12	17	10	18	11	14	15	15
<i>A. tenuissima</i>	08	15	15	17	14	15	15	16
Lipase Production								
<i>A. alternata</i>	21	15	20	17	20	24	20	17
<i>A. citri</i>	20	20	24	20	19	18	20	29
<i>A. crassa</i>	17	20	18	22	25	21	18	20
<i>A. dianthicola</i>	15	17	20	19	21	25	22	31
<i>A. macrospora</i>	18	17	20	18	15	27	24	20
<i>A. tenuissima</i>	20	28	21	25	24	25	25	29
Protease Production								
<i>A. alternata</i>	16	18	17	20	20	15	19	19
<i>A. citri</i>	20	14	15	18	17	20	24	20
<i>A. crassa</i>	18	19	18	14	17	19	20	15
<i>A. dianthicola</i>	20	17	17	13	20	16	22	18
<i>A. macrospora</i>	21	23	24	20	19	14	14	20
<i>A. tenuissima</i>	17	20	17	13	20	22	24	21

Activity zone in mm

Table 3 Effect of phosphorus sources on enzyme production of *Alternaria* species

Species of <i>Alternaria</i>	Sources of phosphorus			
	Potassium di Hydrogen Orthophosphate (c)	Di.sod. Hydrogen Orthopho. Dihydrate.	Sod.dihydrogen Orthosphosph Ate.	Diammonium Phosphate.
Amylase Production				
<i>A. alternata</i>	10	11	12	12
<i>A. citri</i>	10	09	10	13
<i>A. crassa</i>	08	08	-	14
<i>A. dianthicola</i>	14	15	10	10
<i>A. macrospora</i>	13	12	-	12
<i>A. tenuissima</i>	13	11	-	15
Lipase Production				
<i>A. alternata</i>	25	27	23	28
<i>A. citri</i>	25	28	26	29
<i>A. crassa</i>	25	29	27	26
<i>A. dianthicola</i>	28	25	30	25
<i>A. macrospora</i>	25	27	23	25
<i>A. tenuissima</i>	25	23	33	29
Protease Production				
<i>A. alternata</i>	18	18	19	18
<i>A. citri</i>	21	28	20	19
<i>A. crassa</i>	20	21	23	24
<i>A. dianthicola</i>	24	26	21	18
<i>A. macrospora</i>	26	23	21	20
<i>A. tenuissima</i>	20	20	20	19

Activity zone in mm

Table 4 Effect of sulphure sources on enzyme production of *Alternaria* species

Species of <i>Alternaria</i>	Sources of sulphure					
	Magnesium Sulphate (c)	Ferrous Sulphate	Sodium sulphate	Calcium sulphate	Zinc sulphate	Copper sulphate
Amylase Production						
<i>A. alternata</i>	08	12	12	10	10	13
<i>A. citri</i>	12	13	14	15	15	15
<i>A. crassa</i>	08	10	17	12	08	18
<i>A. thivla</i>	10	11	14	09	10	12
<i>A. macrospora</i>	07	12	15	14	14	14

<i>A. tenuissima</i>	11	10	10	12	12	11
Lipase Production						
<i>A. alternata</i>	20	29	27	23	25	21
<i>A. citri</i>	18	24	29	25	25	20
<i>A. crassa</i>	19	20	22	18	23	25
<i>A. dianthicola</i>	22	20	20	25	21	20
<i>A. macrospora</i>	25	27	25	28	28	22
<i>A. tenuissima</i>	24	20	28	25	20	25
Protease Production						
<i>A. alternata</i>	24	18	28	19	19	24
<i>A. citri</i>	23	20	20	20	18	18
<i>A. crassa</i>	23	28	25	29	26	20
<i>A. dianthicola</i>	20	27	20	24	22	25
<i>A. macrospora</i>	30	32	24	28	20	24
<i>A. tenuissima</i>	25	18	19	18	20	15
Activity zone in mm						

## Acknowledgement

Authors are grateful to Professor and Head Dept of Botany, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad (M.S.) for giving research facilities.

## References

- Hislop, E.C., J.L. Paver and J.P.R. Keon (1982). An acid protease produced by *Monilinia fructigena* in vitro and in infected apple fruits and its possible role in pathogenesis, *J. Gen. Microbiol.* 128: 799-807.
- Kakde R.B. and A.M. Chavan (2009). International Conference on Biological Science: Darwin to Dolly and Beyond, Nanded (M.S., India). Abstract. pp:51
- Kesare U.T. (2009) Studies on seed-mycoflora of soybean. Ph. D. Thesis, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad.
- Neergaard Paul (1973). Detection of seed borne pathogen by culture tests. *Seed Sci. and technol.* 1: 217-254.
- Patil and Shastri, N.V. (1982). Effect of carbon and nitrogen sources on extracellular production of protease by *Alternaria alternata* (Fr.) Kiessl. *Indian Jour. Microbiol.* 22(2): 115-122.
- Rajamani, S. (1990). Factors influencing the production of extracellular protease by selected phytopathogenic fungi. *J. Indian Bot. Soc.* 69: 285-287.
- Rathod S.R.(2007) Studies on seed-borne species of *Alternaria* in different crops. Ph. D. Thesis, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad.
- Sharma D.P. and T. Satyanarayana (1980). Production of protease by some moulds. *Ind. Phytopath.* 33: 359-395.
- Sandikar S.N. and Mukadam D.S. (1992). Lipase production by Seed-borne fungi. *Ind. Phytopath.* 44-6.
- Sierra, J. M. (1957) A simple method for the detection of lipolytic activity of microorganisms and some observations on the influence of the contact between cells and fatty substances. *Antonie van Leeuwenhoek Ned. Tijdschr. Hyg.* 23: 15- 25.
- Singh, B.S. and S.B. Saxena (1988). Extracellular proteolysis by certain species of *Penicillium*. *J. Indian Bot. Soc.* 67: 78-79.