



SHORT COMMUNICATION

GAMMA RAYS INDUCED MUTAGENIC STUDIES IN *CATHARANTHUS ROSEUS* (L.) G. DON., AN IMPORTANT ORNAMENTAL PLANT

S. MURUGAN, D. DHANAVEL*, S. KASTHURI

Division of Cytogenetics and Mutation Breeding, Department of Botany, Faculty of Science, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India

ABSTRACT

A study was conducted to estimate the effect of gamma rays on growth and yield traits in periwinkle (*Catharanthus roseus* (L.) G. Don), an important horticulture plant. The seeds were treated with different levels of gamma rays (20, 25 and 30). Morphological parameters were analysed on the 30th day, like plant height, days of first flower, plant height, fresh weight per plant, dry weight per plant, root fresh weight per plant and root dry weight per plant. The morphological parameters were decreased with increasing levels of gamma rays doses.

INTRODUCTION

Catharanthus roseus is an important horticultural plant belonging to the family Apocynaceae [1]. Apart from its horticultural and landscaping potential, the plant is famous for its secondary metabolite contents and medicinal uses [2]. There are numerous researches about its medicinal potential [3, 4], but the methods of increasing or exploring the genetic variations and mutation chances in this plant is scanty.

Mutation breeding is the fast growing area in all crop plants and also ornamental plants [5]. The wide chances of mutations in variety releasing and exploring new hybrids are widely studied [6-9]. Radiations like gamma can be effectively utilized as a mutation inducing agent and can alter the morphological and physiological characteristics of the plant [10,11,12]. Also the methods like chemical mutagenesis by employing chemicals like EMS also reported from crop plants [13].

The present investigation was conducted to study the effect of gamma rays in inducing mutation in *Catharanthus roseus* (L.) G. Don., an important ornamental plant.

MATERIALS AND METHODS

The dry seeds of periwinkle were collected from J. P. Laboratories, Virudhunagar District, Tamil Nadu. The

gamma ray was subjected with 05, 10, 15, 20, 25, 30, 35, 40, 45 and 50KR of CO60 at Indra Gandhi Centre for Atomic Research, Kalpakkam, Tamil Nadu, India and untreated dry seeds served as control. These treated seeds were sown on 50 seeds with spacing of 30 × 20 cm in plots. The experiment was conducted at Department of Botany, Annamalai University during the year 2016. The experiment was set up in a completely randomized design with eleven treatments in three replications, observations regarding germination percentage and seedling vigour index (SVI) were recorded on 10th day after sowing and the results were analysed statistically [14].

RESULTS AND DISCUSSION

The treatment reduced the germination percentage in periwinkle with dosage of gamma rays. This type of germination reduction was earlier reported blackgram [15], soybean [16] and rice [17]. Effect of gamma rays on growth and yield parameters in periwinkle in M1 generation was given in table 1. Packiaraj [18] reported the growth reduction in gamma rays treatments in cowpea varieties. In the present experiment, 30th day plant height, days of first flower, plant height, fresh weight per plant, dry weight per plant, root fresh weight per plant and root dry weight per plant showed a decreasing trend when compared to the control (table 1) and days to first flower increased with increase in dose of gamma rays (table 1)

Received 11 October 2017; Accepted 21 December 2017

*Corresponding Author

D. Dhanavel

Division of Cytogenetics and Mutation Breeding, Department of Botany, Faculty of Science, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India

Email: ddhana2005@yahoo.co.in

©This article is open access and licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted, use, distribution and reproduction in any medium, or format for any purpose, even commercially provided the work is properly cited. Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

Table 1: Effect of gamma rays on growth and yield parameters in periwinkle in M1 generation

Gamma rays doses	30 th day plant height	Days of first flower	Plant height	Fresh weight per plant	Dry weight per plant	Root fresh weight per plant	Root fresh weight per plant
control	8.25±0.24	41.00±1.20	80.50±2.41	158.00±4.74	47.56±2.98	23.20±0.69	8.96±0.52
20KR	6.30±0.18	47.05±1.41	70.89±2.12	150.68±4.52	43.86±2.72	16.83±0.50	8.05±0.42
25 KR	5.13±0.15	48.65±1.45	62.24±1.86	146.32±4.38	41.18±2.63	15.33±0.45	7.16±0.13
30 KR	4.20±0.12	51.05±1.53	58.76±1.76	141.77±4.25	38.29±2.52	14.13±0.42	5.45±0.29

REFERENCES

1. Nejat N, Valdiani A, Cahill D, Tan YH, Maziah M, Abiri R. Ornamental exterior versus therapeutic interior of Madagascar Periwinkle (*Catharanthus roseus*): the two faces of a versatile herb. *The Scientific World Journal*. 2015;2015.
2. Yamamoto K, Takahashi K, Mizuno H, Anegawa A, Ishizaki K, Fukaki H, Ohnishi M, Yamazaki M, Masujima T, Mimura T. Cell-specific localization of alkaloids in *Catharanthus roseus* stem tissue measured with Imaging MS and Single-cell MS. *Proceedings of the National Academy of Sciences*. 2016;113:3891-6.
3. Moreno PR, van der Heijden R, Verpoorte R. Cell and tissue cultures of *Catharanthus roseus*: a literature survey. *Plant cell, tissue and organ culture*. 1995;42:1-25.
4. Svoboda GH, Johnson IS, Gorman M, Neuss N. Current status of research on the alkaloids of *Vinca rosea* Linn. (*Catharanthus roseus* G. Don). *Journal of pharmaceutical sciences*. 1962 Aug 1;51:707-20.
5. Ahloowalia BS, Maluszynski M, Nichterlein K. Global impact of mutation-derived varieties. *Euphytica*. 2004;135:187-204.
6. Adamu, A. K. and H. Aliyu, 2007. Morphological effects of sodium azide on tomato (*Lycopersicon esculentum* Mill.). *Science World Journal*, 2: 9-12.
7. Kozgar, M. I., S. Goyal and S. Khan, 2011. EMS induced mutational variability in *Vigna radiata* and *Vigna mungo*. *Res. J. Bot.*, 6:31-37.
8. Tah, P. R., 2006. Induced macromutation in mungbean (*Vigna radiata* (L.) Wilczek). *Int. J. Bot.*, 2:219-228.
9. Khan, S. and S. Goyal, 2009. Improvement of mungbean varieties through induced mutations. *Afr. J. Plant Sci.*, 3:174-180.
10. Kiong, A., A. Ling Pick, S. H. Grace Lai and A. R. Harun, 2008. Physiological responses of orthosiphon stamineus plantlets to gamma irradiation. *Am-Eurasian J. Sustain Agric.*, 2: 135-149.
11. Arulbalachandran D, Mullainathan L. Chlorophyll and morphological mutants of black gram (*Vigna mungo* (L.) Hepper) derived by gamma rays and EMS. *Journal of Phytology*. 2009;1:236-241.
12. Pavadai P, Girija M, Dhanavel D. Effect of gamma rays on some yield parameters and protein content of soybean in M2, M3 and M4 generation. *Journal of Experimental Sciences*. 2010;1:8-11.
13. Pavadai P, Girija M, Dhanavel D. Effectiveness Efficiency and biochemical content of Physical and Chemical Mutagens in Soybean (*Glycine max* (L.) Merr.). *Journal of Phytology*. 2009;1:444-447.
14. Panse, V. G. and P. V. Sukhatme, 1985. Statistical Methods for Agricultural Work. ICAR, New Delhi.
15. Ramasamy, M. M., 1973. Investigation on induced mutagenesis in blackgram *Ph. D. Thesis*, Tamil Nadu Agric. Univ., Coimbatore.
16. Balakrishnan, P. C., 1991. Induced mutagenesis in soybean (*Glycine max* (L.) Merrill) *Ph. D. Thesis*, Tamil Nadu Agric. Univ., Coimbatore.
17. Ramesh, D. V. and T. V. V. Seetharami Reddi, 2002. Induced morphological variability among three genotypes of rice. *Cytol. Genet. J.*, 3:115-20.
18. Packiaraj, D., 1988. Studies on induced mutagenesis of parents and hybrid in cowpea (*Vigna unguiculata* (L.) Walp.). *M. Sc. (Ag.) Thesis*, Tamil Nadu Agric. Univ., Coimbatore.