



ISSN: 2075-6240

The essential oil constituents of *Artabotrys* species – A review

Subban Ravi*, Kaveri Sundaram

Department of Chemistry, Karpagam Academy of Higher Education, Coimbatore-641021, Tamil Nadu, India

Received: June 04, 2020

Revised: July 16, 2020

Accepted: July 18, 2020

Published: July 23, 2020

*Corresponding Author:

Subban Ravi,

Email: ravisubban@rediffmail.com **KEYWORDS:** *Artabotrys* species, Annonaceae, essential oil composition, ssquiterpenes, hydrodistillation

ABSTRACT

Artabotrys species which belongs to Annonaceae family are pleasant smelling and it is attributed to the presence of mono and sesquiterpenoids present in the essential oil of the plant. The objective of the present work is to review the chemical composition of the essential oils reported from twenty different *Artabotrys* species from various parts of the world. In the various *Artabotrys* species, the major compounds are monoterpene and sesquiterpene hydrocarbons and oxygenated sesquiterpenes. The frequently and most commonly identified constituents are β -caryophyllene, caryophyllene oxide, 3-Carene, cyperene, cyperenone and 1,5-epoxy-salvial4(14)-ene. Other constituents seems to be more specific to the respective *Artabotrys* species.

INTRODUCTION

Aromatic plants generally grown in subtropical and tropical countries possess essential oils. Various parts of these plants like barks, roots, rhizomes, seeds, buds, fruits flowers and leaves yield essential oils. Among the various methods available to obtain the essential oils like cold pressing, solvent extraction, hydrodistillation and supercritical fluid extraction, the preferable method is hydrodistillation [1]. Low molecular weight pleasant smelling aromatic and aliphatic compounds and terpenes/terpenoids are the major components of essential oils. Usually the essential oils will have two or three major compounds in relatively larger concentrations along with many minor compounds. The compounds present in larger amounts determine the biological activities of the essential oil [2].

The use of essential oils for medicinal treatment has reduced during the midway of the twentieth century when compared with their use in food and cosmetics industry, because of the coming up of synthetic medicinal compounds. However as a consequence of consumer's awareness about the toxic nature of the synthetic compounds, the demand for alternative and safe natural medicines has risen. As a result recently essential oils have gained more attention as a possible alternative and in the modern era they also finds their use in various products such as air fresheners, cosmetics, hygiene products, household cleaning products, food, and agriculture as well as in medical uses. Further they are used in aromapathay and in other para-medical practices. Essential oils possess various bioactivities like antimicrobial, anti-inflammatory, antioxidant, antiviral and

ant carcinogenic. Consequently in an attempt to find alternative medicines, the pharmacological activities evaluation studies of the essential oils have become active in recent years [3].

Artabotrys species belongs to the Tropical Annonaceae family and comprises of about 105 species [4]. About 70% of the species are present in Asia and the remaining 30% of the species are found in Africa. Among the climbing climbers of Annonaceae family *Artabotrys* species are unique woody climbers. The uniqueness is due to the presence of special persistent flower hooks which helps them in climbing [5]. Leaves are simple and alternate, without hairs. The inflorescence often bear many flowers, but are single flowers in some species. To the best of our knowledge for 20 plants the composition of essential oils (Table 1) from the different parts of the plants have been investigated and reported in the literature. They are *A. hexapetalus*, *A. rupestris*, *A. ordratissimus*, *A. modestus*, *A. hongkongensis*, *A. vinhensis*, *A. lastoursvillensis*, *A. pallens*, *A. petelotii*, *A. intermedius*, *A. harmanddii*, *A. taynguyenensis*, *A. insignis*, *A. pierianus*, *A. rutilus*, *A. thomsoni*, *A. jollyanus*, *A. oliganthus*, *A. velutinus* and *A. venustus*. Qualitative and Quantitative variations in chemical composition of the essential oils obtained from *Artabotrys* species cultivated in India and elsewhere. This may be ascribed to factors such as the age and nature of the plant, climatic conditions, and place of collection and handling procedures.

Essential oils from *Artabotrys* species

A. hexapetalus, commonly known in India as manoranjini is a shrub and its flowers are yellow in colour and have renowned

Copyright: © The authors. This article is open access and licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted, use, distribution and reproduction in any medium, or format for any purpose, even commercially provided the work is properly cited. Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

Table 1: List of *Artabotrys* species for which the composition of the essential oils were determined by GC-MS

Plant Name	Plant part/method of extraction	Place	Volatile Compounds identified	Ref
<i>A. hexapetalus</i>	Flower/simple headspace-solvent trapping technique	Thailand	Ethyl acetate (47.3%), isobutyl acetate (26.8%), ethyl isobutanoate (9.2%) and ethyl butanoate (9.7%)	[6]
	Flower/Hydrodistillation	Thailand	β -gurjunene (30.0%), β -caryophyllene (10.1%), and globulol (13.8%)	[6]
	Flower/solvent extraction	Thailand	Isopentyl acetate (12.6%), 3-methylbutanol (5.7%), limonene (5.7%), linalool (7.7%), benzene ethanol (5.3%), 2-methylbutanol (3.9%), ethyl butanoate (3.1%), 2-methylbutyl acetate (7.7%)	[6]
	Flower/solid-phase microextraction	Thailand	Isobutyl acetate (39.5%), Ethyl acetate (12.8%), ethyl isobutanoate (4.9%), ethyl 2-methyl butanoate (2.3%), ethyl isovalerate (11.3%), ethyl 3-methyl-2-butenate (6.8%), isobutyl isovalerate (2.1%) and β caryophyllene (5.4%)	[6]
<i>A. hexapetalus</i>	Flower/hydrodistillation	Vietnam	Caryophyllene oxide (31.5%) and β -caryophyllene (11.4%)	[10]
<i>A. hexapetalus</i>	Stem bark/ Hydrodistillation	Tanzania	Elemene (5.57%), β -caryophyllene oxide (45.62%), 1,3,3-trimethyl-2-(2-methylcyclopropyl)-cyclohexene (8.62%), 4 α ,8dimethyl-2-(1-methylethylidene)naphthalene (7.97%)	[11]
<i>A. hexapetalus</i>	Leaves/ Hydrodistillation	Tanzania	cadinol (10.89%), β caryophyllene oxide 11hexadecynal (8.41%),	[11]
	Stem bark,	Vietnam	Caryophyllene oxide (36.2%), α -copaene (10.9%), β -caryophyllene (7.8%) and β -asarone (6.1%)	[12]
<i>A. rupestris</i>	Root/ Hydrodistillation	Tanzania	First two fractions:6-(p-tolyl)-2-methyl-2-heptanol (44.49% and 42.25%, respectively), 7,11,15-trimethyl-3-methylene-hexadeca1,6,10,14-tetraene (25.36%) Third fraction: (-)-spathulenol. (43.6%) 1-(1,5-dimethyl-4hexenyl)-4-methyl-benzene (22.11%)	[11]
<i>A. ordratissimus</i>	Leaves/ Hydrodistillation	Bangalore	3-Carene (44.91%), β -caryophyllene (19.17%), α humulene (8.78%), α -copaene (6.59%) and caryophyllene oxide (5.55%)	[13]
<i>A. ordratissimus</i>	Fruit & leaves/ Hydrodistillation	South India	β -caryophyllene (14.7 %), trans nerolidol (8.2%, δ -cadinene (7.3%), α -copaene (6.4%), trans- α -farnesene (5.8%), τ -cadinol (4.3%) and caryophyllene oxide (3.2%)	[14]
<i>A. ordratissimus</i>	Leaves/ Hydrodistillation	South India	β -caryophyllene (17.3%), trans nerolidol (1.9%), δ -cadinene (4.2%), α -copaene (9.3%), trans, trans- α -farnesene (7.4%), τ -cadinol (2.9% and caryophyllene oxide (6.8%)	[14]
<i>A. ordratissimus</i>	Leaves/ Hydrodistillation	India, Calicut	Terpinen-4-ol (38.6%), β Caryophyllene oxide (11.8%)	[15]
<i>A. modestus</i>	Stem bark/ Hydrodistillation	Tanzania	Linalool (11.2%) (20.5% sesquiterpenes)	[16]
<i>A. hongkongensis</i>	Leaves/ Hydrodistillation	Vietnam	cadinol, cubenol, β -Caryophyllene oxide, (-) spathulenol and 7-methyl-4-methylene-1-(1methylethyl)-naphthalene.	[17]
<i>A. vinhensis</i>	Leaves/ Hydrodistillation	Vietnam	Spathulenol (13.1%), cadinene (6.3%), β -caryophyllene (6.6%), δ - γ -elemene (6.3%) and benzyl benzoate (5.6%).	[18]
<i>A. lastoursvillensis</i>	Stem bark/Hydrodistillation	Gabon	Germacrene D (14.4%), α -pinene (16.7%), limonene (15.4%), benzyl benzoate (8.8%) and β -pinene (7.5%).	[8]
<i>A. pallens</i>	Leaf/ Hydrodistillation	Vietnam	Cyperene (25.9%), cyperenone (11.1%), caryophyllene oxide (6.7%), P-elemene (5.5%) and P-guaiene (5.0%), cadalene (5.6%)	[19]
<i>Artabotrys petelotii</i>	Leaf/hydrodistillation	Vietnam	α -phellandrene (20.1%), α -gurjunene (21.9%) and bicycloelemene (9.6%), with significant proportions of α -pinene (5.2%), (E)- β -ocimene (4.4%) and isolekene (3.6%)	[20]
<i>A. intermedius</i>	Stem bark/ hydrodistillation	Vietnam	Elemol (19.4 %), cis- β -guaiene (9.2 %), δ -elemene (5.5 %) and δ -cadinene (8.4%)	[20]
	Leaf/ hydrodistillation	Vietnam	Elemol (32.7 %), δ -cadinene (11.9 %) and spathulenol (9.6 %)	[20]
<i>A. harmanddii</i>	Stem bark/ hydrodistillation	Vietnam	δ -3-carene (19.1 %), α -gurjunene (10.7 %) and α -zingiberene (6.3 %)	[20]
	Leaf/ hydrodistillation	Vietnam	δ -3-carene (29.9 %), germacrene D (15.1 %) and α -amorphene (8.0 %)	[20]
<i>A. taynguyenensis</i>	Leaf/ hydrodistillation	Vietnam	spathulenol (17.4 %), aromadendrene epoxide (12.2 %), γ -elemene (7.1 %) and isospathulenol (5.6 %)	[20]
	Leaves/ hydrodistillation	Vietnam	valencene (40.1%) along with δ -selinene (8.8%), α -pinene (6.7%), α -muurolene (5.1%) and α -panasinsene (5.1%)	[21]
<i>A. insignis</i>	Stem bark/ hydrodistillation	Zaire	bicycloelemene (25.1%), bicyclogermacrene (23.7%) and spathulenol (12.8%)	[21]
	Root bark/ hydrodistillation	Zaire	Caryophyllene oxide (31.4%), 6-cadinol (10.7%), caryophyllenol (8.2%), 1,5-epoxy-salvial4(14)-ene (8.1%), camphene (9.4%), p-elemene (2.4%), cyperenone(4.4%)	[22]
<i>A. pierianus</i>	Stem bark/ hydrodistillation	Zaire	camphene(5.1%),1,5-epoxy-salvial4(14)-ene (27.3%), widdrol (7.5%), spathulenol (4.7%), 6-cadinol(4.3%),linalool (3.4%), cyperenone (3.2%)	[22]
	Stem bark/ hydrodistillation	Zaire	Cyperene (13.6%), caryophyllene oxide (9.4%), cadalene(6.4%), cyperenone (13.5%)	[22]
<i>A. rufus</i>	Root bark/ hydrodistillation	Zaire	a-copaen-1 1 -0l (6.3%), p-caryophy llene (20.5%), caryophyllene oxide (22.6%), 1,5-epoxy-salvial4(14)-ene (12.7%), caryophyllenol (5.2%)	[22]
<i>A. thomsoni</i>	Stem bark/ hydrodistillation	Zaire	Camphene (9.6%), cyperene (3.4%), palustrol (7.9%), spathulenol (6.9%), 1,5-epoxy-salvial4(14)-ene (12.1%), 6-cadinol (4.7%), cyperenone (7.7%)	[22]
<i>A. venustus</i>	Stem bark/ hydrodistillation	Malaysia	Camphene (10.1%), ledol (7.3%), 1,5-epoxy-salvial4(14)-ene (34.5%), β -caryophy llene (3.8%), caryophyllene oxide (3.9%)	[22]

(Contd...)

Table 1: (Continued)

Plant Name	Plant part/method of extraction	Place	Volatile Compounds identified	Ref
<i>A. jollyanus</i>	Leaves/ hydrodistillation	southern Côte d'Ivoire, Brazil	Trans-calamenene (15.7%), -copaene (14.8%), -cubebene (10.4%), cadina-3,5-diene (10.3%), (E)- caryophyllene (6.3%) and cadina-1,4-diene (6.1%).	[22]
<i>A. oliganthus</i>	Leaf/ hydrodistillation	Brazil	Δ -3-carene (60.2 %), myrcene (10.6 %), β -selinene (1.9 %). β -pinene (5.9 %), α -pinene (4.6 %) and γ -terpinene (4.6 3 %)	[23]
	Stem bark oil/ hydrodistillation	Brazil	Δ -3-carene (57.2 %), β -selinene (5.7 %), myrcene (2.8 %). β -pinene (7.7 %), α -pinene (6.1 %) and γ -terpinene (5.3 %)	[23]
	Fruit/ hydrodistillation	Brazil	Myrcene (59.2 %), Δ -3-carene: 16.3 %, β -farnesene (3.7 %) and β -selinene (2.2 %)	[23]
<i>A. velutinus</i>	leaves	Bonou	benzyl benzoate (61.2%) and E- β -caryophyllene (9.1%).	[9]

Table 2: Classification of *Artabotrys* species based on their major compounds

S. No.	Major compounds	Artabotrys species
1	β -caryophyllene and caryophyllene oxide	<i>A. hexapetalus</i> (Vietnam) flower <i>A. hexapetalus</i> (Tanzania), stem bark, leaves <i>A. ordratissimus</i> (South India), leaves & Fruits <i>A. modestus</i> (Juma), stem bark
2	6-(p-tolyl)-2-methyl-2-heptanol	<i>A. rupestris</i> (Tanzania), root
3	(-)-spathulenol	<i>A. rupestris</i> (Tanzania), root <i>A. hongkongensis</i> (Vietnam) <i>A. harmandii</i> (Vietnam), leaves & stem bark <i>A. taynguyenensis</i> root & stem bark
4	3-Carene	<i>A. ordratissimus</i> (India-Bangalore), leaves <i>A. intermedius</i> (Vietnam), leaf and stem bark <i>A. oliganthus</i> (Brazil), leaves, stem bark, fruit
5	Terpinen-4-ol	<i>A. ordratissimus</i> (India-Calicut), leaves
6	cyperene and cyperenone	<i>A. lastoursvillensis</i> (Gabon) <i>A. insignis</i> (Zaire), root, stem bark <i>A. pierianus</i> (Zaire), stem bark <i>A. thomsoni</i> (Zaire), root, stem bark
7	α -gurjunene	<i>A. hexapetalus</i> (Thailand) <i>A. pallens</i> (Vietnam), leaves
8	elemol	<i>A. petelotii</i> (Vietnam), leaf
9	trans-calamenene	<i>A. jollyanus</i> (Brazil) leaves
10	benzyl benzoate	<i>A. velutinus</i> (Bonou) leaves
11	α -pinene	<i>A. vinhensis</i>
12	1,5-epoxy-salvia14(14)-ene	<i>A. rufus</i> (Zaire), root bark <i>A. venustus</i> (Malaysia) stem bark <i>A. thomsoni</i> <i>A. insignis</i> (Zaire), root, stem bark

exotic fragrance. In the beginning the flowers are green in colour and on ageing it turns yellow and are long lasting with pleasant fruity smell. When young it is a shrub and after attaining a height of 2 meters it turns into a climber. Flowers and the essential oil possess antifungal activity and are used in the perfume industry, for preparing tea like beverage, in aromapathy and even it is used as a substitute for Ylang Ylang oil.

The essential oil was extracted from the flowers, leaves and stem bark of *A. hexapetalus* from countries like Thailand, Vietnam and Tanzania. The volatile constituents are dominated by sesquiterpene hydrocarbons and oxygenated sesquiterpenoids and mainly consists of β -caryophyllene oxide and or β -caryophyllene as the common constituent. An attempt was made to match the odour of the flowers from *A. hexapetalus* [6] from Thailand for which they extracted the

volatile constituents by four different methods. Seventeen components were obtained from Simple headspace-solvent trapping technique and they are dominated by ethyl and isobutyl esters, 47.3% ethyl acetate, 9.2% ethyl isobutanoate, 26.8% isobutyl acetate and 9.7% ethyl butanoate. By hydrodistillation method thirty one components were identified of which β -caryophyllene, β -gurjunene and globulol (10.1, 30.0 and 13.8% respectively) are the major compounds. By solvent extraction method thirty one components were identified comprising of benzene ethanol (5.3%), limonene (5.7%), linalool (7.7%), 3-methylbutanol (5.7%), isopentyl acetate (12.6%) and 2-methylbutyl acetate (7.7%),) are the major compounds. By solid phase micro extraction method, thirty-nine components were identified with a chemical composition of 12.8% Ethyl acetate, 4.9% ethyl isobutanoate, 39.5% isobutyl acetate, 2.3% ethyl 2-methyl butanoate, 11.3% ethyl isovalerate, 6.8% of ethyl 3-methyl-2-butenate, 2.1% isobutyl isovalerate and 5.4% β caryophyllene (5.4%). The odour of the solution from this technique was similar to that of the fresh flower.

A. odoratissimus is also a large woody climber originated in India, Philippines, South China and Burma (Myanmar). Its flowers are either solitary, or in clusters of two or three, axillary and turns greenish yellow in color on ageing. Its name in Bengali is 'Kanthali champa' (jackfruit-champa) since it gives a strong smell resembling that of ripened jackfruit. The flowers are available throughout the year. Because of its huge size it is not suitable for small gardens. It is used as an anti-depressant, mood elevator, hypotensive and as a hair vitalizer. The oil is also extensively used in deodorants, perfumes, shampoos, beauty soaps, hair and skin lotions and creams, hair oils etc., in blends or alone. Essential oil constituents were reported from the leaves and fruits of *A. odoratissimus* from three different locations Tamilnadu, Calicut and Bangalore in India. The major compounds identified are 3-Carene (44.91%), β -caryophyllene (17.3%) and Terpinen-4-ol (38.6%) from the essential oils from Tamilnadu, Calicut and Bangalore respectively. β -caryophyllene and caryophyllene oxide are the two other common compounds present from these three essential oils.

A. rupestris from Tanzania was collected and three different fractions of the essential oil was obtained at interval of 50 min using hydrodistillation method. In the first two fractions 6-(p-tolyl)-2methyl-2-heptanol was the major component (44.49% and 42.25%, respectively) and in the third fraction spathulenol (43.60%) was the major compound. It also contains

a high amount of alcoholic and unsaturated sesquiterpenes. Essential oil from an endemic species *A. modestus*, from Tanzania consists of cadinol, β -Caryophyllene oxide, cubenol, 7-methyl-4-methylene-1-(1-methylethyl)-naphthalene, (-)-spathulenol and it was commercially exploited for its mosquito repellence activity. Spathulenol (13.1%), δ -cadinene (6.3%), benzyl benzoate (5.6%), β -caryophyllene (6.6%) and α -elemene (6.3%) were reported from *A. hongkongensis*. Germacrene D (14.4%), α -pinene (16.7%), limonene (15.4%), benzyl benzoate (8.8%) and β -pinene (7.5%) were found to be richer in the oil of *A. vinhensis*. The essential oil from the leaves of Ivoirian *A. jollyanus* consists largely of sesquiterpene hydrocarbons like trans-calamenene, copaene, cubebene and 7-hydroxycalamenene. When the composition of the essential oil was analysed during its phenological cycle, the same sesquiterpenes remains as the dominant compounds but the quantity of the component varies. Another Ivoirian species *A. oliganthus* differs from the essential oil from the leaves of *A. jollyanus* by dominating in monoterpenes like 3-carene, 60.2% and myrcene, 10.6% [7]. In turn, these two oils vary considerably from the oils obtained from the bark of Gabonese *A. lastourvillensis* Pell consisting of cyperene, 25.9% and cyperenone, 11.1% as the major compounds [8] and Beninese *A. velutinus*. [9] which is richer in aromatic components (62.6%) and sesquiterpene hydrocarbons (29.9%). It consists of 30 compounds representing 98.9% of the oil and the major components are benzyl benzoate (61.2%) and E- β -caryophyllene (9.1%).

The oil of *A. pallens* comprised of 39.8% monoterpenes and 58.0% sesquiterpenoid compounds with significant proportions of α -phellandrene (20.1%), α -gurjunene (21.9%), bicycloelemene (9.6%), α -pinene (5.2%), (E)- β -ocimene (4.4%) and isodenedene (3.6%). The oils of *A. intermedius* and *A. oliganthus* are similar with respect to having a high content of δ -3-carene (19.1-29.9% and 57.2-60.2% respectively) but differ due to its low content of myrcene (0.7-0.8% and 15.7-59.2% respectively). One of the chemotypes of *A. ordratissimus* also consists of 3-carene (44.91%), as the major compound. From *A. harmandii* forty-one compounds comprising of seven monoterpene hydrocarbons (6.5%), five oxygenated monoterpenes (2.8%), nineteen sesquiterpene hydrocarbons (35.8%), nine oxygenated sesquiterpenes (44.1%) and a non-terpene (3.6%) was reported in the leaf oil. The major compounds are aromadendrene epoxide (12.2%), spathulenol (17.4%), γ -elemene (7.1%) and isospathulenol (5.6%).

Classification of Essential Oils

Based on the above data the following chemical patterns of essential oils from *Artabotrys* oils may be arrived at.

1. Oils mainly with sesquiterpenes,
2. Oil dominated by monoterpenes
3. Oils consisting mainly of monoterpenes and sesquiterpenes.
4. Oils with 1,5-epoxysalvial-4-(14)-ene as the main compound.
5. Oils with cyperene and cyperenone as the main compounds

Oils dominated by sesquiterpenes are from *A. hexapetalus*, *A. odoratissimus*, *A. lastoursvillensi*, *A. insignis*, *A. pierreanus*,

A. rufus, *A. thomsoni*, *A. venustus*, *A. hongkongensis*, leaf and stem oils of *A. petelotti* and leaf oil of *A. harmandii*.

Oils rich in both monoterpenes and sesquiterpenes are from *A. vinhensis*, the leaf and stem oils of *A. intermedius*. Oils consisting of 1,5-epoxysalvial-4-(14)-ene is in a significant amount are from *A. pierianus*, *A. thomsoni*, and *A. venustus*. Oils with considerable quantity of cyperene and cyperenone are from the bark oil of *A. lastourvillensis* from Gabon (cyperene (25.9%) and cyperenone (11.1%) and leaf and stem bark oils from *A. insignis*, *A. rtabotrys pierianus*, *A. thomsoni*, and *A. venustus*.

The chemical patterns of both the leaf and stem oils of *A. petelotti* are similar to first chemical class with a dominance of sesquiterpene compounds. Also, could be classified into the third class due to the abundance of mono- and sesquiterpene as the major compounds. The leaf oil of *A. taynguyenensis* Ban was richer the sesquiterpenoid valencene (40.1%) along with α -pinene (6.7%), δ -selinene (8.8%), α -panasinsene (5.1%) and α -muurolene (5.1%). However, bicycloelemene (25.1%), spathulenol (12.8%) and bicyclogermacrene (23.7%) are the principal compounds in the stem oil. Further based on the presence of the major compound responsible for the activity of *Artabotrys* species can be classified as in table-2. If sufficient number of data are available then it can be narrowed down.

CONCLUSIONS

We have presented an overview of all the published reports on the chemical composition of the essential oils from twenty different *Artabotrys* species. The essential oils from *Artabotrys* species have been extensively studied during the last decades, because they are richer in chemical compounds of interest in the perfume and food industries. We have restricted to the main constituents identified in the oils. The study will be useful to the fragrance, cosmetics and food industry to monitor quality and consumer safety by developing species biomarkers. In addition traceability, adulterations of drugs and foods and other quality control aspects could be better addressed by using the present work.

AUTHOR'S CONTRIBUTIONS

Both the authors contributed equally in collecting the data, validating and in preparing the manuscript. The first author SR also participated in the review of the manuscript.

REFERENCES

1. Hamdy AE, Shaaban Ahmed H, Ghorab EL, Takayuki Shibamoto. Bioactivity of essential oils and their volatile aroma components: Review. *Journal of Essential Oil Research*. 2012; 24:2, 203-212.
2. Marianne Perricone, Ersilla Arace, Maria R. Corbo, Milena Sinigaglia, and Antonio Bevilacqua. Bioactivity of Essential Oils: A review on their interaction with food components. *Frontiers in Biology*. 2015; 6:76,1-12
3. Baber Ali, Nasar Ali Al Wabel, Saiba Shams, Aftab Ahamad, Shah Alam Khan, Firoz Anwar. Essential Oil used in Aromapathy: A systemic Review. *Asian Pacific Journal of Tropical Biomedicine*. 2015; 5(8): 601-611.
4. Guo X, Tang CC, Thomas DC, Couvreur TLP, Saunders RMK. A

- mega-phylogeny of the Annonaceae: taxonomic placement of five enigmatic genera and support for a new tribe, Phoeniciantheae. *Scientific Reports* 7: 2017: 7323. <https://doi.org/10.1038/s41598-017-07252-2>.
5. Mabberley D. Mabberley's plant-book: a portable dictionary of plants, their classifications, and uses. Cambridge University Press, Cambridge. 2008: 1021.
 6. Mahidol C, Chimneys N, Chokchaichamnankit D, Techasakul S. Identification of volatile constituents in *Artabotrys hexapetalus* flowers using simple headspace solvent-trapping technique in combination with Gas Chromatography-Mass Spectrometry and Retention Indices. *Acta Hort.*. ISHS 2005: 677.
 7. Ouattara ZA, Brice Boti J, Ahibo AC, Félix Tomi, Bighelli A. *Artabotrys oliganthus* Engl. & Diels from Ivory Coast: Composition of Leaf, Stem Bark and Fruit Oils. *Journal of Essential Oil Bearing Plants*. 2015: 14(1): 95-100
 8. Menut CG, Lamaty JM, BessiL.re CE, Mve-Mb, Affane-Nguema JP. Aromatic plants of Tropical Central Africa. W. The essential oil of *Artabotrys lastourutlensis* Pell. from Gabon. *J. Essent. Oil Res.* 1992: 4: 305- 307.
 9. Yovo M, Alitonou GA, Yedomonhan H, Tchobo F, Dedome O, Sessou P, Avlessi F, Menut C, Sohounhloué D. First report on chemical composition and antimicrobial activity of *Artabotrys velutinus* extracts against some clinical strains in Benin. *American Journal of Applied Chemistry*. 2016: 4 (3), 71-76. doi: 10.11648/j.ajac.20160403.11
 10. Phan GM, Phan ST, König WA. Chemical composition of the flower essential oil of *Artabotrys hexapetalus* (L. f.) Bhandare of Vietnam, *Journal of Essential Oil Research*, 2007: 19:6, 523-524. DOI: 10.1080/10412905.2007.9699321
 11. Suleiman RA., Mgani QA and Nyandoro SS. Chemical compositions and mosquito repellency of essential oils from *Artabotrys hexapetalus* and *Artabotrys rupestris*. *Int. J. Biol. Chem. Sci.* 2014: 8(6): 2804-2812.
 12. Rang DD, Huong HH. The primary study of the chemical composition of the *Artabotrys hexapetalus* flower in Ninh Binh Province. *Journal of Science University of Hanoi, Vietnam*. 2004: 4: 70-73.
 13. Venkatesh HN., Sudharshana TN, Manjunath BNK, Mohana DC. Antifungal and antifumonisin activities of chemically characterized essential oil of *Artabotrys odoratissimus*. *Journal of Herbs, Spices & Medicinal Plants*. 2018. DOI: 10.1080/10496475.2018.1520776
 14. Jirovetz L, Buchbauer G, Puschmann CH, leischhacker W. .Analysis of the essential oils of *Artabotrys odoratissimus* fruits and leaves from South-India. *Journal of Essential Oil Bearing Plants*. 1998:1 : 2-3.
 15. Garg S., Siddiqui N. Chemical composition of the essential oil of *Artabotrys odoratissimus*. *Indian Journal of Chemistry Section B*. 1999: 38(10):1229-1230
 16. Juma SR. Chemical composition and biomedical properties of Annonaceous and other essential oils. PhD. Thesis, 2009. University of Dar es Salaam, Tanzania.
 17. Thang TD, Dai DN, Hoi TM, Ogunwande AI. Essential Oils from Five Species of Annonaceae from Vietnam, *Natural Product Communications*. 2013: 8(2): 239-242.
 18. Dai DN, Thang TD, Hoi TM, Dung NX. Chemical constituents of essential oil from *Artabotrys vinhensis* of Vietnam. *Journal of Science and Technology, Vietnam*. 2010: 48:946-950.
 19. Thang TD, Dai DN, Hoi TM, Ogunwande IA. Chemical compositions of the leaf essential oils of some Annonaceae from Vietnam. *Journal of Essential oil Research*. 2013: 25(2): 85-91
 20. Nguyen H, Hung, DN, Dai, Doan M, Dung, Truong TB, Tran DG, Thang, Ogunwande IA. Chemical Composition of Essential Oils of *Artabotrys petelotii* Merr., *Artabotrys intermedius* Hassk., and *Artabotrys harmandii* Finet & Gagnep. (Annonaceae) from Vietnam. *Journal of Essential Oil Bearing Plants*. 2014: 17(6): 1105-1111.
 21. Thang TD, Dai N, Ogunwande IA. Study on the chemical constituents of essential oils of two Annonaceae plants from Vietnam: *Millettia sinensis* and *Artabotrys taynguyenensis*, 2014.
 22. Abbas GF, Roblot HF, Leboeuf, André Cavé M, Brigitte Charles. Essential Oils of Annonaceae. Part VI. Volatile Constituents of the Essential Oils from Five *Artabotrys* Species. *Journal of Essential oil Research*. 1997: 9 (2): 145-149.
 23. Gooré SG, Ouattara ZA, Yapi AP, Yves-AlainBékro, Bighelli A, Mathieu Paoli, FelixTomi. Chemical composition of the leaf oil of *Artabotrys jollyanus* from Côté d'Ivoire. *Brazilian Journal of Pharmacognosy*. 2017: 27 : 414–418.