

SHORT COMMUNICATION

Effect of physical factors on development of anthracnose of mango fruits

Gulab M. Rathod

Department of Botany, Shrikrishna Mahavidyalaya, Gunjoti, Dist.Osmanabad (M. S.), India

CORRESPONDENCE

Gulab M. Rathod, Department of Botany,
Shrikrishna Mahavidyalaya, Gunjoti, Dist.
Osmanabad (M. S.), India

E-mail:

EDITOR

Gadgile D.P.

CB Volume 2, Year 2011, Pages 15-16

Anthracnose is most important post harvest fungal disease of mango. It is the major disease, limiting fruit production in all countries where mangoes are grown, especially where high humidity prevails during the cropping season (Gadgile et al., 2009). Environmental factors such as temperature and Relative Humidity (R.H.) play vital role in the development and spread of post-harvest fungal diseases of fruits. (Bagwan and Yeole, 2003; Bagwan and Meshram; 2003; Cherian and Mani, 2007, Gadgile et al, 2009, Gadgile et al, 2009a, Gadgile et al, 2009b, Gadgile and Chavan, 2010). Severity of post-harvest fungal diseases depends upon environmental factors like temperature and relative humidity. Present paper deals with the study of impact of environmental factors on development of anthracnose of mango fruits. Mango fruits Alphonso variety was collected from Osmanabad fruits market, India and surface sterilized with 0.1 % HgCl₂. Two mm sized injury was made. That fruits were

dipped in spore suspension of *Colletotrichum gloeosporioides* for 2 minutes and placed in sterilized polythene bags as on fruit per bags. It was incubated to different level of temperature and R.H. percentages adjusted level were maintained (Buxton and Mellanby, 1934). On 8th day of incubation severity of rot was recorded as percent fruit area infected.

At 25°C and at 100% R.H. anthracnose severity was highest. Disease development was absent at 10°C and at 30% R.H., so at this physical environment there is a very less rotting of mango fruits. Percentage of severity was increased from 30 to 100% R.H. (Table 1). Gadgile et al. (2009b) reported similar findings.

It can be concluded at 10°C and at 30% R.H.; anthracnose is less severe in mango fruits. Whereas, at 25°C and at 100% R.H. anthracnose development is maximum.

Table 1 Effect of temperature and R.H. on disease severity anthracnose of mango fruits

Temperature (°C)	Disease development %	R.H. (%)	Disease development %
10	0.0	30	19.5
25	63.0	50	30.4
30	52.5	80	50.9
40	33.4	100	61.3

Acknowledgement

Thanks are due to UGC, (WRO, Pune) New Delhi, India for financial support in the form of minor research project. Author is also grateful to Principal, Shrikrishna Mahavidyalaya, Gunjoti, Dist.Osmanabad (M. S.), India for providing research facilities.

References

- Bagwan ,N.B. and Yeole, R.D. (2003). Effect of temperature and relative humidity on incidence of post- harvest rot of mango. J.Mycol.Pl.Pathol. 33(1):136- 137.
- Bagwan, N.B. and Meshram, B.M. (2003). Effect of temperature and relative humidity of on post- harvest rots of banana fruits. J.Mycol.Pl.Pathol. 33(2): 299-300.
- Buxton, P.A. and Mellanby, K. (1934). The measurement and control of humidity. Bull.Ent.Res. 25:171-175.

- Cherian, T.T. and Mani Varghese, K.I. (2007). Effect of temperature , relative humidity and injury on development and spread of post- harvest rot of banana. Indian Phytopath. 60(1):112-114.
- Gadgile, D.P., Chavan, A.M., Pangrikar, P.P., Hatti, A.D. and Kakde, R.B. (2009) .Impact of Temperature and Relative Humidity on development of *Rhizopus* rot of mango fruits. Nat. J.of Life sci. 6(2): 215-217.
- Gadgile, D.P., Chavan, A. M. Kakde, R.B., Hatti, A.D., Pangrikar, P.P. and Gaikwad, R.S. (2009a). Development of *Botryodiplodia theobromae* Rot (Stem End Rot) of Mango Fruits Under The Influence of Different Temperature And Relative Humidity. The Ecotech. 1(2):184-186.
- Gadgile, D.P., Chavan, A. M. Kakde, R.B., Hatti, A.D., Pangrikar, P.P. and Gaikwad, R.S. (2009b). Influence of

Environmental factors on the development of *Colletotrichum gloeosporioides* rot (Anthracnose) of Mango fruits. Environmental Conservation J. 10 (3):73-75.

Gadgile, D.P. and Ashok M. Chavan. (2010). Impact of temperature and relative humidity on development of *Aspergillus flavus* rot of mango fruit. Recent Research in Science and Technology. 2(3): 48-49.