



REGULAR ARTICLE

HETEROTIC EXPRESSION IN INBREDS DERIVED FROM FOUR DIFFERENT BASE POPULATIONS IN MAIZE (*ZEA MAYS* L.)

B. BHARATHIVEERAMANI¹, M. PRAKASH^{2*}, A. SEETHARAM³

¹Foliage Crop Solutions Private Limited, R and D Farm, Attur, Tamilnadu 636102, India

²Department of Genetics and Plant Breeding, Faculty of Agriculture, Annamalai University, Annamalai Nagar 608002, Tamilnadu, India

³Former Project Coordinator (Small millets), Indian Council of Agricultural Research, GKVK campus, Bangalore 560065, India

ABSTRACT

The concepts of combining ability and heterosis are the fundamental tools for enhancing productivity of different crops. The main objective is to study the usefulness of inbreds derived from four different base populations namely advanced generations of single cross hybrids, three way cross hybrids, hybrid mixtures and composites by analyzing the combining ability of inbreds and crosses derived and the heterosis obtained for important characters. One hundred and forty four maize inbreds derived from four different base populations namely advanced generations of single cross hybrids, three way cross hybrids, hybrid mixtures and composites were studied for their heterotic expression. The results indicated that composite and hybrid mixture base populations shall be of great use in deriving genetically divergent inbreds and single cross hybrids with significant standard heterosis suitable for commercial exploitation.

Keywords: Maize, Heterosis, Inbreds

INTRODUCTION

Maize is an extensively investigated crop for combining ability and heterosis breeding. After realizing the advantages of single cross hybrids in maize improvement, the thrust at present has been on this direction. With this reorientation towards breeding of single cross hybrids, it has become imperative now to use diverse source populations for deriving inbreds not only divergent but also heterotic and productive. It is well recognized that the crosses between genetically diverse parents show greater heterosis compared to crosses between closely related parents. The superiority of inbreds directly depend on the presence of desirable genes and gene complexes in the base population [1]. Populations of narrow genetic base have been considered as preferential germplasms in breeding programs compared to open pollinated varieties, since the latter are little improved. Besides single cross hybrids, elite line synthetics/composites, F₂ populations, backcross populations, pools and experimental varieties are also used as source materials [2]. Arshad *et al.* [3] suggested that, for successful breeding program, successful selection of superior genotypes is essential. Usually the breeders select genetically narrow-base types for developing recombination lines from F₂ of commercial single cross hybrids for maize [4]. Though widely followed

by maize breeders, the study about the genetic divergence and usefulness of inbreds derived from such narrow-base populations are very limited.

In the development of a desirable hybrid, it is necessary to identify the potential inbred lines which have high combining ability for the characters under consideration in hybrid combinations [5]. The main objective of the present investigation is to assess the combining ability of inbreds and crosses derived and the heterosis obtained for important characters of inbreds derived from four different base populations namely advanced generations of single cross hybrids, three way cross hybrids, hybrid mixtures and composites.

MATERIALS AND METHODS

The material for the study comprised of 144 inbreds originated from different base populations of unknown pedigree *viz.* advanced generations of single cross hybrids, three way cross hybrids, hybrid mixtures and composites. The number of inbreds representing different base populations and their accession number is given in table 1. The experiment was laid in a 12 x 12 simple lattice design with two replications at the RandD Farm, Foliage Crop Solutions Private Limited, Attur, TamilNadu, India. The data were recorded on 19 characters *viz.*, days to 50%

Received 14 March 2018; Accepted 11 April 2018

*Corresponding Author

M. Prakash

Professor, Department of Genetics and Plant Breeding, Faculty of Agriculture, Annamalai University, Annamalai Nagar 608002, Tamilnadu, India

Email: geeth_prakash@yahoo.co.in

©This article is open access and licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted, use, distribution and reproduction in any medium, or format for any purpose, even commercially provided the work is properly cited. Attribution – You must give appropriate credit, provide a link to the license, and indicate if changes were made.

tasseling, days to 50% silking, anthesis-silking interval, plant height (cm), ear height (cm), number of leaves, leaf length (cm), leaf width (cm), tassel length (cm), number of tassel branches, ear length (cm), ear circumference (cm), number of kernel rows, number of kernels/row, number of kernels/ear, days to maturity, hundred seed weight (g), shelling percentage and grain yield/plant (g). The statistical analysis was carried out using mean values of five plants over two replications for each character.

Based on *per se* performance of 144 inbreds, fifteen lines and four testers were chosen for the studying heterosis. The details of selected inbreds are presented in table 2. The experiment was conducted at the RandD farm, Foliage Crop Solutions Private Limited, Attur, Tamilnadu, India. Fifteen lines, four testers and the sixty hybrids derived by crossing the chosen lines and testers were raised in a randomized block design with three replications. The hybrids were randomized in a separate block and the parents in the adjacent block of the same layout as suggested by Arunachalam [6]. Three commercial hybrids namely 900M Gold, CP 818 and NK 6240 were included as standard checks for comparison and were randomized with the hybrids. Observations were recorded on 10 characters *viz.* days to 50% tasseling, days to 50% silking, days to maturity, ear length (cm), ear circumference (cm), number of kernel rows, number of kernels/row, number of kernels/ear, hundred seed weight(g) and grain yield (tonnes/ha). Line x tester analysis was done using the method suggested by Kempthorne (1957) and heterosis for individual characters was computed by the formulae suggested by Turner (1953) and Hayes *et al.* [7]. Statistical analysis for heterosis was carried out by WINDOSTAT software developed by Indostat services, Hyderabad, India. The mean values of observations on 10 consecutive plants were used for data analysis.

RESULTS

The results of heterosis (%) of 60 hybrids are presented in table 3. Except the cross “FI-113 x FI-109”, all the remaining 59 crosses exhibited significant heterosis over mid parent in the preferred negative direction for days to 50 % tasseling. As many as 13 hybrids showed significant heterosis over the best commercial check NK 6240. Of these, “FI-139 x FI-143” was the only hybrid which recorded significant negative heterosis with a magnitude of-6.0% while the remaining 12 hybrids had significant positive heterosis for days to 50% tasseling. The magnitude of standard heterosis for days to 50% tasseling ranged from-6.0 to 9.3%. Fifty nine crosses over mid parent and 58 crosses over better parent exhibited significant negative heterosis for days to 50 % silking. A range of-5.1% to 8.3% standard heterosis was observed over the best commercial check NK 6240. Out of 60 crosses, 14 recorded significant standard heterosis for days to 50% silking, of which in 13 crosses it was in positive direction and in one in the negative direction. The hybrid “FI-139 x FI-143” which recorded significant negative heterosis for days to 50%

tasseling was the only hybrid to record significant heterosis in the preferred negative direction.

Among the 60 crosses, 35 crosses exhibited significant negative heterosis over mid parent while six cross combinations showed significant positive heterosis for days to maturity. Of 42 hybrids which had significant heterosis over better parent, 40 were in the negative direction. The heterosis observed over the best commercial check NK 6240 ranged from-4.4 to 14.9. As many as 37 hybrids exhibited significant heterosis over the best check in which six were in the desired negative direction. The cross “FI-139 x FI-142” was the best combination recording negative significant standard heterosis of-4.4% followed by “FI-139 x FI-143” (-4.0%) and “FI-5 x FI-143” (-4.0%). Seventeen hybrids exhibited significantly positive heterosis over the mid parent while 10 hybrids (nine in positive and one in negative direction) showed significant heterosis over the better parent for the character number of kernel rows. The check 900M gold was found to be the best check with highest number of kernel rows. Out of 14 hybrids which had significant heterosis over 900M gold, four hybrids showed heterosis in the positive direction. This included 3 hybrids involving the tester FI-143, *viz.* “FI-54 x FI-143” (27.5%), “FI-5 x FI-143” (17.3%), “FI-101 x FI-143” (13.5%) and another hybrid “FI-127 x FI-142” (13.5%) involving the tester FI-142. The range of standard heterosis observed for number of kernel rows was from-19.4 and 27.5%. Except the cross “FI-139 x FI-142”, all the 59 crosses showed significant positive heterosis over mid parent for number of kernels per row. Out of 60 crosses, 55 exhibited significant positive heterosis over the better parent. A heterosis range of-28.3 to 8.3% was recorded over the best check CP 818. However, 17 hybrids exhibited significant negative heterosis over CP 818 and none of the hybrids registered significant positive heterosis.

As many as 42 hybrids exhibited significant heterosis over mid parent for hundred seed weight and it was in the desired positive direction in 41 hybrids and in the negative direction in one hybrid. While 19 hybrids showed significant positive heterosis, it was significantly negative over better parent in 10 hybrids. Out of 35 hybrids showing significant heterosis over the best check CP818, only in three hybrids it was in the positive direction. The hybrids which showed significantly positive standard heterosis were “FI-139 x FI-109” (20.4%), “FI-104 x FI-142” (11.9%) and “FI-104 x FI-109” (10.2%). Among the 60 hybrids under study, 58 showed significant positive heterosis for grain yield over mid parent. Over the better parent, 58 hybrids exhibited significant heterosis out of which 57 were in positive direction and in one it was in negative direction. A range of-63.4 to 14.6% standard heterosis was observed over the best check CP 818. In all, 51 hybrids had significantly negative heterosis over CP 818. In the order of merit, three hybrids *viz.* “FI-24 x FI-142” (14.6%), “FI-54 x FI-109” (14.0%) and “FI-54 x FI-142” (10.8%) exhibited significant positive heterosis for grain yield over the best check CP 818.

Table 1: Details of 144 maize inbreds used for diversity analysis

| S. No. | Source population | No. of inbreds | Inbred numbers |
|--------|---|----------------|--|
| 1 | Advanced generation of Single cross hybrids | 7 4 20 | FI-1, FI-2, FI-3, FI-4, FI-5, FI-6, FI-7 FI-29, FI-30, FI-31, FI-32 FI-33, FI-34, FI-35, FI-36, FI-37, FI-38, FI-39, FI-40, FI-41, FI-42, FI-43, FI-44, FI-45, |

| | | | |
|---|--|---------|---|
| | a. Chola | | FI-46, FI-47, FI-48, FI-49, FI-50, FI-51, FI-52 |
| | b. Ashoka | | |
| | c. SCH 55 | | |
| 2 | Advanced generation of Three way cross hybrids | 4 17 | FI-25, FI-26, FI-27, FI-28 FI-8, FI-9, FI-10, FI-11, FI-12, FI-13, FI-14, FI-15, FI-16, FI-17, FI-18, FI-19, FI-20, FI-21, FI-22, FI-23, FI-24 |
| | a. C555 | | |
| | b. TWH 001 | | |
| 3 | Hybrid mixtures | 62 | FI-53, FI-54, FI-55, FI-56, FI-57, FI-58, FI-59, FI-60, FI-61, FI-62, FI-63, FI-64, FI-65, FI-66, FI-67, FI-68, FI-69, FI-70, FI-71, FI-72, FI-73, FI-74, FI-75, FI-76, FI-77, FI-78, FI-79, FI-80, FI-81, FI-82, FI-83, FI-84, FI-85, FI-86, FI-87, FI-88, FI-89, FI-90, FI-91, FI-92, FI-93, FI-94, FI-95, FI-96, FI-97, FI-98, FI-99, FI-100, FI-101, FI-102, FI-103, FI-104, FI-105, FI-106, FI-107, FI-108, FI-109, FI-110, FI-111, FI-112, FI-113, FI-114 |
| 4 | Composite-CMP001 | 30 | FI-115, FI-116, FI-117, FI-118, FI-119, FI-120, FI-121, FI-122, FI-123, FI-124, FI-125, FI-126, FI-127, FI-128, FI-129, FI-130, FI-131, FI-132, FI-133, FI-134, FI-135, FI-136, FI-137, FI-138, FI-139, FI-140, FI-141, FI-142, FI-143, FI-144 |

Table 2: Details of fifteen lines and four testers chosen for study

| S. No. | Inbred no | Base population | Days to 50% tasseling | Days to 50% silking | Days to maturity | No. of kernel rows | No. of kernels/row | No. of kernels/ear | Hundred seed wt. (g) | Grain yield/plant (g) |
|----------------|-----------|------------------------|-----------------------|---------------------|------------------|--------------------|--------------------|--------------------|----------------------|-----------------------|
| Lines | | | | | | | | | | |
| 1 | FI-5 | Single cross hybrid | 59 | 61 | 106 | 13.2 | 26.2 | 344.0 | 33.0 | 107.8 |
| 2 | FI-7 | Single cross hybrid | 58 | 59 | 97 | 13.0 | 25.2 | 327.4 | 29.7 | 92.9 |
| 3 | FI-24 | Three way cross hybrid | 60 | 60 | 102 | 13.4 | 21.7 | 292.3 | 31.1 | 89.9 |
| 4 | FI-49 | Single cross hybrid | 63 | 65 | 104 | 13.8 | 27.3 | 376.8 | 28.9 | 102.1 |
| 5 | FI-54 | Hybrid mixture | 61 | 63 | 97 | 16.7 | 25.2 | 419.8 | 28.2 | 101.9 |
| 6 | FI-59 | Hybrid mixture | 63 | 64 | 102 | 14.0 | 26.0 | 365.3 | 26.9 | 93.8 |
| 7 | FI-101 | Hybrid mixture | 58 | 60 | 96 | 14.2 | 20.9 | 297.0 | 31.5 | 86.4 |
| 8 | FI-104 | Hybrid mixture | 56 | 58 | 101 | 13.0 | 27.7 | 359.6 | 29.6 | 113.4 |
| 9 | FI-113 | Hybrid mixture | 66 | 68 | 128 | 12.4 | 24.9 | 308.2 | 36.6 | 94.9 |
| 10 | FI-114 | Hybrid mixture | 64 | 67 | 118 | 14.6 | 23.1 | 337.4 | 32.1 | 90.9 |
| 11 | FI-127 | Composite | 58 | 61 | 96 | 11.0 | 22.5 | 247.2 | 33.0 | 80.2 |
| 12 | FI-130 | Composite | 59 | 60 | 102 | 14.4 | 21.2 | 305.3 | 31.8 | 78.9 |
| 13 | FI-139 | Composite | 59 | 60 | 95 | 11.0 | 21.3 | 234.5 | 40.5 | 88.1 |
| 14 | FI-141 | Composite | 61 | 63 | 107 | 13.0 | 23.7 | 307.1 | 31.1 | 99.8 |
| 15 | FI-144 | Composite | 62 | 65 | 105 | 15.2 | 19.6 | 297.2 | 26.8 | 85.5 |
| Testers | | | | | | | | | | |
| 1 | FI-109 | Hybrid mixture | 59 | 61 | 106 | 13.0 | 24.9 | 323.5 | 36.8 | 112.3 |
| 2 | FI-140 | Composite | 59 | 61 | 97 | 14.2 | 23.7 | 337.7 | 24.1 | 74.7 |
| 3 | FI-142 | Composite | 59 | 61 | 94 | 13.4 | 19.7 | 264.0 | 35.2 | 88.5 |
| 4 | FI-143 | Composite | 57 | 59 | 102 | 16.2 | 27.7 | 452.9 | 21.1 | 87.5 |

Table 3: Heterosis (%) of 60 hybrids over mid parent, better parent and the best commercial check

| S. No. | Hybrids | Days to 50% tasseling | | | Days to 50% silking | | | Days to maturity | | | Number of kernel rows | | |
|--------|----------------|-----------------------|---------------|---------|---------------------|---------|---------|------------------|---------------|---------|-----------------------|---------------|-----------|
| | | Mid Parent | Better Parent | NK 6240 | Better Parent | CP 818 | NK 6240 | Mid Parent | Better Parent | NK 6240 | Mid Parent | Better Parent | 900M Gold |
| 1 | FI-5 x FI-109 | -8.5** | -8.5** | 0.0 | -4.7* | -5.3* | 3.9 | -2.4 | -5.2** | 5.1** | 13.8* | 6.5 | -6.8 |
| 2 | FI-5 x FI-140 | -8.4** | -10.0** | 2.0 | -10.0** | -11.8** | 0.6 | -10.3** | -11.8** | -2.2 | 15.4** | 13.0* | -1.1 |
| 3 | FI-5 x FI-142 | -9.5** | -12.1** | 2.0 | -12.5** | -14.9** | -1.3 | -10.9** | -12.8** | -3.3* | 6.5 | 4.2 | -4.6 |
| 4 | FI-5 X FI-143 | -9.8** | -9.8** | -1.3 | -7.0** | -7.0** | 1.9 | -10.0** | -13.4** | -4.0* | 21.3** | 10.8* | 17.3** |
| 5 | FI-7 x FI-109 | -10.5** | -12.3** | 0.0 | -6.7** | -8.5** | 3.2 | 1.4 | 0.3 | 6.9** | 3.7 | -5.1 | -12.7* |
| 6 | FI-7 x FI-140 | -11.4** | -11.7** | 0.7 | -10.7** | -11.2** | 1.3 | -5.1** | -5.4** | 1.5 | 6.7 | 2.0 | -6.1 |
| 7 | FI-7 x FI-142 | -12.5** | -13.2** | 0.7 | -13.2** | -14.4** | -0.6 | -7.0** | -7.1** | -1.1 | -4.9 | -5.1 | -12.7* |
| 8 | FI-7 x FI-143 | -11.6** | -13.5** | -1.3 | -9.5** | -10.8** | 0.6 | -6.8** | -8.5** | -2.5 | 5.3 | -1.6 | 4.2 |
| 9 | FI-24 x FI-109 | -8.3** | -10.9** | 3.3 | -5.2** | -7.8** | 5.8* | -1.0 | -4.8** | 7.6** | 5.8 | -2.3 | -11.8* |
| 10 | FI-24 x FI-140 | -11.1** | -12.1** | 2.0 | -9.2** | -9.5** | 3.9 | -4.0** | -6.4** | 5.8** | 10.9 | 7.0 | -3.4 |
| 11 | FI-24 x FI-142 | -12.1** | -12.1** | 2.0 | -12.8** | -13.3** | 0.6 | -7.1** | -9.9** | 1.8 | -0.2 | -0.9 | -9.3 |
| 12 | FI-24 x FI-143 | -10.7** | -13.2** | 0.7 | -9.7** | -11.7** | 1.3 | -9.2** | -13.5** | -2.2 | -0.8 | -8.1 | -2.6 |
| 13 | FI-49 x FI-109 | -5.3** | -9.0** | 8.0** | -4.3* | -7.7** | 7.7** | 3.2* | 1.3 | 9.8** | 12.2* | 3.3 | -6.1 |
| 14 | FI-49 x FI-140 | -8.6** | -10.7** | 6.0** | -7.8** | -8.8** | 6.4** | -9.6** | -10.0** | -2.5 | 2.3 | -1.6 | -10.6 |
| 15 | FI-49 x FI-142 | -10.2** | -11.2** | 5.3* | -10.7** | -11.0** | 3.9 | -0.7 | -1.7 | 6.5** | 3.6 | 3.2 | -5.5 |
| 16 | FI-49 x FI-143 | -6.4** | -10.1** | 6.7** | -6.0** | -8.8** | 6.4** | -1.0 | -3.7* | 4.4** | 7.6 | 0.0 | 5.9 |
| 17 | FI-54 x FI-109 | -5.7** | -6.6** | 4.0 | -5.5** | -6.9** | 3.9 | 1.6 | 1.4 | 6.2** | 26.2** | 19.9** | 1.7 |
| 18 | FI-54 x FI-140 | -7.4** | -8.2** | 4.0 | -6.8** | -7.9** | 5.1* | -5.3** | -6.4** | 0.4 | 21.0** | 20.4 | 2.1 |
| 19 | FI-54 x FI-142 | -8.5** | -10.3** | 4.0 | -9.3** | -11.1** | 3.2 | -8.6** | -9.2** | -3.6* | 14.4* | 10.1 | 0.8 |

| | | | | | | | | | | | | | |
|----|-----------------|---------|---------|-------|---------|---------|-------|---------|---------|--------|--------|--------|---------|
| 20 | FI-54 x FI-143 | -7.6** | -8.4** | 2.0 | -6.1** | -6.9** | 3.9 | -6.3** | -7.3** | -2.9 | 33.7** | 20.4 | 27.5** |
| 21 | FI-59 x FI-109 | -8.7** | -10.5** | 2.0 | -5.0** | -5.8** | 3.9 | 4.2** | 4.2** | 8.7** | 6.6 | -1.9 | -11.0 |
| 22 | FI-59 x FI-140 | -9.1** | -9.4** | 3.3 | -9.1** | -10.7** | 1.9 | -0.7 | -2.0 | 5.1** | 3.4 | -0.5 | -9.7 |
| 23 | FI-59 x FI-142 | -9.6** | -10.3** | 4.0 | -9.4** | -11.6** | 2.6 | -2.9* | -3.8* | 2.2 | -1.9 | -2.3 | -10.6 |
| 24 | FI-59 x FI-143 | -6.9** | -8.8** | 4.0 | -3.8* | -4.1 | 5.8* | -6.1** | -6.9** | -2.9 | 3.0 | -4.4 | 1.3 |
| 25 | FI-101 x FI-109 | -11.0** | -13.3** | 0.0 | -7.3** | -8.1** | 1.3 | -0.2 | -0.4 | 4.4** | 9.0 | -0.9 | -7.6 |
| 26 | FI-101 x FI-140 | -12.0** | -12.7** | 0.7 | -10.3** | -11.8** | 0.6 | -0.2 | -1.4 | 5.8** | 10.0 | 4.5 | -2.5 |
| 27 | FI-101 x FI-142 | -13.0** | -13.2** | 0.7 | -13.3** | -15.5** | -1.9 | -3.8** | -4.4** | 1.5 | -0.5 | -1.4 | -8.0 |
| 28 | FI-101 x FI-143 | -12.2** | -14.5** | -1.3 | -10.2** | -10.5** | -1.3 | -0.4 | -1.4 | 3.3* | 14.0** | 7.2 | 13.5* |
| 29 | FI-104 x FI-109 | -8.6** | -11.0** | 2.7 | -5.7** | -8.8** | 5.8* | 4.8** | 2.3 | 12.0** | -0.4 | -5.7 | -19.4** |
| 30 | FI-104 x FI-140 | -11.4** | -12.1** | 1.3 | -9.8** | -10.5** | 3.9 | -5.0** | -6.0** | 2.9 | 2.6 | 1.7 | -13.1* |
| 31 | FI-104 x FI-142 | -9.5** | -9.8** | 4.7* | -11.1** | -11.1** | 3.2 | -5.9** | -7.3** | 1.5 | -1.3 | -4.6 | -12.7* |
| 32 | FI-104 x FI-143 | -11.0** | -13.3** | 0.0 | -9.1** | -11.6** | 2.6 | -5.3** | -8.3** | 0.4 | 8.9 | -1.6 | 4.2 |
| 33 | FI-113 x FI-109 | -2.7** | -4.7* | 8.7** | -2.0 | -4.0 | 8.3** | 7.2** | 5.4** | 13.8** | 6.8 | -2.3 | -10.1 |
| 34 | FI-113 x FI-140 | -7.9** | -8.2** | 4.7* | -7.9** | -8.4** | 4.5 | -3.4** | -3.7* | 4.0* | 2.6 | -1.8 | -9.7 |
| 35 | FI-113 x FI-142 | -9.0** | -9.8** | 4.7* | -9.2** | -10.5** | 3.9 | 0.5 | -0.3 | 7.6** | -0.9 | -1.2 | -9.1 |
| 36 | FI-113 x FI-143 | -6.9** | -8.8** | 4.0 | -6.6** | -8.0** | 3.9 | 3.3* | 0.7 | 8.7** | 8.6 | 1.4 | 7.4 |
| 37 | FI-114 x FI-109 | -8.1** | -12.2** | 5.3* | -6.4** | -11.2** | 7.1** | -1.9 | -11.5** | 14.9** | 11.8 | 8.4 | -11.8* |
| 38 | FI-114 x FI-140 | -8.0** | -10.6** | 7.3** | -8.7** | -11.2** | 7.1** | -10.7** | -18.4** | 5.8** | 22.0** | 20.1** | 0.8 |
| 39 | FI-114 x FI-142 | -10.7** | -12.2** | 5.3* | -12.2** | -13.8** | 3.9 | -10.3** | -18.4** | 5.8** | 11.3 | 5.1 | -3.8 |
| 40 | FI-114 x FI-143 | -9.9** | -13.9** | 3.3 | -7.5** | -11.7** | 6.4** | -6.4** | -16.2** | 8.7** | 13.6* | 0.4 | 6.3 |
| 41 | FI-127 x FI-109 | -8.3** | -8.5** | 0.0 | -6.1** | -8.0** | 3.9 | 3.9** | 2.4 | 10.1** | 15.3* | 14.1 | -11.0 |
| 42 | FI-127 x FI-140 | -8.1** | -10.0** | 2.0 | -9.0** | -9.6** | 3.2 | -8.9** | -9.1** | -2.2 | 18.2** | 14.1* | -4.2 |
| 43 | FI-127 x FI-142 | -11.6** | -14.4** | -0.7 | -12.6** | -13.8** | 0.0 | -8.8** | -9.4** | -2.5 | 33.8** | 24.0** | 13.5* |
| 44 | FI-127 x FI-143 | -8.9** | -9.2** | -0.7 | -7.8** | -9.1** | 2.6 | -8.3** | -10.4** | -3.6* | 2.3 | -11.2* | -5.9 |
| 45 | FI-130 x FI-109 | -8.4** | -9.5** | 1.3 | -5.3** | -5.9** | 3.2 | 1.2 | -0.3 | 7.3** | 12.2* | 0.4 | -3.0 |
| 46 | FI-130 x FI-140 | -11.8** | -12.4** | -0.7 | -10.0** | -11.8** | 0.6 | -3.2* | -3.4* | 4.0* | 6.5 | -0.4 | -3.8 |
| 47 | FI-130 x FI-142 | -10.5** | -12.1** | 2.0 | -10.2** | -12.7** | 1.3 | -2.0 | -2.7 | 4.7** | -4.5 | -7.0 | -10.1 |
| 48 | FI-130 x FI-143 | -10.2** | -11.3** | -0.7 | -8.2** | -8.2** | 0.6 | -2.8* | -5.1* | 2.2 | 8.3 | 3.6 | 9.7 |
| 49 | FI-139 x FI-109 | -11.0** | -12.9** | -0.7 | -10.5** | -12.0** | -1.3 | 0.7 | 0.7 | 5.1** | 3.9 | -1.2 | -6.2** |
| 50 | FI-139 x FI-140 | -12.6** | -12.9** | -0.7 | -12.2** | -12.9** | -0.6 | -8.2** | -9.5** | -2.9 | 4.5 | 4.0 | -11.8* |
| 51 | FI-139 x FI-142 | -4.9** | -5.8** | 9.3** | -7.3** | -8.8** | 5.8* | -9.1** | -9.9** | -4.4** | -2.9 | -6.5 | -14.4* |
| 52 | FI-139 x FI-143 | -15.8** | -17.5** | -6.0* | -14.5** | -15.4** | -5.1* | -7.2** | -8.0** | -4.0* | 5.3 | -5.2 | 0.4 |
| 53 | FI-141 x FI-109 | -8.2** | -11.4** | 4.0 | -4.7* | -5.8** | 4.5 | 1.4 | 0.0 | 7.3** | 12.1* | 4.8 | -8.0 |
| 54 | FI-141 x FI-140 | -10.4** | -11.9** | 3.3 | -9.4** | -10.7** | 1.9 | -2.0 | -2.0 | 5.1** | 16.5** | 13.9* | 0.0 |
| 55 | FI-141 x FI-142 | -11.4** | -11.9** | 3.3 | -10.2** | -12.2** | 1.9 | -5.9** | -6.4** | 0.4 | 0.2 | -1.8 | -10.1 |
| 56 | FI-141 x FI-143 | -10.0** | -13.1** | 2.0 | -5.8** | -6.4** | 3.9 | -1.2 | -3.4* | 3.6* | 0.2 | -8.4 | -3.0 |
| 57 | FI-144 x FI-109 | -8.3** | -10.9** | 3.3 | -5.4** | -8.8** | 6.4** | -1.0 | -2.4 | 4.7** | 10.6 | 0.0 | -5.5 |
| 58 | FI-144 x FI-140 | -11.1** | -12.1** | 2.0 | -11.1** | -12.1** | 2.6 | -5.4** | -5.4** | 1.5 | 8.7 | 2.7 | -3.0 |
| 59 | FI-144 x FI-142 | -10.9** | -10.9** | 3.3 | -10.7** | -11.0** | 3.9 | -3.9** | -4.4** | 2.5 | 5.2 | 3.6 | -2.1 |
| 60 | FI-144 x FI-143 | -10.7** | -13.2** | 0.7 | -9.4** | -12.1** | 2.6 | -3.6** | -5.7** | 1.1 | -1.5 | -6.8 | -1.3 |

Table 3: (contd.)

| S. No. | Hybrids | Number of kernels/row | | | Number of kernels/ear | | | 100 seed weight | | | Grain yield | | |
|--------|----------------|-----------------------|---------------|---------|-----------------------|---------------|-----------|-----------------|---------------|---------|-------------|---------------|---------|
| | | Mid Parent | Better Parent | CP 818 | Mid Parent | Better Parent | 900M Gold | Mid Parent | Better Parent | CP 818 | Mid Parent | Better Parent | CP 818 |
| 1 | FI-5 x FI-109 | 24.5** | 22.9** | -9.3 | 41.7** | 34.2* | -14.6 | 8.3 | 0.7 | -3.5 | 92.2** | 87.1** | -13.5** |
| 2 | FI-5 x FI-140 | 32.5** | 22.3** | -12.1* | 52.4** | 37.9** | -12.2 | -5.4 | -7.2 | -20.5** | 99.0** | 55.6** | -28.0** |
| 3 | FI-5 x FI-142 | 29.7** | 23.4** | -11.3 | 38.0** | 34.2* | -14.5 | 25.2** | 22.8** | 1.2 | 87.2** | 86.1** | -13.9** |
| 4 | FI-5 X FI-143 | 34.8** | 29.0** | -7.3 | 63.9** | 56.3** | 9.8 | 15.3* | -13.2* | -28.5** | 65.9** | 58.1** | -26.9** |
| 5 | FI-7 x FI-109 | 33.4** | 29.6** | 1.5 | 37.8** | 22.7 | -10.6 | 18.9** | 7.8 | 3.3 | 94.7** | 90.8** | -12.9** |
| 6 | FI-7 x FI-140 | 35.0** | 19.9** | -6.1 | 43.1** | 22.2 | -10.9 | 12.4* | 7.2 | -8.1 | 126.3** | 77.8** | -18.8** |
| 7 | FI-7 x FI-142 | 36.3** | 24.6** | -2.4 | 29.4* | 18.0 | -14.0 | 37.1** | 35.9** | 7.7 | 102.7** | 102.5** | -7.4 |
| 8 | FI-7 x FI-143 | 35.2** | 24.2** | -2.7 | 43.2** | 40.6** | 2.5 | 17.6** | -9.7 | -29.7** | 64.0** | 57.2** | -28.2** |
| 9 | FI-24 x FI-109 | 65.3** | 46.7** | 8.3 | 76.8** | 69.3** | -3.6 | -4.7 | -9.8* | -13.6** | 110.6** | 92.8** | -15.6** |
| 10 | FI-24 x FI-140 | 61.5** | 56.6** | -4.7 | 79.7** | 78.8** | -6.8 | -1.9 | -2.0 | -16.0** | 158.0** | 121.4** | -19.4** |
| 11 | FI-24 x FI-142 | 68.8** | 58.7** | 3.1 | 68.5** | 57.2** | -5.5 | 22.5** | 18.1** | 0.9 | 179.1** | 150.6** | 14.6** |
| 12 | FI-24 x FI-143 | 65.5** | 54.8** | 1.6 | 63.1** | 42.1** | -0.2 | 3.7 | -22.9** | -34.1** | 53.1** | 43.0** | -40.1** |
| 13 | FI-49 x FI-109 | 32.5** | 30.8** | -3.5 | 48.9** | 38.8** | -8.6 | -5.4 | -7.8 | -11.6* | 80.3** | 78.2** | -20.1** |
| 14 | FI-49 x FI-140 | 20.3** | 11.1 | -20.1** | 23.4 | 10.1 | -27.5** | -12.9** | -15.3** | -23.1** | 48.4** | 17.3 | -47.4** |
| 15 | FI-49 x FI-142 | 35.2** | 28.7** | -7.5 | 40.2** | 34.1* | -11.6 | 19.0** | 11.4* | 1.2 | 88.3** | 86.5** | -14.7** |
| 16 | FI-49 x FI-143 | 24.0** | 18.6* | -14.7* | 34.5** | 30.3* | -8.5 | 1.6 | -25.9** | -32.7** | 60.8** | 55.5** | -30.2** |
| 17 | FI-54 x FI-109 | 32.8** | 31.5** | -2.9 | 67.7** | 60.7** | -0.2 | 0.3 | -2.8 | -6.8 | 156.5** | 152.8** | 14.0** |
| 18 | FI-54 x FI-140 | 36.8** | 25.9** | -8.9 | 65.8** | 51.8** | -5.7 | -6.8 | -9.0 | -18.2** | 166.4** | 110.2** | -5.2 |
| 19 | FI-54 x FI-142 | 42.1** | 34.8** | -2.4 | 62.7** | 60.1** | -0.6 | 17.0 | 10.1* | -1.1 | 144.0** | 142.2** | 10.8* |
| 20 | FI-54 x FI-143 | 51.3** | 44.3** | 4.5 | 103.5** | 91.7** | 34.7** | -2.4 | -28.6** | -35.8** | 70.8** | 64.7** | -25.7** |
| 21 | FI-59 x FI-109 | 25.6** | 22.8** | -9.4 | 34.1** | 26.1 | -18.6* | -7.7 | -15.5** | -19.0** | 30.9** | 21.0* | -37.6** |
| 22 | FI-59 x FI-140 | 44.8** | 34.9** | -4.9 | 49.4** | 34.4* | -13.2 | -2.6 | -6.0 | -19.4** | 114.1** | 61.2** | -16.9** |
| 23 | FI-59 x FI-142 | 54.9** | 48.7** | 4.9 | 52.0** | 46.7** | -5.2 | 27.3** | 27.0** | 1.2 | 85.8** | 75.3** | -9.6* |
| 24 | FI-59 x FI-143 | 42.7** | 37.7** | -2.9 | 47.2** | 41.3** | -0.7 | 29.6** | -1.3 | -21.4** | 36.5** | 23.8** | -36.2** |

| | | | | | | | | | | | | | |
|----|-----------------|--------|--------|---------|---------|--------|---------|--------|---------|---------|---------|---------|---------|
| 25 | FI-101 x FI-109 | 27.8** | 18.1* | -12.9* | 40.3** | 38.0* | -18.7* | 14.3** | 1.5 | -2.7 | 109.2** | 84.0** | -19.4** |
| 26 | FI-101 x FI-140 | 40.3** | 38.4** | -13.5* | 53.7** | 44.2** | -15.1 | 15.0** | 7.4 | -7.9 | 189.8** | 158.5** | -14.0** |
| 27 | FI-101 x FI-142 | 33.0** | 30.5** | -15.2* | 32.2* | 30.9* | -21.3* | 37.5** | 33.2** | 5.6 | 112.9** | 83.9** | -15.9** |
| 28 | FI-101 x FI-143 | 23.4** | 20.5* | -20.9** | 40.5** | 29.2* | -9.3 | 22.6** | -4.4 | -28.9** | 61.8** | 45.1** | -39.2** |
| 29 | FI-104 x FI-109 | 31.6** | 28.7** | -5.0 | 31.6* | 27.4 | -22.5* | 29.6** | 14.9** | 10.2* | 107.0** | 99.8** | -12.5** |
| 30 | FI-104 x FI-140 | 38.3** | 28.8** | -9.1 | 42.0** | 31.2* | -20.2* | 17.7** | 9.8 | -5.9 | 148.7** | 103.9** | -16.9** |
| 31 | FI-104 x FI-142 | 34.9** | 29.5** | -8.6 | 33.2** | 32.4* | -19.5* | 45.8** | 41.2** | 11.9** | 112.7** | 101.1** | -8.0 |
| 32 | FI-104 x FI-143 | 40.6** | 35.7** | -4.2 | 54.1** | 43.8** | 1.0 | 39.7** | 9.1 | -19.0** | 78.0** | 75.5** | -26.5** |
| 33 | FI-113 x FI-109 | 43.9** | 31.8** | -2.7 | 54.8** | 54.5** | -11.8 | 2.5 | -10.5* | -14.2** | 71.6** | 67.3** | -22.9** |
| 34 | FI-113 x FI-140 | 54.7** | 54.0** | -5.4 | 58.5** | 50.9** | -13.9 | 13.4** | 4.0 | -10.8* | 91.1** | 49.6** | -31.0** |
| 35 | FI-113 x FI-142 | 46.4** | 42.5** | -7.5 | 44.6** | 41.0** | -15.2 | 34.1** | 27.6** | 1.1 | 64.3** | 63.7** | -24.6** |
| 36 | FI-113 x FI-143 | 50.4** | 45.6** | -4.4 | 62.7** | 47.5** | 3.6 | 38.7** | 9.7 | -21.5** | 72.9** | 65.1** | -23.9** |
| 37 | FI-114 x FI-109 | 40.8** | 13.5 | -16.2** | 59.0** | 31.3* | -25.3** | 11.0* | 1.8 | -2.4 | 106.7** | 105.8** | -9.9* |
| 38 | FI-114 x FI-140 | 64.8** | 43.5** | -12.7* | 100.9** | 72.6** | -10.9 | 3.4 | -0.1 | -14.3** | 119.1** | 75.3** | -23.9** |
| 39 | FI-114 x FI-142 | 70.6** | 44.6** | -6.1 | 87.7** | 51.7** | -8.7 | 26.7** | 26.1** | 0.9 | 110.7 | 105.3 | -6.1 |
| 40 | FI-114 x FI-143 | 75.5** | 48.1** | -2.8 | 94.3** | 48.4** | 4.3 | 27.9** | -2.8 | -22.2** | 106.6 | 103.0 | -11.9** |
| 41 | FI-127 x FI-109 | 44.2** | 36.3** | 0.6 | 66.1** | 58.6** | -9.7 | 15.3* | 8.2 | 3.8 | 94.0** | 88.9** | -12.7** |
| 42 | FI-127 x FI-140 | 50.4** | 44.9** | -4.8 | 78.3** | 78.0** | -7.8 | 2.2 | 1.3 | -13.2** | 140.5** | 88.1** | -13.1** |
| 43 | FI-127 x FI-142 | 46.5** | 45.6** | -4.3 | 97.6** | 83.9** | 10.6 | 22.1** | 18.5** | -0.2 | 88.9** | 88.0** | -13.1** |
| 44 | FI-127 x FI-143 | 40.3** | 40.2** | -7.9 | 43.6** | 24.8 | -12.4 | 13.7* | -15.0** | -28.4** | 36.3** | 29.9** | -39.9** |
| 45 | FI-130 x FI-109 | 14.8* | 11.4 | -17.7** | 29.1* | 18.7 | -19.4* | 19.4** | 3.0 | -1.3 | 86.0** | 80.9** | -20.8** |
| 46 | FI-130 x FI-140 | 32.4** | 24.2** | -13.7* | 40.5** | 23.6 | -16.1 | 16.3** | 5.3 | -9.7 | 132.6** | 89.6** | -21.5** |
| 47 | FI-130 x FI-142 | 17.0* | 13.1 | -21.3** | 11.7 | 5.3 | -28.5** | 41.5** | 32.8** | 5.3 | 85.1** | 76.3** | -19.4** |
| 48 | FI-130 x FI-143 | 42.2** | 38.2** | -3.9 | 53.9** | 51.3** | 6.3 | 22.6** | -2.0 | -31.9** | 90.6** | 89.5** | -20.6** |
| 49 | FI-139 x FI-109 | 32.9** | 28.2** | -5.4 | 38.0** | 35.7** | -20.0* | 45.0** | 25.6** | 20.4** | 109.6** | 104.0** | -10.6* |
| 50 | FI-139 x FI-140 | 47.7** | 39.3** | -4.4 | 54.2** | 44.6** | -14.8 | 29.2** | 17.5** | 0.8 | 164.2** | 115.1** | -10.8* |
| 51 | FI-139 x FI-142 | 7.4 | 4.5 | -28.3** | 5.5 | 4.4 | -37.2** | 17.4** | 10.7 | -12.3** | -16.1 | -20.0* | -63.4** |
| 52 | FI-139 x FI-143 | 47.5** | 44.3** | -1.0 | 55.4** | 42.9** | 0.4 | 47.8** | 17.8** | -17.3** | 110.1** | 109.1** | -12.4** |
| 53 | FI-141 x FI-109 | 42.3** | 38.6** | 2.3 | 59.8** | 53.4** | -5.0 | 0.2 | -5.7 | -9.6* | 90.2** | 68.0** | -4.0 |
| 54 | FI-141 x FI-140 | 54.8** | 44.7** | 1.2 | 80.0** | 65.0** | 2.1 | 2.2 | 1.6 | -12.9** | 97.3** | 43.7** | -17.9** |
| 55 | FI-141 x FI-142 | 50.8** | 45.4** | 1.7 | 51.3** | 49.2** | -7.7 | 25.9** | 21.9** | 3.2 | 109.0** | 88.2** | 7.5 |
| 56 | FI-141 x FI-143 | 24.5** | 20.7* | -15.6** | 25.9* | 18.4 | -16.8 | 2.7 | -23.4** | -35.1** | 0.4 | -13.0 | -50.3** |
| 57 | FI-144 x FI-109 | 31.3** | 22.9** | -9.3 | 47.0** | 41.5** | -13.0 | 24.9** | 6.7 | 2.3 | 120.8** | 93.9** | -15.1** |
| 58 | FI-144 x FI-140 | 39.5** | 35.6** | -12.7* | 51.5** | 39.3** | -14.4 | 17.1** | 4.9 | -10.1* | 179.5** | 149.8** | -17.3** |
| 59 | FI-144 x FI-142 | 20.9** | 20.4* | -21.8** | 27.0* | 25.6 | -22.7* | 41.8** | 31.7** | 4.4 | 119.7** | 89.4** | -13.4** |
| 60 | FI-144 x FI-143 | 34.8** | 33.5** | -12.4* | 33.3** | 25.0* | -12.2 | 40.4** | 13.2* | -23.1** | 85.4** | 66.0** | -30.5** |

*-significant at 5% level; **-significant at 1% level

DISCUSSION

Earliness for days to 50% tasseling, days to 50% silking and days to maturity are considered as desirable traits to ensure better hybrid performance under rainfed conditions. The hybrid “FI-139 x FI-143” recorded significantly negative heterosis over mid parent, better parent and the standard check for these characters. Both the inbreds involved in this cross were derived from composite base populations. Also another 5 hybrid combinations showed significant negative heterosis over mid parent, better parent and standard check. It was found that significant negative heterosis was observed in crosses which involve testers from composite base population confirming their suitability in developing early inbreds and hybrids. Bhavana *et al.* [8] observed significant negative heterosis for days to 50% tasseling and days to 50% silking. Xiacong Zhang *et al.* [9] evaluated the parental populations and 21 crosses.

Four crosses *viz.* “FI-54 x FI-143” (27.5%), “FI-5 x FI-143” (17.3%), “FI-101 x FI-143” (13.5%) and “FI-127 x FI-142” (13.5%) recorded significant positive standard heterosis for number of kernel rows. These hybrids had parents derived from single cross hybrid, hybrid mixture and composite base population. Among these 2 hybrids, “FI-54 x FI-143” which is a cross between inbreds derived from hybrid mixture and composite base population registered significant positive standard heterosis also. Three hybrids which had both the parents as good general combiners *viz.* “FI-139 x FI-109”, “FI-104 x FI-142” and “FI-104 x FI-109” exhibited significant positive heterosis over the standard

check. All the inbreds involved in the crosses were derivatives of composite and hybrid mixture base population. Obtaining higher grain yield is the ultimate objective of any maize breeding program. Sixty hybrids over mid parent and 58 over better parent exhibited significant positive heterosis for grain yield which substantiates the importance and usefulness of heterosis breeding in maize yield improvement. Three hybrids *viz.* “FI-24 x FI-142”, “FI-54 x FI-109” and “FI-54 x FI-142” recorded significant positive standard heterosis for grain yield.

It was noticed that among the five superior cross combinations which had significant heterosis for grain yield, “FI-24 x FI-142” was the only cross which involved inbreds derived from three-way cross hybrid and composite base population. Two cross combinations namely “FI-54 x FI-142” and “FI-113 x FI-142” were the crosses between inbred parents derived from hybrid mixtures and composite base population. The remaining two hybrids *viz.* “FI-54 x FI-109” and “FI-139 x FI-143” involved the cross between the inbreds derived from same base population namely hybrid mixture and composite respectively. Wende Abera *et al.* [10] reported almost similar results.

This clearly indicated that composites and hybrid mixtures could be important source populations for deriving inbred parents for utilization in single cross hybrid development. Further, superiority of crosses between inbreds derived from the same base population for grain yield indicated the possibilities of deriving heterotic inbred lines from the

same base population to develop single cross hybrids for successful commercial exploitation.

ACKNOWLEDGEMENT

The support provided by M/S. Foliage crop solutions private limited, Chennai, India to carry out the research work is gratefully acknowledged.

REFERENCES

1. Duvick, DN, 1984. Genetics contribution to yield gain in U. S hybrid maize. In: Genetic contribution to yield gain of five major crop plants. pp: 15-47. Crop Science Society of America.
2. Kaul J, SainDass R, Saikumar R, PayalBansal, Om Prakash, Usha Nara, Bilal Ahmad, 2011. Single cross hybrid breeding technology in maize. Directorate of maize research, Pusa campus, New Delhi. Technical bulletin 2011/9.
3. Arshad M, Bakhsh A, Zubair M, Ghafoor A, 2003. Genetic variability and correlation studies in chickpea (*Cicer arietinum* L.). Pak J Bot 35: 605-611.
4. Chandel U, Mankotia BS, Thakur KS, 2014. Assesment of recombinant lines of maize hybrids for inbred development. Bangl J Bot 43: 363-366.
5. Melania MD, Carena MJ, 2005. Alternative maize heterotic patterns for the northen corn belt. Crop Sci 45:2186-2194.
6. Arunachalam V, 1974. The fallacy behind the use of a modified line x tester design. Indian J Genet Pl Br 34: 280-287.
7. Hayes HK, Immer FR, Smith DC, 1955. Methods of Plant Breeding. Mc Graw Hill Book Co Inc, New York.551 pp.
8. Bhavana P, Singh RP, Gadag RN, 2011. Gene action and heterosis for yield and yield components in maize (*Zea mays* L.). Indian J Agr Sci 81: 163-166.
9. Xiacong Zhang, Hongjun Yong, Zhiqiang Zhou, Chaoshu Zhang, Ming Lu, Qi Sun, Lin Zhang, Mingshun Li, Degui Zhang, Jianfeng Weng *et al.*, 2017. Heterosis and combining ability of seven maize germplasm populations. Euphytica 213:45.
10. Wende Abera, Shimelis Hussein, John Derera, Mosisa Worku, Mark Lain, 2016. Heterosis and combining ability of elite maize inbred lines under northern corn leaf blight disease prone environments of the mid-altitude tropics. Euphytica 208: 391-400.