

Antioxidants: elixir of life

Sadaf Kalam*, Rupal Singh, Abin Mani, Jaswant Patel, Firoz Naem Khan and Adarsh Pandey#

Centre for Scientific Research and Development (C.S.R.D.), People's Group, Bhopal-462037, MP, India.

Department of Botany, SS (P.G.) College, Shahjahanpur-242001, UP, India

Abstract

There has been a current upsurge in the medical implications of free radicals and related species during the past several decades. These chemical species are integral components produced during normal biochemical and physiological processes but leads to oxidative stress when produce in excess and causes potential damage to cells. A wide range of non-enzymatic and enzymatic antioxidant defenses exists to counteract the damaging effects of free radicals. There exist epidemiological evidences correlating higher intake of antioxidant rich foodstuffs with greater free radical neutralizing potential to lower incidence of several human morbidities or mortalities. Gene therapy to produce more antioxidants in the body, novel biomolecules and the use of functional foods enriched with antioxidants are milestones to newer approaches to reduce free radical damage. This paper reviews the biology of reactive species, their pathways through which they relate to the pathology of various diseases and discusses the putative roles that antioxidants, from different sources, play in controlling oxidative stress and reduce the incidence of concerned diseases.

Keywords: Antioxidants, Free radicals, Oxidative stress, Diseases.

INTRODUCTION

Greek physician Hippocrates back in 400 B.C., said, "Let food be your medicine and medicine be your food." Today, good nutrition is more important than anything else. At least four of the 10 leading causes of death in the world like heart disease, cancer, stroke and diabetes are directly related to the way we eat. Recent research has shown a significant relationship between free radicals and these diseases. It is paradoxical that oxygen, which is an element indispensable for life, can have severely deleterious effects on the human body under certain situations. Most potential harmful effects of oxygen can be credited to the formation and activity of a number of chemical compounds, known as reactive oxygen species (ROS), which have a tendency to donate oxygen to other substances. Many such reactive species are free radicals and have a surplus of one or more free-floating electrons rather than having matched pairs and are, therefore, unstable, highly reactive and responsible for deleterious cellular damage. To stabilize and deactivate these species, organisms have developed a complex antioxidants protection system capable of critically maintaining normal cellular and physiological functions.

Free radicals

Free Radicals are highly reactive molecular species with an unpaired electron, which causes them to capture electrons from other substances in order to neutralize themselves. Free radicals

derivatives of oxygen like superoxide free radical anion ($O_2^{\cdot-}$), hydroxyl free radical (OH^{\cdot}), nitric oxide radical (NO^{\cdot}), lipid peroxy (LO^{\cdot}), lipid alkoxyl (LOO^{\cdot}) and lipid peroxide ($LOOH$) as well as non-radical derivatives such as hydrogen peroxide and singlet oxygen are collectively known as reactive oxygen species (ROS). The hydroxyl radical ($^{\cdot}OH$), which is the neutral form of hydroxide ion, has a high reactivity, making it very dangerous radical with a very short *in vivo* half-life of approximately 10^{-9} seconds [1]. These potentially damaging molecules are capable of attacking healthy cells of the body causing them to lose their structure and function [2].

Although the initial attack causes the free radical to become neutralized, another free radical is formed in the process, causing a chain reaction to occur and consequently leads to oxidative stress. Such unstable molecules react easily with essential molecules of our body including DNA, fats and proteins [3]. They are formed as a part of our natural metabolism but also by certain environmental factors-including smoking, pesticides, pollution and radiation. Cell damage caused by free radicals appears to be a major contributor to aging and to degenerative diseases such as cancer, cardiovascular disease, cataracts, immune system decline and brain dysfunction [4, 5]. Overall, free radicals have been implicated in the pathogenesis of at least 50 diseases [6].

Reactive oxygen species (ROS) and reactive nitrogen species (RNS)

ROS and RNS formed *in vivo* such as superoxide anion, hydroxyl radical and hydrogen peroxide are highly reactive and potentially damaging transient chemical species. These are continuously produced during the body's metabolic reactions in small amount, as they are essential for energy supply, detoxification, chemical signaling and immune function. ROS are regulated by endogenous enzymes superoxide dismutase, glutathione peroxidase and catalase but due to overproduction of reactive species, induced by exposure to external oxidant substances or a failure in the defense mechanism, damage to cell structure, DNA, lipids and

Received: Oct 5, 2011; Revised: Dec 20, 2011; Accepted: Dec 25, 2011

*Corresponding Author

Sadaf Kalam
Centre for Scientific Research and Development (C.S.R.D.), People's Group,
Bhopal-462037, MP, India.

Tel: +91-9502403852
Email: sadaf2577@gmail.com

proteins [7] occur which increased risk of more than 30 different disease processes [8]. The most notorious among them being neurodegenerative conditions like Alzheimer's disease [9, 10], mild cognitive impairment [11] and Parkinson's disease [12]. Other diseases include highly disabling vascular pathologies like cardiovascular disease and cardiac failure [13], alcohol induced liver disease [14] and ulcerative colitis [15] and cancer caused by a complex of different causes, of which RNS/ROS is a component. Because ROS/RNS production is a naturally occurring process, a variety of enzymatic and non-enzymatic mechanisms have evolved to protect cells against these species [16]. The RNS, for example nitric oxide ($\text{NO}\cdot$) radical, is an abundant reactive radical which acts as an important oxidative biological signaling molecule in a large variety of physiological processes, including neurotransmission, blood pressure regulation, defense mechanisms, smooth muscle relaxation and immune regulation [17]. Examples of such species are listed in Table 1 [18].

Table 1. Examples of Reactive Oxygen Species (ROS)

The Radicals	The Non radicals
Superoxide ($\text{O}_2^{\cdot-}$)	Hydrogen peroxide (H_2O_2)
Hydroxyl ($\text{OH}\cdot$)	Hypochlorous acid (HOCl)
Peroxy, alkoxy ($\text{RO}_2\cdot$, $\text{RO}\cdot$)	Ozone (O_3)
Oxides of nitrogen ($\text{NO}\cdot$, $\text{NO}_2\cdot$)	Singlet oxygen ($^1\text{O}_2$)

Pathways leading to generation of free radicals

Free radicals/ROS/RNS are produced mainly from two important sources [19-21] in the biological system i.e. cellular metabolism like mitochondrial electron transport chain, endoplasmic reticulum oxidation, NADPH oxidase, xanthine oxidase, prostaglandin synthesis, reduced riboflavin, nitric oxide synthetase, reperfusion injury, cytochrome P_{450} , activated neutrophils and phagocytic cells and environmental sources like drugs, pesticides, transition metals, tobacco smoke, alcohol, radiations and high temperature. The other metabolic sources of free radicals include ionising radiation which causes lysis of water, reaction of transition metals with O_2 or H_2O_2 , and respiratory burst of macrophages [22].

Prooxidants and oxidative stress

Prooxidants are chemicals that induce oxidative stress either through ROS or inhibiting antioxidant systems. Normally, there is equilibrium between a free radical/ROS formation and endogenous antioxidant defense mechanisms, but if this balance is disturbed, it can produce oxidative stress [23, 24]. Oxidative stress is the shift towards the pro-oxidants in pro-oxidant: antioxidant balance that can occur due to increase in oxidative metabolism. It is characterized by a disturbance in the balance between ROS production on one hand and ROS removal and repair of damaged complex molecules (such as proteins or DNA) on the other [25]. Increased oxidative stress at the cellular level may occur due to exposure to alcohol, medications,

trauma, cold, infections, poor diet, toxins, radiation and strenuous exercise. Under certain conditions, such as acute or chronic alcohol exposure, ROS production is enhanced or the level or activity of antioxidants is reduced. Oxidative stress is associated with numerous deleterious consequences for the cell such as lipid peroxidation or even cell death.

Damage due to free radicals

Oxidative damage to DNA, proteins and other macromolecules has been implicated in the pathogenesis of a wide variety of diseases. Conditions associated with oxidative damage are heart diseases, cancer, pulmonary disorders, cataracts, diabetes, neurological disorders and aging [26]. Mitochondrial dysfunction may occur due to ROS produced as a consequence of increased oxidative stress and insufficient antioxidant defenses [27, 28]. ROS are potentially damaging transient chemical species which exert oxidative stress towards human cells rendering each cell to face about 10,000 oxidative hits per second [29, 30]. Scientists have exhaustively reviewed the effects of free radicals and antioxidants in normal physiological functions and human diseases [31]. There have been reports that the most hazardous hydroxyl radical reacts with all components of the DNA molecule, damaging both the purine and pyrimidine bases and also the deoxyribose backbone [32]. Not only DNA is being attacked by ROS but they also attack other cellular components involving polyunsaturated fatty acid residues of phospholipids, side chains of all amino acids residues of proteins, in particular cysteine and methionine residues [33, 34].

Antioxidants

Antioxidants are a broad group of compounds which constitute the first line of defense against free radical damage thus are essential for maintaining optimum health and well-being. They are protective agents, capable of stabilizing or deactivating free radicals before they attack cells. Being beneficial compounds, they control free radical formation naturally and help organisms to deal with oxidative stress caused by free radicals [35]. Exposure to free radicals increases the need for intake of antioxidants. The free radical exposures is increased due to pollution, cigarette smoke, drugs, illness, stress and even exercise. Diets supplemented with antioxidants are now-a-days recognized as an imperative way of protecting cells from adverse effects of free radicals. Identification of pharmacologically potential antioxidant compounds has increased tremendously as they exhibit no side effects for use in preventive medicine and food industry. Antioxidant compounds are present in vegetables, fruits and many natural beverages like tea. Balanced diets are naturally rich in antioxidants. Antioxidants display an array of benefits like they support kidney function, maintain good dental health, improve reproductive function, improve nervous system functioning, have anti-aging effect, protect liver, support immune system and improve defense power of the body. They also reduce obesity, offer protection against digestive disorders, maintain healthy vision and improve quality of sleep. Figure 1 depicts the relative functions of antioxidants.




Fig 1. Functions of antioxidants

Antioxidants are a family of compounds considered best to fight against a number of age-related problems such as Alzheimer's disease [36]. Antioxidants can be obtained from a host of fruits,

beverages, mushrooms etc. each doing some specific task in shielding the body from the harmful effects of the free radicals. Figure 2 demonstrates mechanism of action of antioxidants.




Fig 2. Mechanism of action of antioxidants

Considered to be wonder element, antioxidants are essential to good health and well-being as the concept of health promotion has become a legitimate part of health care. The ability to utilize oxygen has provided humans with the benefit of metabolizing fats, proteins and carbohydrates for energy [37]. Several studies have been directed towards the evaluation of antioxidant properties of several naturally occurring botanicals and herbs, potentially useful as nutraceutical ingredients [38].

Antioxidant in plant cells

Antioxidant in plant cells mainly include glutathione, ascorbate, tocopherol, proline, betaine and others which are also information rich redox buffers and important redox signaling components that interact with cellular compartments. As an unfortunate consequence of aerobic life for higher plants, ROS are formed by partial reduction of molecular oxygen. The enzymatic and non-enzymatic antioxidants in higher plant cells can protect their cells from oxidative damage by scavenging ROS. Thus, special attention is given to ROS and ROS-antioxidant interaction as a metabolic interface for different type of signals derived from metabolisms and from the changing environment [39]. Besides exacerbating cellular damage, ROS act as ubiquitous signal molecules in higher plants as well as a central component in stress responses [40].

Shielding effect by antioxidants

Humans have evolved a highly complicated antioxidant defense system to combat the damaging effects of free radicals. These includes both endogenous and exogenous components which function interactively and synergistically to neutralize free radicals [41]. These components are (i) Antioxidant enzymes (Superoxide Dismutase, Glutathione peroxidase and Catalase), (ii) Metal binding proteins (Ferritin, Lactoferrin, Albumin and Ceruloplasmin), (iii) Nutrient-derived antioxidants (Ascorbic acid, Tocopherols, Tocotrienols, Carotenoids, Glutathione and Lipoic acid), and (iv) Phytonutrients (Flavonoids, Phenolic acid, Stilbenes, Tannins and Carotenoids). The activity or amount of antioxidant is generally calculated as ORAC values i.e. Oxygen Radical Absorbance Capacity which is a measure of the ability of foods to subdue harmful oxygen free radicals that can damage human body. The ORAC values for two cups of black tea are equivalent to that of one glass red wine or seven glasses orange juice or consuming twenty glasses apple juice. Additionally, dark chocolates snack within a balanced diet can improve DNA resistance to oxidative stress in healthy subjects [42].

Antioxidants and diseases

Antioxidants are intimately involved in the prevention of cellular damage- the common pathway for cancer, aging, and a variety of diseases. Damage to DNA and other cellular organelles by free radicals occupies highest position in the onset and development of diseases. Some diseases have been mentioned below indicating free radicals as their precipitating factor.

Cardiovascular disease

Atherosclerosis and its complications most notably Coronary

Heart Disease (CHD) continues to be the major cause of premature death in the development world. Polyunsaturated fatty acid residues in lipoproteins have a chemical structure that makes them a particularly vulnerable target for free radical oxidation [43]. ROS-induced oxidative stress in cardiac and vascular myocytes has been linked with cardiovascular tissue injury [44]. Regardless of the direct evidence for a link between oxidative stress and cardiovascular disease, ROS-induced oxidative stress plays a role in various cardiovascular diseases such as atherosclerosis, ischemic heart disease, hypertension, cardiomyopathies, cardiac hypertrophy and congestive heart failure [45]. The major sources of oxidative stress in cardiovascular system include: (i) the enzymes xanthine oxidoreductase (XOR), (ii) NAD(P)H oxidase (multisubunit membrane complexes), (iii) NOS, (iv) the mitochondrial cytochromes and (v) hemoglobin [46, 47]. NOSs and hemoglobin are also principal sources of RNS, including NO• and SNOs (NO-modified cysteine thiols in amino acids, peptides, and proteins), which convey NO• bioactivity. Oxidative stress is associated with increased formation of ROS that modifies phospholipids and proteins leading to peroxidation and oxidation of thiol groups [48]. Studies have shown an inverse relationship between vitamin C intake and cardiovascular disease mortality in humans [49]. In hypercholesterolemic persons, supplementation with combination of vitamin E and slow-release vitamin C slows down atherosclerotic progression [50].

Human cancer

Under conditions of oxidative stress, abnormal expression of cellular genes can be induced by a wide variety of carcinogens [51]. The by-products of normal metabolism, which increase inflammation and exposure to exogenous sources like nitrogen oxide pollutants, smoking, certain drugs (e.g., acetaminophen, bleomycin) and radiation can induce cancer-causing mutations, lipids and proteins oxidation and alteration in signal-transduction pathways that enhance cancer risk [52]. Antioxidants also show promise in cancer therapy by their palliative action, reducing painful side effects associated with treatment [53]. The major goal of antioxidants in maintaining health is to protect normal cells from events that trigger or enhance cancer and to prevent long-term damage that may occur during cancer therapy and give rise to secondary malignancies in later years [54]. Many cancer patients who are undergoing therapy take antioxidant supplements in an effort to alleviate treatment toxicity and improve long term outcome. The results of epidemiological studies have suggested that consumption of diet rich in carotenoids was associated with reduction in breast cancer risk [55], and a reduced prostate cancer risk was seen with intake of foodstuff that are rich in lycopene [56]. Similarly, intake of allium vegetables (garlic, onions) containing high levels of antioxidant organosulfur compounds is linked with reduced gastrointestinal cancer risk [57] while polyphenols rich green tea has been considerably shown to reduce the risk of breast cancer and ovarian cancer in Asian women [58, 59]. Dietary antioxidants other than vitamin E, vitamin C and the carotenoids may also be of significance in the prevention of degenerative diseases and the maintenance of good health. The role of the antioxidant minerals in the etiology of cancer has been investigated and selenium is emerging as a major prophylactic factor against cancer [60].

Diabetes

Increased oxidative stress plays a major role in the development and advancement of diabetes and its complications [61, 62]. Generally, diabetes is accompanied by increased production of free radicals [63, 64] or impaired antioxidant defenses [65, 66]. Increased oxidative stress has been proposed to be one of the major causes of the hyperglycemia-induced trigger of diabetic complications. Diabetic Hyperglycemia in an organism stimulates ROS formation from a variety of sources. These sources include glucose autooxidation, polyol pathway and non-enzymatic glycation of proteins [67].

There have been found changes in oxidative stress biomarkers including superoxide dismutase, catalase, glutathione reductase, glutathione peroxidase, glutathione levels, vitamins, lipid peroxidation, nitrite concentration and nonenzymatic glycosylated proteins in diabetic individuals suffering from complications [68]. However, oxidative stress is relatively low in Non-insulin dependent Diabetes mellitus (NIDDM) when compared to insulin dependent Diabetes mellitus (IDDM) suggesting metabolic differences between the two types of diabetes [69]. Various studies show that the antioxidants can help in reducing oxidative stress and alleviate these diabetic complications [70]. Along with antioxidants, advanced glycation end products inhibitors, either introduced endogenously or exogenously, may counteract with the deleterious effects of the ROS/RNS and thus, helpful in prevention or treatment paradigms of this devastating disease [71].

Ischemia-reperfusion (IR) injury

IR injury activated when liver or other organs are briefly subjected to reduce blood supply followed by reperfusion, ultimately leading to cell damage and tissue failure. It has been revealed that ROS are generated during ischemia and reperfusion and may represent pivotal mediators of the resultant pathological complications [72]. Although, there is low oxygen tension during ischemia, moderate ROS generation is substantiated to occur most probably from a mitochondria source [73, 74]. Scientists presented the experimental and clinical evidence about the role of antioxidants in modulating hepatic ischemia-reperfusion injury [75]. The protective effects of antioxidants against brain I/R-induced injury may be mediated by down-regulation of nuclear factor-kappa B (NF- κ B) activity in rats [76]. A surplus of antioxidant agents against hepatic IR injury [77] and myocardial IR injury [78] has been tested *in vitro* and *in vivo* and the results emphasized the importance of antioxidative mechanism in cardiac and hepatic protection. Moreover, Ascorbate - a major antioxidant in brain seems to be helpful in scrutinizing chronic cerebral hypoperfusion [79] and resveratrol could protect intestinal tissue against IR injury with its potent antioxidant properties [80].

Alcoholic liver disorder (ALD)

Alcohol promotes the generation of ROS [81] and/or interferes with the body's normal defense mechanisms against free radicals. The ensuing alcohol-induced oxidative stress results in compromised antioxidant defense system, particularly in the patients of ALD [82]. Alcohol breakdown in the liver results in the formation of molecules whose further metabolism in the cell leads to ROS generation. Alcohol also stimulates the activity of enzymes called cytochrome

P₄₅₀s, and alters the level of certain metals in the body, thereby facilitating ROS production. This ROS production and resulting oxidative stress in liver cells play a major role in the development of ALD [83]. Nevertheless, numerous studies have found that administering antioxidants agents can prevent cells from the toxic effects of alcohol. For example, in the intragastric infusion model, the antioxidant vitamin E; the chemical Ebselen, which mimics the actions of glutathione peroxidase; the copper-zinc or manganese SODs; or a GSH precursor, all can significantly prevent or ameliorate alcohol's lethal effects in ALD patients [84-88].

Neurodegenerative disorders (NDD)

NDD are a diverse group of diseases of the nervous system, including the brain, spinal cord and peripheral nerves with different etiologies. Many are hereditary, some are secondary to toxic or metabolic processes, and others result from infections [89]. The brain is vulnerable to oxidative damage because of its high oxygen utilization, high content of oxidizable polyunsaturated fatty acids and the presence of redox-active metal ions [90]. The increased incidence of NDD such as Alzheimer's disease (AD), amyotrophic lateral sclerosis (ALS), cerebellar disorders, Parkinson's disease (PD), Huntington's disease (HD) and stroke in aged populations may be related to the accumulation of oxidative damage in neurons either primarily or secondarily [91]. Neuroprotective antioxidants can act as protective agent in NDD and are considered a promising approach to slowing the progression and limiting the extent of neuronal cell loss in these disorders [92, 93].

Alzheimer's disease (AD)

The brains of patients with AD exhibit a significant extent of oxidative damage associated with a marked accumulation of amyloid- β peptide (A β), the main constituent of senile plaques in brain, as well as deposition of neurofibrillary tangles and neurophil threads [94]. The cognitive function in such patients is inversely associated with systemic oxidative stress and vitamin E can be considered as an effective treatment of this disease [95].

Parkinson's disease (PD)

PD involves a selective loss of neurons in an area of the midbrain called the *substantia nigra* [96]. There is evidence that there are relatively high levels of basal oxidative stress in the *substantia nigra pars compacta* (SNc) in the normal brain, but this has been found to be increases in PD patients [97]. A majority of studies indicate the effect of oxidative stress that contributes to the cascade of events leading to dopamine cell degeneration in PD [98]. Supplementation with antioxidants may prevent or reduce the rate of progression of this disease such as L-dihydroxyphenylalanine (L-dopa), the agents used in the treatment of PD, combined with high levels of multiple antioxidants may improve the efficacy of L-dopa therapy [99].

Rheumatoid arthritis (RA)

Rheumatoid arthritis is a systemic autoimmune disease that causes chronic inflammation of the joints and tissue around the joints with infiltration of macrophages and activated T cells [100]. Cellular proliferation of the synoviocytes and neo-angiogenesis leads to

formation of pannus which destroys the articular cartilage and the bone [101]. Some studies provide evidences for the involvement of free radicals in the pathogenesis of rheumatoid arthritis [102-103]. The pathogenesis of this disease is linked predominantly with the formation of free radicals at the site of inflammation. Antioxidant supplementation or intake of natural dietary antioxidants possibly plays a protective and an adjuvant role in the treatment of RA [104].

Oxidative Damage to Nucleic Acid

Nucleic acids damage by oxygen free radicals results in producing many different types of lesions. These lesions can be grouped into strand breaks and base modification products [105]. Superoxide produced from the xanthine/ xanthine oxidase system has been shown to cause DNA strand breaks [106]. Hydroxyl free radicals react readily with nucleic acids, yielding many different kinds of products including strand breaks, which are probably the most frequent single lesion. These strand breaks are a result of the attack of the hydroxyl free radicals on the sugar portion, probably carbons 3' and 4', of the macromolecule. Strand breaks have important implications in terms of the development of pathological states because they must be repaired for the cell to function properly. Decrease in total antioxidant capacity of the aged rat plasma and myocardium shows that this decrease may be involved in the mechanisms of free radical- induced damage to DNA during aging [107]. Such oxidative damage to nucleic acids can be decline by supplementation with pure antioxidants or with foods rich in antioxidants [108].

Aging

The process of aging may be characterized as a progressive decline in the physiological functions of an organism with time and results in a decreased resistance to multiple forms of stress as well as an increased susceptibility to various diseases [109]. The free radical theory of aging was first introduced in 1956 by Denham Harman who proposed the concept of free radicals playing a role in the aging process [110]. This theory asserts that many of the changes, such as damage to DNA, protein cross-linking and other changes, that occur in the body with age has been attributed to free radicals. Over time, these damages accumulate and cause aging process. Intense research has been triggered into the field of role of free radicals in biological systems. Superoxide generated adventitiously by the mitochondrial respiratory chain can give rise to much more reactive radicals, resulting in random oxidation of all classes of macromolecules. However, it has become apparent that many downstream consequences of free radical damage can also be caused by processes not involving oxidation. Damage associated with lipid peroxidation and cellular damage by active oxygen species

contributes to the pathological changes attributed to aging. Scientists reported a decrease in the rate of free radical mediated lipid peroxidation on supplementation with vitamin E [111].

Longevity genes

The genetics of aging has made substantial strides in the past decade. This progress has been confined primarily to model organisms, such as filamentous fungi, yeast, nematodes, fruit flies, and mice, in which some thirty-five genes that determine life span have been cloned. These genes encode a wide array of cellular functions, indicating that there must be multiple mechanisms of aging. Nevertheless, some generalizations are already beginning to emerge. There are at least four broad physiological processes that play a role in aging: metabolic control, resistance to stress, gene dysregulation, and genetic stability [112].

Researchers proposed that p66^{shc} is part of a signal transduction pathway that regulates stress apoptotic responses and life span in mammals [113]. The SHC1 gene (named after its Src and Collagen homologous regions) is an important candidate gene for longevity in humans as it was found that mice in which p66SHC specific region of the SHC gene is deleted live 30% longer without apparent disease and these mice have lower levels of oxidative stress and apoptosis, both of which have been linked to old age survival in man [114]. Apoptosis and reactive oxygen species have been suggested to play a role in human longevity [115, 116]. Resveratrol, the first gene known to activate complement longevity, prolongs lifespan and retards the expression of age-dependent traits in a short-lived vertebrate [117]. Scientists have reported that the longevity of both invertebrates and vertebrates can be altered through genetic manipulation and pharmacological intervention. Most of these interventions involve alterations of one or more of the following: insulin/IGF-I signaling pathway, caloric intake, stress resistance and nuclear structure [118].

Phytochemicals

Since decades, plants have been recognized to contain many natural substances. Phytochemicals are natural chemical compounds found in plant-derived foods. These natural compounds are widely distributed and some times present in surprisingly higher concentration in plants exhibiting potent antioxidant activity. A wide range of phytochemicals, including flavonoids, carotenoids, polyphenols and organosulfur compounds are antioxidants and dietary intake of such antioxidants plays an important role in the protection of the human beings against free radicals and several other diseases like cardiovascular diseases and cancer [119]. Table 2 lists phytochemicals present in different vegetables and fruits [120].

Table 2. Phytochemicals present in different vegetables and fruits.

Food	Phytochemicals
Allium vegetables (Garlic, onions, chives, leeks)	Allyl sulfides
Cruciferous vegetables (broccoli, cauliflower, cabbage, Brussels sprouts, kale, turnips, kohlrabi)	Indoles/ glucinolates, Sulfaforaphane, Isothiocyanates/ thiocyanates/ thiols
Solanaceous vegetables (tomatoes, peppers, capsicum)	Lycopene, Solasodine, Capsaicin
Umbelliferous vegetables (carrots, celery, parsley, parsnips)	Carotenoids, Phthalides, Polyacetylenes
Composite plants (artichoke)	Silimarin
Citrus fruits (oranges, lemons, grapefruit)	Monoterpenes (limonene), Carotenoids, Tangeritin, Naringin
Other fruits (grapes, berries, cherries, apples, watermelon, pomegranate)	Ellagic acid, Phenols, Flavonoids,

Beans, grains, seeds (soybeans, oats, barley, brown rice, whole wheat, flax seed)	Flavonoids (isoflavones), Phytic acid Saponins
Herbs, spices (ginger, mint, rosemary, thyme, oregano, sage, basil, fennel, garlic, turmeric)	Gingerols, Zingerone, Flavonoids, Monoterpenes (limonene), Ferulic acid, Alicin, Curcumin
Licorice root, Green tea	Glycyrrhizin, Epigallocatechin gallate (EGCG)
Other plants, <i>Hypericum</i> sp., Rosemary, <i>Myrsine</i> sp., <i>Macuna</i> sp., Spinach, Maize	Hypericin, Carnosol, Embelin, Macunadine, Lutein, zeaxanthin

Many clinical and epidemiological studies show a connection between the antioxidant activity of the substances present in the diet and the prevention from such diseases as cardiovascular diseases or carcinogenesis [121, 122].

In traditional medicine and healing, many herbs and medicinal plants are one of the richest reservoirs of antioxidants. In many countries, screening studies were carried out for the comparison of antioxidant activities of medicinal plants typical for the respective country [123, 124]. Plants are known to produce secondary metabolites which possess antioxidant potential which represent a potential source of novel and lucrative compounds with antioxidant activity. Traditional herbal medicines thus constitute an important integral part of the Indian healthcare system. These include tomatoes, yams, carrot, leafy green vegetables, blue-berries, black-berries, cherries, *Ginko biloba*, garlic and green tea [125]. The antioxidant effects in plants are mainly due to the presence of phenolic compounds such as flavonoids, phenolic acids, tannins and phenolic diterpenes [126]. Several epidemiological studies suggest that plants rich in antioxidant play a protective role in health and against disease and their consumption lowered risk of cancer, heart disease, hypertension and stroke [127, 128]. Major groups of phytochemicals contributing to the total antioxidant capacity of plant include polyphenols and vitamins (C & E).

Phenolic compounds

Phenolic compounds of plants are hydroxylated derivatives of benzoic acid and cinnamic acids and have been reported to possess antioxidant effects [129]. Several reports indicate that the antioxidant potential of medicinal plants may be related to the concentration of their phenolic compounds which include phenolic acids, flavonoids, anthocyanins and tannins [130].

Flavonoids

Many studies suggest that dietary factors based on cereals, pulses, spices, dark green leafy vegetables (kale and spinach), citrus fruits, crude palm oil, soybean oil, cod liver oil, sprouts, peppers, whole grain, honey, walnuts and black tea can significantly increase the hepatic antioxidant enzymes and their supplementation reduces the risk of coronary heart disease effectively and safely particularly phenolic compounds like flavonoids. They improve endothelial function and inhibit platelet aggregation in humans [131]. Flavonoids regarded as “super antioxidants” scavenge free radicals associated with oxygen and iron; or by inhibiting oxidative enzymes. Over 4000 flavonoids have been found and fall in four different groups: flavones, flavanones, catechins, and anthocyanins. Certain fruits, flowers, roots, stems, tea, wine, grains and vegetables harbor high quantities of flavonoids. Being rich in flavonoids – apples protect against lung cancer and Blueberries improves memory in animals [132]. Isoflavones of soybeans seems to starve cancer cells and inhibit tumor growth. It also lower blood cholesterol and protect cardiac arteries.




Fig 3. Flavonoids

Bioflavonoids

Oligomeric Proanthocyanidins (Bioflavonoids) have the ability to attach to cells and their proteins for up to 72 hours protecting them from oxidation and free radical damage and replenish antioxidant effect of eye fluid, which help protect the eye against UV rays [133]. Grape seed, red wine, grape skin, peanut skin, and some white wines and fruits are rich in bioflavonoids.

Polyphenols and resveratrol

Polyphenols are another important group of phytochemicals which includes resveratrol (3,4',5-trihydroxystilbene) and protects against cancer by inhibiting cell growth and against heart disease by limiting clot formation. The richest natural sources of resveratrol are dark grape extracts and giant knotweed (*Polygonum cuspidatum*, a perennial shrub). Resveratrol and its related polyphenols also protect the cell's DNA, block cancer-causing chemicals and radiation and fight free radicals and inflammation. In one study, resveratrol along with quercetin and curcumin (from turmeric), emerged as the most powerful anticancer agents from 22 compounds that were subjected to a battery of tests, including their ability to suppress ornithine decarboxylase, scavenge free radicals and counteract carcinogens [134].




Fig 4. Resveratrol

Carotenoids

Carotenoids are a group of red, orange and yellow pigments found in plant foods, particularly fruits and vegetables. Carotenoids like β -carotene act as a precursor of vitamin A. Both *in vitro* and *in*

vivo studies show that β -carotene is an effective antioxidant [135]. It is one of the most powerful singlet oxygen quenchers as it can dissipate the energy of singlet oxygen, thus preventing this active molecule from generating free radicals. Its other antioxidant properties include the scavenging of free radicals. Unlike other nutrient antioxidants, β -carotene acts efficiently even at low oxygen pressure [136]. Besides β -carotene, other important dietary carotenoids include α -carotene, lycopene, lutein, zeaxanthin and β -

cryptoxanthin. Some studies have confirmed that lycopene has a powerful role as an antioxidant [137]. Tomatoes, with their abundant lycopene, defend against cancer by protecting DNA from oxidative damage [138]. Broccoli sprouts contain an abundance of the cancer-fighting phytochemical sulforaphane. Citrus fruits contains considerable amount of limonene which possess inhibitory activity against human gastric cancer [139]. Table 3 represents antioxidants and their respective color/ pigments in different veggies and fruits.

Table 3. Major antioxidants and their respective color/ pigments from different sources

Antioxidant	Color/Type	Sources
Beta-Carotene	Orange Colors	Sweet potatoes, carrots, squash, apricots, pumpkin, and mangoes
Lutein	Green Colors.	leafy vegetables/greens, spinach
Lycopene	Red Colors	Tomatoes, watermelon, guava, papaya, apricots, pink grapefruit.
Vitamin E	Fat Soluble	almonds, oils (corn and soybean), mangoes, nuts, and broccoli
Vitamin A	Fat Soluble	liver, sweet potatoes, carrots, milk, egg yolks and mozzarella cheese
Vitamin C	Water Soluble	Many fruits and vegetables, cereals, beef, poultry and fish.
Selenium (Mineral)	-	Seafood, poultry and meat products

Naturally occurring antioxidants in leafy vegetables and seeds, such as ascorbic acid, vitamin E and phenolic compounds, possess the ability to reduce the oxidative damage associated with many diseases, including cancer, cardiovascular disease, cataracts, atherosclerosis, diabetes, arthritis, immune deficiency diseases and aging [140, 141] and so they are important in the prevention of human diseases. Antioxidant compound may efficiently function as free radical scavengers, completers of pro-oxidant metals, reducing agents and quenchers of singlet oxygen formation. Antioxidants such as tocopherols and phenolic compounds are often added in oils and fatty acids to retard their auto-oxidation [142].

Medicinal herbs as antioxidants

Indian medicinal herbs were extensively investigated as vital sources of antioxidants. *Aerva lanata* exhibits 1,1-diphenyl-2-picrylhydrazyl (DPPH) scavenging properties [143] and are rich in flavonoids such as Kaempferol 3-rhamnoside and Kaempferol 3-rhamnolactoside [144]. Its inhibitory effect on microsomal cytochrome P₄₅₀ enzyme or on lipid peroxidation is due to the presence of alkaloids [145]. Similarly, the ability of *Amaranthus paniculatus* extract to act as a free radical scavenger or hydrogen donor was revealed by DPPH radical scavenging activity assay [146]. *Coccinia indica* leaf extract also possess an antioxidant activity which may be attributed its protective action on lipid peroxidation and to the enhancing effect on cellular antioxidant defense contributing to the protective action against oxidative damage [147]. Presence of polyphenolic compounds with antioxidant activity such as quercetin 3-glucuronide isoquercetin and rutin has been reported in seeds of *Coriandrum sativum* while confirmation through DPPH assay [148] and the presence of berberina and phenolic compounds was analyzed in *Coscinium fenestratum* [149]. *Evolvulus alsinoides* ethanolic extract and water infusion have antioxidant effects as depicted by the inhibition of lipid peroxidation and ABTS assay [150]. *Fagona cretica* is known to exhibit strong free radical scavenging properties against reactive oxygen [151]. *Gymnema montanum* also shows antioxidant activity due to the presence of ascorbic acid and α -tocopherol [152]. *Hygrophila auriculata* is an indigenous plant with free radical scavenging and lipid peroxidation activities and inhibits carbon tetrachloride induced oxidative stress and proliferative

response [153].

Various species of *Phyllanthus* are known to exhibit potent antioxidant activity. *P. amarus* is rich in polyphenols, tannins, quercetin, catechin and epigallocatechin as potential antioxidants [154]. The n-hexane extract of *P. maderaspatensis* exhibits *in vitro* radical scavenging activity and inhibition of lipid peroxidation [155]. Furthermore, methanolic and aqueous extract of *P. niruri* showed inhibition of membrane lipid peroxidation, scavenging of DPPH radical and inhibition of ROS *in vitro* [156] as well as it inhibit carbon tetrachloride induced formation of lipids peroxide in the liver of rats *in vivo* [157]. Hence, herbal medicines exhibit healing power and are one of the important cultural and traditional recipes for health care and sustainability. The antioxidant activities of these herbal plants and their extracts have a promising role in radical scavenging and in protecting cells from cellular damage.

Garlic

Garlic is rich in antioxidant phytochemicals that comprise organosulfur compounds as well as flavonoids such as allixin, which are capable of scavenging free radicals [158]. The antioxidant compounds present in aged, old garlic extract called "Kyolic" are known to contain radio-protective properties. This aged garlic extract is an odorless preparation made from organic garlic by a process of extraction and aging [159]. It is rich in antioxidant and more effective as an immunomodulating agent compared to fresh garlic [160]. Its major antioxidant compound is S-allyl cysteine, which is well effective in shielding cells from oxidant damage by radiation and anticancer drugs [161]. Garlic antioxidants have been also linked with blood cholesterol lowering properties, cardio-protective aspects, as well as anti-aging properties [162].

Curcumin

Various experimental studies confirm that Curcumin, a yellow pigment in *Curcuma longa*, acts as a potent antioxidant and radical scavenger [163]. Curcumin is also known to possess cancer-preventive antioxidant where it induces apoptosis in colon cancer cells and multiple myeloma cells [164] as well as ovarian cells [165]. It acts as a free radical scavenger, inhibiting lipid peroxidation [166]

and oxidative DNA damage and thought to play a vital role against an array of chronic pathological complications such as cancer, atherosclerosis and neurodegenerative diseases [167].

Silibinin

Silibinin is a powerful antioxidant being the product of Milk thistle (*Silbum marianum*) [168]. It is known to induce apoptosis in prostate cancer cells suggesting a potential role in the control of disease [169].

Herbal Tea

There is growing interest in drinking tea all over the world, which could be connected with antioxidative activity, fighting the harmful influence of environmentally generated free radicals. The analysis of DPPH• free radicals scavenging ability by green and black tea aqueous extracts was conducted and found that the green tea aqueous extract showed higher radicals scavenging efficiency than black tea extract [170]. Black tea contains lower levels of catechins and higher levels of tannins [171]. The best abilities of DPPH• free radicals scavenging was affirmed for (±)-catechin, quercitin and rutin, considerably higher than BHT and α-tocopherol.




Fig 5. Epigallocatechin Gallate (EGCG)

The major components of black tea include theaflavins (orange pigments) and thearubigins (red pigments). Green tea is rich in polyphenols and flavonoids and contains a characteristic Catechin-Epigallocatechin Gallate (EGCG) which is a superior antioxidant known to inhibit cancer and also is an anti-obese (calorie burning) compound. The antioxidant present in tea reduces cholesterol and risk of heart diseases and boosts immune system [172].

Pomegranate: The “Rising Superfruit”

Antioxidants found in fruits and vegetables are known to provide our bodies with many health benefits. Pomegranate (*Punica granatum*) is one such fruit that contains high levels of antioxidants including polyphenols, such as tannins and anthocyanins. The most abundant polyphenols and a major antioxidant compound present in pomegranate juice are hydrolyzable tannins called Punicalagins that have free radical scavenging properties in laboratory experiments [173]. Most therapeutically beneficial pomegranate constituents are ellagic acid, ellagitannins (including punicalagins), punicic acid, flavonoids, anthocyanidins, anthocyanins, and estrogenic flavonols and flavones [174]. Pomegranate extracts have been shown to scavenge free radicals and decrease macrophage oxidative stress and lipid peroxidation in animals [175] and increase plasma antioxidant capacity in elderly humans [176].




Fig 6. Ellagic Acid




Fig 7. Punicalagin

Mushrooms as a source of antioxidants

Mushrooms have been highly valued across the globe as both food and medicine since times immemorial. They are a group of fleshy macroscopic fungi which until recently as other fungi, were introduced in the plant kingdom because of cell wall and spores [177]. Edible mushrooms are found to contain a noticeable amount of total phenols and can therefore play a major role in exhibiting antioxidant potential. They can be used as easily accessible source of natural antioxidants and as a possible food supplement or as a therapeutic in pharmaceutical industry.

A naturally occurring metabolite is Ergothioneine (ERT) or 2-mercaptohistidine trimethylbetaine found in dietary mushrooms which is a unique powerful antioxidant and naturally occurring amino acid. *Grifola frondosa*, *Pleurotus ostreatus* and *Lentinula edodes* harbor highest amount of ERT. Since it scavenges strong oxidants such as hydroxy radicals, hypochlorite and peroxynitrite *in vitro*, and chelates redox-active bivalent cations such as Cu^{2+} , ERT has been considered to function primarily as an antioxidant within cells. *Ganoderma lucidum*, a medicinally potent mushroom is known to profoundly increase the catalase gene expression and enzyme activities in mouse liver [178]. Functionally catalases are related to peroxidases, both promote H_2O_2 oxidation by mechanisms involving ferryl intermediates [179]. Aqueous extracts from wild *Cordyceps sinensis* could scavenge hydroxyl radicals [180]. Similarly, *Cordyceps militaris* resisted the damage from superoxide anion mainly with superoxide dismutase and resisted the damage from hydrogen peroxide with both catalase and antioxidant compounds [181]. Several commercial mushrooms are reported to be effective against an array of diseases by virtue of their antioxidant potential. The antioxidant activity of *Pleurotus pulmonarius* (Fr.) was found to be due to the presence of phenols [182] and that of *Russula delica* could be credited to the presence of phenols and flavonoids [183].

Endogenous defense mechanisms

Body relies on several endogenous defense mechanisms to help protect against free radical-induced damage. Antioxidant enzymes are scavenger of reactive oxygen intermediates and are involved in many cellular defense systems [184]. Enzymes like glutathione peroxidase, catalase and superoxide dismutase (SOD)

metabolize oxidative toxic intermediates and necessitate micronutrient cofactors such as selenium, iron, copper, zinc, and manganese for optimum catalytic activity. During normal physiological process, reactive species are continuously formed but can be removed by antioxidant defense mechanism [185]. Table 4 depicts different endogenous antioxidants along with their functions [186].

Table 4. Antioxidant enzymes and human diseases

Disease	Specification	Main key enzyme/s
Allergy	Intolerance to aspirin	GPX
	Intolerance to other drugs	SOD
	Intolerance to some foods	GPX
	Reaction in skin tests	SOD
Cancer	Bowel	CAT, GPX, SOD
	Breast	GPX
	Colorectal	COX-2
	Kidney	CAT, GPX, SOD
	Leukemia	CAT, GPX, SOD
	Liver	CAT, GPX, SOD
	Skin	GPX
Cardiological and vessels injuries	Ischemia	SOD
	Atherosclerosis	SOD
Infectious disease	Arthritis	COX-2
	<i>Helicobacter pylori</i>	SOD
	Hepatitis	GPX
	HIV	GPX
	Influenza virus	CAT, GPX, SOD
	Chronic granulomatous disease	CAT
Genetic disorder	Down's syndrome	SOD
	Allergic encephalomyelitis	NOS
Neurodegenerative disease	Alzheimer's disease	SOD
	Amyotrophic lateral sclerosis	SOD
	Huntington's disease	SOD
	Parkinson's disease	GPX
	Prion disease	SOD
	Diabetes	CAT, SOD
Metabolic malfunction	Cataract	CAT, SOD

GPX: Glutathione peroxidase; SOD: Superoxide dismutase; CAT: Catalase; COX-2: Cyclooxygenase-2; NOS: Nitric oxide synthase.

The role of catalase in deterring tumor promotion and enhancing cell survival during periods of oxidative stress supports the antioxidant function of this enzyme [187]. Free radicals such as superoxide radical ($O_2^{\cdot-}$), Hydroxyl radical (OH^{\cdot}) and other reactive oxygen species (ROS) are associated with cellular necrosis and a variety of pathological conditions such as cancer, degenerative disease in neurons, hepatopathies, atherosclerosis and even aging [188]. The defense systems against free radical in human are a proof of the main role of antioxidant enzymes in blood cells detoxification, showing the coordinated enzymatic mechanism, and the interrelationships between all these enzymatic activities [189].

Synthetic antioxidants

Supplementation with antioxidant could represent an important therapeutic potential to minimize the damage caused by free radicals [190]. Scientists are attempting to develop novel synthetic antioxidants to retard the effects of free radical induced damage in different foodstuffs and human body cells [191]. Many such synthetic compounds have been evaluated for their efficacy as radical scavengers or for their other inhibitory effects. Among them, only four synthetic antioxidants are extensively used in food preservation; namely, butylated hydroxyanisole (BHA), butylated

hydroxytoluene (BHT), propyl gallate (PG) and tert-butylhydroquinone (TBHQ) [192, 193]. These synthetic antioxidants are broadly used in food industry but now-a-days use of such compounds in foods has been decreased due to their possible action as promoters of carcinogenesis [194].

CONCLUSION

Free radicals and related species have been implicated in the etiology of myriad of major diseases. Recent research is directed towards formulation of strategies to protect cardinal tissues and organs against oxidative damage induced by free radicals and related chemical entities. The antioxidant defense systems are importance in maintaining normal cellular physiology and fighting against diseases. There are endogenous protective antioxidant mechanisms and dietary antioxidants that are intimately involved in the prevention of cellular damage caused by free radicals. The traditional Indian diet, spices and medicinal plants are rich sources of natural antioxidants. Attention is focused on ROS/RNS- linked pathogenesis of cancer, cardiovascular disease, atherosclerosis, hypertension, ischemia/reperfusion injury, diabetes, neurodegenerative diseases, rheumatoid arthritis and aging. An array of natural products are endowed with antioxidant potential, thus

it is a time, after periods of booming research on oxidants and antioxidants to critically reflect on this related subject. Diet rich in antioxidants such as fresh and unprocessed foods, raw fruit and vegetables and other antioxidant supplements are now being renowned as an important means of reducing the toxic effects imposed by free radicals. Therefore, the endogenous antioxidant enzymes and cofactors as well as the extensive assortment of natural antioxidants are concurrently presenting an enormous possibility in improving free radical protection and in the treatment of human diseases.

REFERENCES

- [1] Sies, H. 1993. Strategies of antioxidant defense. *Eur. J. Biochem.* 215(2):213-219.
- [2] Cheeseman, K. H., T. F. Slater. 1993. An introduction to free radical biochemistry. In: K. H. Cheeseman and T. F. Slater (Eds.), *Free radical in medicine*, New York: Churchill Livingstone, pp. 481-493.
- [3] Blasiak, J., A. Sikora, A. Czechowska and J. Drzewoski. 2003. Free radical scavengers can modulate the DNA damaging action of alloxan. *Acta Biochim. Pol.* 50(1):205-210.
- [4] Tandon, V. R., S. Verma, J. B. Singh and A. Mahajan. 2005. Antioxidants and cardiovascular health. *J. K. Sci.* 7(2):61-64.
- [5] Sies, H., W. Stahl and A. R. Sundquist. 1992. Antioxidant functions of vitamins: Vitamin E and C, beta-carotene and other carotenoids. *Ann. N. Y. Acad. Sci.* 669:07-20.
- [6] Halliwell, B. 1994. Free radicals, antioxidants and human disease: Curiosity, cause or consequences. *Lancet.* 344(8924):721-724.
- [7] Valko, M., C. J. Rhodes, J. Moncol, M. Izakovic and M. Mazur. 2006. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem. Biol. Interact.* 160(1):01-40.
- [8] Aruoma, O. I. 1998. Free radicals, oxidative stress and antioxidants in human health and disease. *J. Am. Oil Chem. Soc.* 75(2):199-212.
- [9] Smith, M. A., G. Perry, P. L. Richey, L. M. Sayrec, V. E. Anderson, M. F. Beal and N. Kowall. 1996. Oxidative damage in Alzheimer's. *Nature.* 382:120-121.
- [10] Smith, M. A., C. A. Rottkamp, A. Nunomura, A. K. Raina and G. Perry. 2000. Oxidative stress in Alzheimer's disease. *Biochim. Biophys. Acta.* 1502(1):139-144.
- [11] Guidi, I., D. Galimberti, S. Lonati, C. Novembrino, F. Bamonti, M. Tiriticco, C. Fenoglio, E. Venturelli, P. Baron, N. Bresolin and E. Scarpini. 2006. Oxidative imbalance in patients with mild cognitive impairment and Alzheimer's disease. *Neurobiol. Aging.* 27(2):262-269.
- [12] Bolton, J. L., M. A. Trush, T. M. Penning, G. Dryhurst and T. J. Monks. 2000. Role of quinones in toxicology. *Chem. Res. Toxicol.* 13(3):135-160.
- [13] Jha, P., M. Flather, E. Lonn, M. Farkouh and S. Yusuf. 1995. The antioxidant vitamins and cardiovascular disease: A critical review of epidemiological and clinical trial data. *Ann. Intern. Med.* 123(11):860-872.
- [14] Arteel, G. E. 2003. Oxidants and antioxidants in alcohol-induced liver disease. *Gastroenterology.* 124(3):778-790.
- [15] Ramakrishna, B. S., R. Varghese, S. Jayakumar, M. Mathan and K. A. Balasubramanian. 1997. Circulating antioxidants in ulcerative colitis and their relationship to disease severity and activity. *J. Gastroenterol. Hepatol.* 12(7):490-494.
- [16] Yu, B. P. 1994. Cellular defenses against damage from reactive oxygen species. *Physiol. Rev.* 74(1):139-162.
- [17] Bergendi, L., L. Benes, Z. Durackova and M. Ferencik. 1999. Chemistry, physiology and pathology of free radicals. *Life Sci.* 65(18-19):1865-1874.
- [18] Mimic-Oka, J., D. V. Simic and T. P. Simic. 1999. Free radicals in cardiovascular diseases. *Facta Univ. Med. Biol.* 6(1):11-22.
- [19] Sen, S., R. Chakraborty, C. Sridhar, Y. S. R. Reddy and B. De. 2010. Free radicals, antioxidants, diseases and phytomedicines: Current status and future prospect. *Int. J. Pharm. Sci.* 3(1):91-100.
- [20] Kelly, F. J., C. Dunster and I. Mudway. 2003. Air pollution and the elderly: Oxidant/antioxidant issues worth consideration. *Eur. Respir. J. Suppl.* 40:70s-75s.
- [21] Mitchell, R. N., R. S. Cotran. 2003. Cell injury, adaptation, and death. In: V. Kumar, R. S. Cotran and S. L. Robbins (Eds.), *Basic Pathology*. 7th edition. New Delhi: Harcourt (India) Pvt. Ltd, pp. 03-33.
- [22] Yamamoto, K. and R. B. Johnston. 1984. Dissociation of phagocytosis from stimulation of the oxidative metabolic burst in macrophages. *J. Exp. Med.* 159(2):405-416.
- [23] Murray, R. K., D. K. Granner, P. A. Mayes, V. W. Rodwell. 2000. Structure and function of lipid soluble vitamins. In: R. K. Murray, D. K. Granner, P. A. Mayes and V. W. Rodwell (Eds.), *Harper's biochemistry*, New York: McGraw-Hill, pp. 642-661.
- [24] Young, I. S. and J. V. Woodside. 2001. Antioxidants in health and disease. *J. Clin. Pathol.* 54(3):176-186.
- [25] Halliwell, B. 1999. Antioxidant defense mechanisms: From the beginning to the end. *Free Radic. Res.* 31(4):261-272.
- [26] Wierusz-Wysocka, B., H. Wysocki, H. Byks, D. Zozulinska, A. Wykretowicz and M. Kazmierczak. 1995. Metabolic control quality and free radical activity in diabetic patients. *Diabetes Res. Clin. Pract.* 27(3):193-197.
- [27] Askew, E. W. 2002. Work at high altitude and oxidative stress: Antioxidant nutrients. *Toxicology.* 180(2):107-119.
- [28] Bailey, S. M. and C. C. Cunningham. 2002. Contribution of mitochondria to oxidative stress associated with alcoholic liver disease. *Free Radic. Biol. Med.* 32(1):11-16.
- [29] Sundaresan, M., Z. X. Yu, V. J. Ferrans, D. J. Sulciner, J. S. Gutkind, K. Irani, P. J. Goldschmidt-Clermont and T. Finkel. 1996. Regulation of reactive oxygen species generation in fibroblast by Rac1. *Biochem. J.* 318(Pt 2):379-382.
- [30] Marangoni, A., S. Accardo, R. Aldini, M. Guardigli, F. Cavrini, V. Sambri, M. Montagnani, A. Roda and R. Cevenini. 2006. Production of reactive oxygen species and expression of inducible nitric oxide synthase in rat isolated Kupffer cells stimulated by *Leptospira interrogans* and *Borrelia burgdorferi*. *World J. Gastroenterol.* 12(19):3077-3081.

- [31] Valko, M., D. Leibfritz, J. Moncol, M. T. Cronin, M. Mazur and J. Telser. 2007. Free radicals and antioxidants in normal physiological functions and human disease. *Int. J. Biochem. Cell Biol.* 39(1):44-84.
- [32] Halliwell, B., J. M. C. Gutteridge. 1999. Oxidative stress and antioxidant protection: Some special cases. In: B. Halliwell and J. M. C. Gutteridge (Eds.), *Free Radicals in Biology and Medicine*, 3rd edition. Oxford: Clarendon Press, pp. 530-533.
- [33] Siems, W. G., T. Grune and H. Esterbauer. 1995. 4-hydroxynonenal formation during ischemia and reperfusion of rat small intestine. *Life Sci.* 57(8):785-789.
- [34] Stadtman, E. R. 2004. Role of oxidant species in aging. *Curr. Med. Chem.* 11(9):1105-1112.
- [35] Bahorun, T., M. A. Soobrattee, V. Luximon-Ramma and O. I. Aruoma. 2006. Free radicals and antioxidants in cardiovascular health and disease. *Internet J. Med. Update.* 1(2):25-41.
- [36] Hajieva, P. and C. Behl. 2006. Antioxidants as a potential therapy against age-related neurodegenerative diseases: Amyloid beta toxicity and alzheimer's disease. *Curr. Pharm. Des.* 12(6):699-704.
- [37] Percival, M. 1998. Antioxidants. *Clin. Nutr. Insights.* 1:01-04.
- [38] Chopra, R. N., S. L. Nayar and I. C. Chopra. 1996. *Glossary of Indian Medicinal Plants*, National Institutes of Science Communication, New Delhi, India.
- [39] Shao, H. B., L. Y. Chu, Z. H. Lu and C. M. Kang. 2007. Primary antioxidant free radical scavenging and redox signaling pathways in higher plant cells. *Int. J. Biol. Sci.* 4(1):08-14.
- [40] Halliwell, B. 2006. Reactive species and antioxidants: Redox biology is a fundamental theme of aerobic life. *Plant Physiol.* 141(2):312-322.
- [41] Liebler, D. C. 1993. The role of metabolism in the antioxidant function of Vitamin E. *Crit. Rev. Toxicol.* 23(2):147-169.
- [42] Spadafranca, A., C. M. Conesa, S. Sirini and G. Testolin. 2010. Effect of dark chocolate on plasma epicatechin levels, DNA resistance to oxidative stress and total antioxidant activity in healthy subjects. *Br. J. Nutr.* 103(7):1008-1014.
- [43] Maxwell, S. R. J. and G. Y. H. Lip. 1997. Free radicals and antioxidants in cardiovascular disease. *Br. J. Clin. Pharmacol.* 44(4):307-317.
- [44] Dhalla, N. S., R. M. Temsah and T. Netticadan. 2000. Role of oxidative stress in cardiovascular diseases. *J. Hypertens.* 18(6):655-673.
- [45] Kukreja, R. C. and M. L. Hess. 1992. The oxygen free-radical system: From equations through membrane-protein interactions to cardiovascular injury and protection. *Cardiovasc. Res.* 26(7):641-655.
- [46] Berry, C. E. and J. M. Hare. 2004. Xanthine oxidoreductase in the cardiovascular disease: Molecular mechanisms and pathophysiological implications. *J. Physiol.* 555(Pt 3):589-606.
- [47] Hare, J. M. and J. S. Stamler. 2005. NO/redox disequilibrium in the failing heart and cardiovascular system. *J. Clin. Invest.* 115(3):509-517.
- [48] Molavi, B. and J. L. Mehta. 2004. Oxidative stress in cardiovascular disease: Molecular basis of its deleterious effects, its detection, and therapeutic considerations. *Curr. Opin. Cardiol.* 19(5):488-493.
- [49] Simon, J.A. 1992. Vitamin C and cardiovascular disease: A review. *J. Am. Coll. Nutr.* 11:107-125.
- [50] Salonen, R. M., K. Nyyssonen, J. Kaikkonen, E. Porkkala-Sarataho, S. Voutilainen, T. H. Rissanen, T. P. Tuomainen, V. P. Valkonen, U. Ristonmaa, H. M. Lakka, M. Vanharanta, J. T. Salonen and H. E. Poulsen. 2003. Six-year effect of combined vitamin C and E supplementation on atherosclerotic progression: The antioxidant supplementation in atherosclerosis prevention (ASAP) study. *Circulation.* 107(7):947-953.
- [51] Borek, C. 1991. Free radical processes in multistage carcinogenesis. *Free Radic. Res. Commun.* 12-13(Pt 2):745-750.
- [52] Gutteridge, J. M. 1993. Free radicals in disease processes: A compilation of cause and consequence. *Free Radic. Res. Commun.* 19(3):141-158.
- [53] Kennedy, M., K. Brunuinga, E. A. Mutlu, J. Losurdo, S. Choudhary and A. Keshavarian. 2001. Successful and sustained treatment of chronic radiation proctitis with antioxidant vitamins E and C. *Am. J. Gastroenterol.* 96(4):1080-1084.
- [54] Coia, L. R. and D. J. Moyland. 1998. *Introduction to Clinical Radiation Oncology*, Medical Physics Publishing, Madison, WI.
- [55] Huang, J. P., M. Zhang, C. D. Holman and X. Xie. 2007. Dietary carotenoids and risk of breast cancer in Chinese women. *Asia Pac. J. Clin. Nutr.* 16(Suppl 1):437-442.
- [56] Nkondjock, A., P. Ghadirian, K. C. Johnson, D. Krewski and Canadian cancer registries epidemiology research group. 2005. Dietary intake of lycopene is associated with reduced pancreatic cancer risk. *J. Nutr.* 135(3):592-597.
- [57] Greenwald, P., C. K. Clifford and J. A. Milner. 2001. Diet and cancer prevention. *Eur. J. Cancer.* 37(8):948-965.
- [58] Wu, A. H., M. C. Yu, C. C. Tseng, J. Hankin and M. C. Pike. 2003. Green tea and risk of breast cancer in Asian Americans. *Int. J. Cancer.* 106(4):574-579.
- [59] Zhang, M., C. W. Binns and A. H. Lee. 2002. Tea consumption and ovarian cancer risk: A case control study in China. *Cancer Epidemiol. Biomarkers Prev.* 11(8):713-718.
- [60] Li, H., M. J. Stampfer, E. L. Giovannucci, J. S. Morris, W. C. Willett, J. M. Gaziano and J. Ma. 2004. A prospective study of plasma selenium levels and prostate cancer risk. *J. Natl. Cancer. Inst.* 96(9):696-703.
- [61] Giacco, F. and M. Brownlee. 2010. Oxidative stress and diabetic complications. *Circulation Res.* 107:1058-1070.
- [62] Ceriello, A. 2000. Oxidative stress and glycemic regulation. *Metabolism.* 49(2 Suppl 1):27-29.
- [63] Chang, K. C., S. Y. Chung, W. S. Chong, J. S. Suh, S. H. Kim, H. K. Noh, B. W. Seong, H. J. Ko and K. W. Chun. 1993. Possible superoxide radical-induced alteration of vascular reactivity in aortas from streptozotocin-treated rats. *J. Pharmacol. Exp. Ther.* 266(2):992-1000.

- [64] Young, I. S., S. Tate, J. H. Lightbody, D. McMaster and E. R. Trimble. 1995. The effects of desferrioxamine and ascorbate on oxidative stress in the streptozotocin diabetic rat. *Free Radic. Biol. Med.* 18(5):833-840.
- [65] Saxena, A. K., P. Srivastava, R. K. Kale and N. Z. Baquer. 1993. Impaired antioxidant status in diabetic rat liver: Effect of vanadate. *Biochem. Pharmacol.* 45(3):539-542.
- [66] McLennan, S. V., S. Heffernan, L. Wright, C. Rae, E. Fisher, D. K. Yue and J. R. Turtle. 1991. Changes in hepatic glutathione metabolism in diabetes. *Diabetes.* 40(3):344-348.
- [67] Johansen, J. S., A. K. Harris, D. J. Rychly and A. Ergul. 2005. Oxidative stress and the use of antioxidants in diabetes: Linking basic science to clinical practice. *Cardiovasc. Diabetol.* 4(1):05.
- [68] Maritim, A. C., R. A. Sanders and J. B. Watkins III. 2003. Diabetes, oxidative stress, and antioxidants: A review. *J. Biochem. Mol. Toxicol.* 17(1):24-38.
- [69] Moussa, S.A. 2008. Oxidative stress in Diabetes mellitus. *Romanian J. Biophys.* 18(3):225-236.
- [70] Rahimi, R., S. Nikfar, B. Larijani and M. Abdollahi. 2005. A review on the role of antioxidants in the management of diabetes and its complications. *Biomed. Pharmacother.* 59(7):365-373.
- [71] Reddy, V. P., X. Zhu, G. Perry and M. A. Smith. 2009. Oxidative stress in diabetes and alzheimer's disease. *J. Alzheimers Dis.* 16(4):763-774.
- [72] Galaris, D., A. Barbouti and P. Korantzopoulos. 2006. Oxidative stress in hepatic ischemia-reperfusion injury: The role of antioxidants and iron chelating compounds. *Curr. Pharm. Des.* 12(23):2875-2890.
- [73] Lombardi, V., L. Valko, S. Stolc, M. Valko, O. Ondrejickova, L. Horakova, J. Placek and A. Troncone. 1998. Free radicals in rabbit spinal cord ischemia: Electron spin resonance spectroscopy and correlation with SOD activity. *Cell Mol. Neurobiol.* 18(4):399-412.
- [74] Becker, L. B. 2004. New concepts in reactive oxygen species and cardiovascular reperfusion physiology. *Cardiovasc. Res.* 61(3):461-470.
- [75] Glantzounis, G. K., H. J. Salacinski, W. Yang, B. R. Davidson and A. M. Seifalian. 2005. The contemporary role of antioxidant therapy in attenuating liver ischemia-reperfusion injury: A review. *Liver transpl.* 11(9):1031-1047.
- [76] Shen, W. H., C. Y. Zhang and G. Y. Zhang. 2003. Antioxidants attenuate reperfusion injury after global brain ischemia through inhibiting nuclear factor-kappa B activity in rats. *Acta Pharmacol. Sin.* 24(11):1125-1130.
- [77] Korontzi, M. I., A. Papalois, I. Kouerinis, A. C. Lazaris, G. Theodoropoulos and G. Zografos. 2010. Redox state and the potential role of antioxidant compounds in liver ischemia/reperfusion injury. *Int. J. Med. Med. Sci.* 2(7):200-209.
- [78] Dhalla, N. S., A. B. Elmoselhi, T. Hata and N. Makino. 2000. Status of myocardial antioxidants in ischemia-reperfusion injury. *Cardiovasc. Res.* 47(3):446-456.
- [79] Kasparova, S., V. Brezova, M. Valko, J. Horecky, V. Mlynarik, T. Liptaj, O. Vancova, O. Ulicna and D. Dobrota. 2005. Study of the oxidative stress in a rat model of chronic brain hypoperfusion. *Neurochem. Int.* 46(8):601-611.
- [80] Ozkan, O. V., M. F. Yuzbasioglu, H. Ciralik, E. B. Kurutas, Z. Yonden, M. Aydin, E. Bulbuloglu, E. Semerci, M. Goksu, Y. Atli, V. Bakan and N. Duran. 2009. Resveratrol, a natural antioxidant, attenuates intestinal ischemia/reperfusion injury in rats. *Tohoku J. Exp. Med.* 218(3):251-258.
- [81] Halliwell, B. and J. M. Gutteridge. 1990. The antioxidant of human extra-cellular fluids. *Arch. Biochem. Biophys.* 280(1):01-08.
- [82] Gupta, S., R. Pandey, R. Katyal, H. K. Aggarwal, R. P. Aggarwal and S. K. Aggarwal. 2005. Lipid peroxide levels and antioxidant status in alcoholic liver disease. *Indian J. Clin. Biochem.* 20(1):67-71.
- [83] Wu, D. and A. I. Cederbaum. 2003. Alcohol, oxidative stress and free radical damage. *Alcohol Res. Health.* 27(4):277-284.
- [84] Nanji, A. A., E. K. Yang, F. Fogt, S. M. Sadrzadeh and A. J. Dannenberg. 1996. Medium chain triglycerides and vitamin E reduce severity of established experimental alcoholic liver disease. *J. Pharmacol. Exp. Ther.* 277(3):1694-1700.
- [85] Kono, H., G. E. Arteel, I. Rusyn, H. Sies and R. G. Thurman. 2001. Ebselen prevents early alcohol-induced liver injury in rats. *Free Radic. Biol. Med.* 30(4):403-411.
- [86] Wheeler, M. D., H. Kono, M. Yin, I. Rusyn, M. Froh, H. D. Connor, R. P. Mason, R. J. Samulski and R. G. Thurman. 2001. Delivery of the Cu/Zn-superoxide dismutase gene with adenovirus reduces early alcohol-induced liver injury in rats. *Gastroenterology.* 12(5):1241-1250.
- [87] Wheeler, M. D., M. Nakagami, B. U. Bradford, T. Uesugi, R. P. Mason, H. D. Connor, A. Dikalova, M. Kadiiska and R. G. Thurman. 2001. Overexpression of manganese superoxide dismutase prevents alcohol-induced liver injury in the rat. *J. Biol. Chem.* 276(39):36664-36672.
- [88] Iimuro, Y., B. U. Bradford, S. Yamashina, I. Rusyn, M. Nakagami, N. Enomoto, H. Kono, W. Frey, D. Forman, D. Brenner and R. G. Thurman. 2000. The glutathione precursor L-2-oxothiazolidine-4-carboxylic acid protects against liver injury due to chronic enteral ethanol exposure in the rat. *Hepatology.* 31(2):391-398.
- [89] Singh, R. P., S. Sharad and S. Kapur. 2004. Free radicals and oxidative stress in neurodegenerative diseases: Relevance of dietary antioxidants. *J. Indian Acad. Clin. Med.* 5(3):218-225.
- [90] Markesbery, W. R. 1997. Oxidative stress hypothesis in Alzheimer's disease. *Free Radic. Biol. Med.* 23(1):134-147.
- [91] Rao, A. V. and B. Balachandran. 2002. Role of oxidative stress and antioxidants in neurodegenerative diseases. *Nutr. Neurosci.* 5(5):291-309.
- [92] Moosmann, B. and C. Behl. 2002. Antioxidants as treatment for neurodegenerative disorders. *Expert. Opin. Investig. Drugs.* 11(10):1407-1435.
- [93] Dugan, L. L., E. G. Lovett, K. L. Quick, J. Lotharius, T. T. Lin and K. L. O'Malley. 2001. Fullerene-based antioxidants and neurodegenerative disorders. *Parkinsonism Relat. Disord.* 7(3):243-246.
- [94] Butterfield, D. A., A. Castegna, C. M. Lauderback and J. Drake. 2002. Evidence that amyloid beta-peptide-induced lipid peroxidation

- and its sequelae in Alzheimer's disease brain contribute to neuronal death. *Neurobiol. Aging*. 23(5):655-664.
- [95] Vina, J., A. Lloret, R. Orti and D. Alonso. 2004. Molecular bases of the treatment of Alzheimer's disease with antioxidants: Prevention of oxidative stress. *Mol. Aspects Med.* 25(1-2):117-123.
- [96] Sayre, L. M., M. A. Smith and G. Perry. 2001. Chemistry and biochemistry of oxidative stress in neurodegenerative disease. *Curr. Med. Chem.* 8(7):721-738.
- [97] Jomova, K., D. Vondrakova, M. Lawson and M. Valko. 2010. Metals, oxidative stress and neurodegenerative disorders. *Mol. Cell Biochem.* 345(1-2):91-104.
- [98] Tretter, L., I. Sipos and V. Adam-Vizi. 2004. Initiation of neuronal damage by complex I deficiency and oxidative stress in Parkinson's disease. *Neurochem. Res.* 29(3):569-577.
- [99] Prasad, K. N., W. C. Cole and B. Kumar. 1999. Multiple antioxidants in the prevention and treatment of Parkinson's disease. *J. Am. Coll. Nutr.* 18(5):413-423.
- [100] Bauerova, K. and A. Bezek. 1999. Role of reactive oxygen and nitrogen species in etiopathogenesis of rheumatoid arthritis. *Gen. Physiol. Biophys.* 18:15-20.
- [101] Edwards, C. R. W. and I. A. D. Bouchier. 1994. Davidson's principles and practice of medicine, 16th edition, Chirchill Livingstone, UK.
- [102] Karatas, F., I. Ozates, H. Canatan, I. Halifeoglu, M. Karatepe and R. Colak. 2003. Antioxidant status and lipid peroxidation in patients with rheumatoid arthritis. *Indian J. Med. Res.* 118:178-181.
- [103] Kamanli, A., M. Naziroglu, N. Aydile and C. Hacieviyagil. 2004. Plasma lipid peroxidation and antioxidant levels in patients with rheumatoid arthritis. *Cell Biochem. Funct.* 22(1):53-57.
- [104] Mahajan, A. and V. R. Tandon. 2004. Antioxidants and rheumatoid arthritis. *J. Indian Rheumatol. Assoc.* 12:139-142.
- [105] Floyd, R. A., J. E. Schneider. 1990. Hydroxyl free radical damage to DNA. In: C. Vigo-Pelfrey (Ed.), *Membrane lipid oxidation*, Vol. 1, Boca Raton, Florida: C.R.C. press, pp. 69-90.
- [106] Brawn, K. and I. Fridovich. 1981. DNA strand scission by enzymically generated oxygen radicals. *Arch. Biochem. Biophys.* 206(2):414-419.
- [107] Siononova, M., Z. Tatarkova, Z. Durackova, D. Dobrota, J. Lehotsky, T. Matakova and P. Kaplan. 2007. Relationship between antioxidant potential and oxidative damage to lipids, proteins and DNA in aged rats. *Physiol. Res.* 56(6):757-764.
- [108] Collins, A. R. and E. Horvathova. 2001. Oxidative DNA damage, antioxidants and DNA repair: Applications of the comet assay. *Biochem. Soc. Trans.* 29(Pt2):337-341.
- [109] Kregal, K. C. and H. J. Zhang. 2007. An integrated view of oxidative stress in aging: Basic mechanisms, functional effects and pathological considerations. *Am. J. Physiol.* 292(1):18-36.
- [110] Harman, D. 1956. Aging: A theory based on free-radical and radiation-chemistry. *J. Gerontol.* 11(3):298-300.
- [111] Wagner, B. A., G. R. Buettner and C. P. Burns. 1996. Vitamin E slows the rate of free radical mediated lipid peroxidation in cells. *Arch. Biochem. Biophys.* 334(2):261-267.
- [112] Jazwinski, S. M. 2000. Aging and longevity genes. *Acta Biochim. Pol.* 47(2):269-279.
- [113] Migliaccio, E., M. Giorgio, S. Mele, G. Pelicci, P. Reboldi, P. P. Pandolfi, L. Lanfrancione and P. G. Pelicci. 1999. The p66shc adaptor protein controls oxidative stress response and life span in mammals. *Nature*. 402 (6759):309-313.
- [114] Mooijaart, S. P., D. van Heemst, J. Schreuder, S. van Gerwen, M. Beekman, B. W. Brandt, P. E. Slagboom and R. G. Westendorp; 'Long Life' study group. 2004. Variation in the SHC1 gene and longevity in humans. *Exp. Gerontol.* 39(2):263-268.
- [115] Nemoto, S. and T. Finkel. 2002. Redox regulation of forkhead proteins through a p66shc-dependent signaling pathway. *Science*. 295 (5564):2450-2452.
- [116] Campisi, J. 2003. Cancer and aging: Rival demons. *Nat. Rev. Cancer*. 3(5):339-349.
- [117] Valenzano, D. R., E. Terzibasi, T. Genade, A. Cattaneo, L. Domenici and A. Cellarino. 2006. Resveratrol prolongs lifespan and retards the onset of age-related markers in a short-lived vertebrate. *Curr. Biol.* 16(3):296-300.
- [118] Warner, H. R. 2005. Longevity genes: From primitive organisms to humans. *Mech. Ageing Dev.* 126(2):235-242.
- [119] Bao, Y. and R. Fenwick. 2004. *Phytochemicals in Health and Disease*, Marcell Dekker Inc, New York.
- [120] Ayoola, G. A., A. D. Folawewo, S. A. Adesegun, O. O. Abioro, A. A. Adepoju-Bello and H. A. B. Coker. 2008. Phytochemical and antioxidant screening of some plants of apocynaceae from South West Nigeria. *African J. Plant Sci.* 2(9):124-128.
- [121] Hughes, D. A. 2000. Dietary antioxidants of human immune function. *Nutr. Bull.* 25(1):35-41.
- [122] Lindsay, D. G. and S. B. Astley. 2002. European research on the functional effects of dietary antioxidants. *Mol. Aspects Med.* 23(1-3):01-38.
- [123] Dastmalchi, K., H. J. Damien Dorman, I. Laakso and R. Hiltunen. 2007. Chemical composition and antioxidative activity of Moldavian balm (*Dracocephalum moldavica* L.) extracts. *Food Sci. Technol.* 40(9):1655-1663.
- [124] Proestos, C., I. S. Bozaris, G. J. E. Nychas and M. Komaitis. 2006. Analysis of flavonoids and phenolic acids in Greek aromatic plants: Investigation of their antioxidant capacity and antimicrobial activity. *Food Chem.* 95(4):664-671.
- [125] Wong, P. Y. Y. and D. D. Kitts. 2006. Studies on the dual antioxidant and antibacterial properties of parsley (*Petroselinum crispum*) and cilantro (*Coriandrum sativum*) extracts. *Food Chem.* 97(3):505-515.
- [126] Polterait, O. 1997. Antioxidants and free-radical scavengers of natural origin. *Current Org. Chem.* 01:415-440.
- [127] Vinson, J. A., X. Su, L. Zubik and P. Bose. 2001. Phenol antioxidant quantity and quality in foods: fruits. *J. Agric. Food Chem.* 49(11):5315-5321.
- [128] Wolfe, K., X. Wu and R. H. Liu. 2003. Antioxidant activity of apple peels. *J. Agric. Food Chem.* 51(3):609-614.

- [129] Muanda, F., D. Kone, A. Dicko, R. Soulimani and C. Younos. 2009. Phytochemical composition and antioxidant capacity of three Malian medicinal plant parts. *Evid. Based Complement. Alternat. Med.* 1-8.
- [130] Djeridane, A., M. Yousfi, B. Nadjemi, D. Boutassouna, P. Stocker and N. Vidal. 2006. Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food Chem.* 97(4):654-660.
- [131] Vita, J. A. 2005. Polyphenols and cardiovascular disease: Effects on endothelial and platelet function. *Am. J. Clin. Nutr.* 81(1 Suppl):292S-297S.
- [132] Hernandez, I., L. Alegre, F. van Breusegem and S. Munne-Bosch. 2009. How relevant are flavonoids as antioxidants in plants. *Trends Plant Sci.* 14(3):125-132.
- [133] Narayana, K. R., M. S. Reddy, M. R. Chaluvadi and D. R. Krishna. 2001. Bioflavonoids classification, pharmacological, biochemical effects and therapeutic potential. *Indian J. Pharmacol.* 33(1):02-16.
- [134] Li, Z. D., Q. Y. Ma and C. A. Wang. 2006. Effect of resveratrol on pancreatic oxygen free radicals in rats with severe acute pancreatitis. *World J. Gastroenterol.* 12(1):137-140.
- [135] Krinsky, N. I. 1993. Actions of carotenoids in biological systems. *Annu. Rev. Nutr.* 13:561-587.
- [136] Tsuchihashi, H., M. Kigoshi, M. Iwatsuki and E. Niki. 1995. Action of β -carotene as an antioxidant against lipid peroxidation. *Arch. Biochem. Biophys.* 323(1):137-147.
- [137] Rao, A. V. and S. Agarwal. 2000. Role of antioxidant lycopene in cancer and heart disease. *J. Am. Coll. Nutr.* 19(5):563-569.
- [138] Porrini, M. and P. Riso. 2000. Lymphocyte lycopene concentration and DNA protection from oxidative damage is increased in women after a short period of tomato consumption. *J. Nutr.* 130(2):189-192.
- [139] Lu, X. G., L. B. Zhan, B. A. Feng, M. Y. Qu, L. H. Yu and J. H. Xie. 2004. Inhibition of growth and metastasis of human gastric cancer implanted in nude mice by d-limonene. *World J. Gastroenterol.* 10(14):2140-2144.
- [140] Lee, K. C., A. E. Mitchell and T. Shibamoto. 2000. Determination of antioxidant properties of aroma extracts from various beans. *J. Agric. Food. Chem.* 48(10):4817-4820.
- [141] Middleton, E. Jr., C. Kandaswami and T. C. Theoharides. 2000. The effects of plant flavonoids on mammalian cells: Implication for inflammation, heart disease and cancer. *Pharmacol. Rev.* 52(4):673-751.
- [142] Choe, E. and D.B. Min. 2006. Mechanisms and factors for edible oil oxidation. *Compr. Rev. Food Sci. F.* 5(4):169-186.
- [143] Shirwaikar, A., D. Issac and S. Malini. 2004. Effect of *Aerva lanata* on cisplatin and gentamicin models of acute renal failure. *J. Ethnopharmacol.* 90(1):81-86.
- [144] Afaq, S. H., S. Tajuddin and R. Afridi. 1991. Bisheri booti (*Aerva lanata*) some lesser known uses and pharmacognosy. *Ethnobotany.* 5:37-40.
- [145] Nevin, K. G. and P. L. Vijayammal. 2005. Effect of *Aerva lanata* against hepatotoxicity of carbon tetrachloride in rats. *Environ. Toxicol. Pharmacol.* 20(3):471-477.
- [146] Amin, I., Y. Norazaidah and K. I. E. Hainida. 2006. Antioxidant activity and phenolic content of raw and balanced *Amaranthus* species. *Food Chem.* 94(1):47-52.
- [147] Venkateswaran, S. and L. Pari. 2003. Effect of *Coccinia indica* leaves on antioxidant status in streptozotocin-induced diabetic rats. *J. Ethnopharmacol.* 84(2-3):163-168.
- [148] Wangensteen, H., A. B. Samuelsen and K. E. Malterud. 2004. Antioxidant activity in extracts from Coriander. *Food Chem.* 88(2):293-297.
- [149] Punitha, I. S., K. Rajendran, A. Shirwaikar and A. Shirwaikar. 2005. Alcoholic stem extract of *Coscinium fenestratum* regulates carbohydrate metabolism and improves antioxidant status in streptozotocin-nicotinamide induced diabetic rats. *Evid. Based Complement. Alternat. Med.* 2(3):375-381.
- [150] Auddy, B., M. Ferreira, F. Blasina, L. Lafon, F. Arredondo, F. Dajas, P. C. Tripathi, T. Seal and B. Mukherjee. 2003. Screening of antioxidant activity of three Indian medicinal plants, traditionally used for the management of neurodegenerative disease. *J. Ethnopharmacol.* 84(2-3):131-138.
- [151] Rawal, A., M. Muddeshwar and S. Biswas. 2004. Effect of *Rubia cordifolia*, *Fagonia cretica* Linn and *Tinospora cordifolia* on free radical generation and lipid peroxidation during oxygen-glucose deprivation in rat hippocampal slices. *Biochem. Biophys. Res. Commun.* 324(2):588-596.
- [152] Ananthan, R., C. Baskar, V. NarmathaBai, L. Pari, M. Latha and K. M. Ramkumar. 2003. Antidiabetic effect of *Gymnema montanum* leaves: Effect on lipid peroxidation induced oxidative stress in experimental diabetes. *Pharmacol. Res.* 48(6):551-556.
- [153] Ahmed, S., A. Rahman, M. Mathur and S. Sultana. 2001. Anti-tumor promoting activity of *Asteracantha longifolia* against experimental hepatocarcinogenesis in rats. *Food Chem. Toxicol.* 39(1):19-28.
- [154] Ali, S. S., N. Kasoju, A. Luthra, A. Singh, H. Sharanbasava, A. Sahu and U. Bora. 2008. Indian medicinal herbs as source of antioxidants. *Food Res. Int.* 41(1):01-15.
- [155] Asha, V. V., S. Akhila, P. J. Wills and A. Subramoniam. 2004. Further studies on the antihepatotoxic activity of *Phyllanthus maderaspatensis* Linn. *J. Ethnopharmacol.* 92(1):67-70.
- [156] Chatterjee, M., K. Sarkar and P. C. Sil. 2006. Herbal (*Phyllanthus niruri*) protein isolate protects liver from nimesulide induced oxidative stress. *Pathophysiology.* 13(2):95-102.
- [157] Harish, R. and T. Shivanandappa. 2006. Antioxidant activity and hepatoprotective potential of *Phyllanthus niruri*. *Food Chem.* 95(2):180-185.
- [158] Nishino, H., A. Nishino, J. Takayasu, A. Iwashima, Y. Itakura, Y. Kodera, H. Matsuura and T. Fuwa. 1990. Antitumor promoting activity of allixin, a stress compound produced by garlic. *Cancer J.* 3(1):20-21.
- [159] Amagase, H., B. L. Petesch, H. Matsuura, S. Kasuga and Y. Itakura. 2001. Intake of garlic and its bioactive compounds. *J. Nutr.* 131(3s):955S-962S.

- [160] Kyo, E., N. Uda, S. Kasuga and Y. Itakura. 2001. Immunomodulatory effects of aged garlic extract. *J. Nutr.* 131(3s):1075S-1079S.
- [161] Borek, C. 2001. Antioxidant health effects of aged garlic extract. *J. Nutr.* 131(3s):1010S-1015S.
- [162] Bagchi, K. and S. Puri. 1998. Free radicals and antioxidants in health and disease. *East. Mediterr. Health J.* 4(2):350-360.
- [163] Ak, T. and I. Gulcin. 2008. Antioxidant and radical scavenging properties of curcumin. *Chem. Biol. Interact.* 174(1):27-37.
- [164] Chauhan, D. P. 2002. Chemotherapeutic potential of curcumin for colorectal cancer. *Curr. Pharm. Des.* 8(19):1695-1706.
- [165] Zheng, L., Q. Tong and C. Wu. 2004. Growth inhibitory effects of curcumin on ovary cells and its mechanism. *J. Huazhong. Univ. Sci. Technol. Med. Sci.* 24(1):55-58.
- [166] Shukla, P. K., V. K. Khanna, M. Y. Khan and R. C. Srimal. 2003. Protective effect of curcumin against lead neurotoxicity in rat. *Hum. Exp. Toxicol.* 22(12):653-658.
- [167] Menon, V. P. and A. R. Sudheer. 2007. Antioxidant and anti-inflammatory properties of curcumin. *Adv. Exp. Med. Biol.* 595:105-125.
- [168] Schonfeld von, J., B. Weisbrod and M. K. Muller. 1997. Silibinin, a plant extract with antioxidant and membrane stabilizing properties, protects exocrine pancreas from cyclosporin A toxicity. *Cell. Mol. Life Sci.* 53(11-12):917-920.
- [169] Tyagi, A., N. Bahtia, M. S. Condon, M. C. Bosland, C. Agarwal and R. Agarwal. 2002. Antiproliferative and apoptotic effects of silibinin in rat prostate cancer cells. *Prostate.* 53(3):211-217.
- [170] Yokozawa, T., E. Dong, T. Nakagawa, H. Kashiwagi, H. Nakagawa, S. Takeuchi and H. Y. Chung. 1998. *In vitro* and *in vivo* studies on the radical-scavenging activity of tea. *J. Agric. Food Chem.* 46(6):2143-2150.
- [171] Arts, M. J., G. R. Haenen, L. C. Wilms, S. A. Beetstra, C. G. Heijnen, H. P. Voss and A. Bast. 2002. Interactions between flavonoids and proteins: Effect on the total antioxidant capacity. *J. Agric. Food Chem.* 50(5):1184-1187.
- [172] Gramza, A., K. Pawlak-Lemanska, J. Korczak, E. Wsowicz and M. Rudzinska. 2005. Tea extracts as free radical scavengers. *Pol. J. Environ. Stud.* 14(6):861-867.
- [173] Kulkarni, A. P., S. M. Aradhya and S. Divakar. 2004. Isolation and identification of a radical scavenging antioxidant - punicalagin from pith and carpellary membrane of pomegranate fruit. *Food Chem.* 87(4):551-557.
- [174] Murthy, K. N. C., G. K. Jayaprakash and R. P. Singh. 2002. Studies on antioxidant activity of pomegranate (*Punica granatum*) peel extract using *in vivo* models. *J. Agric. Food Chem.* 50(17):4791-4795.
- [175] Rosenblat, M., N. Volkova, R. Coleman and M. Aviram. 2006. Pomegranate byproduct administration to apolipoprotein e-deficient mice attenuates atherosclerosis development as a result of decreased macrophage oxidative stress and reduced cellular uptake of oxidized low-density lipoprotein. *J. Agric. Food Chem.* 54(5):1928-1935.
- [176] Guo, C., J. Wei, J. Yang, J. Xu, W. Pang and Y. Jiang. 2008. Pomegranate juice is potentially better than apple juice in improving antioxidant function in elderly subjects. *Nutr. Res.* 28(2):72-77.
- [177] Badole, S. L., P. A. Thakurdesai and S. L. Bodhankar. 2008. Antioxidant activity of aqueous extracts of *Pleurotus pulmonarius* (Fries) Quel-Champ. *Pharmacologyonline.* 2:26-41.
- [178] Park, J., B. R. Lee, L. H. Jin, C. K. Kim, K. S. Choi, J. H. Bahn, K. S. Lee, H. Y. Kwon, H. W. Chang, N. I. Baek, E. H. Lee, J. H. Kang, S. W. Cho and S. Y. Choi. 2001. The stimulatory effects of *Ganoderma lucidum* and *Phellinus linteus* on the antioxidant enzyme catalase. *J. Biochem. Mol. Biol.* 34(2):144-149.
- [179] Dawson, J. H. 1988. Probing structure-function relation in heme-containing oxygenases and peroxidases. *Science.* 240(4851):433-439.
- [180] Zheng, L. P., L. W. Gao, J. Q. Zhou, Y. H. Sima and J. W. Wang. 2008. Antioxidant activity of aqueous extract of a *Tolypocladium* sp. fungus isolated from wild *Cordyceps sinensis*. *Afr. J. Biotechnol.* 7(17):3004-3010.
- [181] Gu, Y., Z. Wang and Q. Yuan. 2006. The varieties of antioxidant activity of *Cordyceps militaris* during the submerged fermentation. *Electro. J. Biol.* 2(2):30-33.
- [182] Ajith, T. A. and K. K. Janardhanan. 2007. Indian medicinal mushrooms as a source of antioxidant and antitumor agents. *J. Clin. Biochem. Nutr.* 40(3):157-162.
- [183] Turkoglu, A., M. E. Duru and N. Mercan. 2007. Antioxidant and antimicrobial activity of *Russula delica* Fr: An edible wild mushroom. *Eurasian J. Anal. Chem.* 2(1):54-67.
- [184] Kumar, A., R. Sivakanesan and S. Singh. 2008. Oxidative stress, endogenous antioxidant and ischemia-modified albumin in normolipidemic acute myocardial infarction patients. *J. Health Sci.* 54(4):482-487.
- [185] Rizzo, A. M., P. Berselli, S. Zava, G. Montorfano, M. Negroni, P. Corsetto and B. Berra. 2010. Endogenous antioxidants and radical scavengers. *Adv. Exp. Med. Biol.* 698:52-67.
- [186] Mates, J. M. and F. Sanchez-Jimenez. 1999. Antioxidant enzymes and their implications in pathophysiologic processes. *Front. Biosci.* 4:339-345.
- [187] Lee, H. Y., W. S. Eum, D. W. Kim, B. R. Lee, C. S. Yoon, S. H. Jang, H. S. Choi, S. H. Choi, N. I. Baek, J. H. Kang, T. C. Kang, M. H. Won, S. W. Cho, K. S. Lee, J. Park and S. Y. Choi. 2003. Isolation and identification of an antioxidant enzyme catalase stimulatory compound from *Ganoderma lucidum*. *J. Biochem. Mol. Biol.* 36(5):450-455.
- [188] Pryor, W. A. 1986. Oxy-radicals and related species: Their information, lifetimes and reactions. *Annu. Rev. Physiol.* 48:657-667.
- [189] Grazioli, V., R. Schiavo, E. Casari, F. Marzatico, R. Rodriguez y Baena and P. Gaetani. 1998. Antioxidant enzymatic activities and lipid peroxidation in cultured human chondrocytes from vertebral plate cartilage. *FEBS Lett.* 431(2):149-153.
- [190] Raza, S. A., A. Rashid, J. William, S. Najaf and M. Arshad. 2009. Effect of synthetic antioxidant on shelf life of locally manufactured butter known as Makhan in Pakistan. *Biharean Biologist.* 3(2):161-162.

- [191] Hussain, H. H., G. Babic, T. Durst, J. S. Wright, M. Flueraru, A. Chichirau, and L. L. Chepelev. 2003. Development of novel antioxidants: Design, synthesis, and reactivity. *J. Org. Chem.* 68(18):7023-7032.
- [192] Shahidi, F. 2000. Antioxidants in food and food antioxidants. *Nahrung.* 44(3):158-163.
- [193] Buck, D. F. and M. K. Edwards. 1997. Antioxidants to prolong shelf-life. *Food Technol. Int.* 29-33.
- [194] Shahidi, F., Y. Zhong. 2005. Antioxidants: Regulatory Status. In: F. Shahidi (Ed.), *Bailey's Industrial Oil and Fat Products*, 6th Edition, 6 Volume Set, John Wiley & Sons, Inc., pp. 491-512.