

# Occurrence and extracellular enzyme potential of Actinomycetes of a thermotolerant, northern region of Karnataka, India

Sreenivasa Nayaka and Vidyasagar G M

Department of Botany, Gulbarga University, Gulbarga-585106, Karnataka, India.

## Abstract

Occurrence and extracellular enzyme activities of a thermotolerant actinomycetes isolated from thermal spring of compost pit area, hair, feathers dumping places and municipal wastes of Gulbarga and Bellary (Northern part of Karnataka) and tested their degradability against hair and feather. A total of 67 actinomycetes were isolated, out of these 57 actinomycetes screened exhibited amylase, protease, chitinase, cellulase and keratinase activity determined in the petriplate using on starch casein agar (SCA) and nutrient agar media (NA). Out of 57 actinomycetes, 50 isolates showed amylase, 54 protease, 50 chitinase, 52 cellulase and 41 isolates showed keratinase activities respectively. The Maximum keratinase activity shows VSAC-12, this study shows that the isolates were found to be capable using keratin substrates as sole source of carbon and energy for further investigation.

**Keywords:** Actinomycetes, Extracellular enzymes, Keratinase

## INTRODUCTION

Actinomycetes play an important role in decomposition of plant and other materials especially in the degradation of complex and relatively recalcitrant polymers. They degrade lignin, cellulose and lignocellulose. There is evidence that actinomycetes are involved in the degradation of many other naturally occurring polymers in soils such as, hemicellulose, pectin, keratin, chitin and fungal cell wall materials (Ali and Royman, 1984). Among the microorganisms, actinomycetes gained special importance due to their capacity to produce bioactive secondary metabolites and enzymes. They are known from many habitats: Soil, compost, sludge, freshwater, marine and atmosphere (Lacey, 1973; Jensen, 1995; Dhevendaran and Annie, 1999; Seng *et al.*, 1999). Actinomycetes, at one time were considered to be microbiological curiosities of no great economic importance. Now it has become the subject of intensive search for sources of new biologically active compounds. Actinomycetes have always been regarded as a strange group of prokaryotes with respect to the diversity of their morphology and of their metabolic products such as, antibiotics, antitumor agents, enzymes, enzyme inhibitors and growth promoting substances (Lechevalier and Lechevalier, 1967). They have been found in all ecological niches such as, soil, water (marine and fresh), compost, extreme environments, marshy places etc. Among them soil has the greatest population density. Extreme environments such as, alkaline, acidic, high temperature, high pressure, complex ecological system, polluted environments, rhizospheres, can be looked for screening new actinomycetes (Gray and Williams, 1975; Goodfellow, 1983).

Most of the interest in actinomycetes group of microorganisms lies in their ability to produce a diverse group of bioactive compounds, having application in medicinal, industries and agricultural fields. Many genera, notably *Streptomyces* produce commercially important antibiotics (Goodfellow and O'Donnell, 1994) and an array of other secondary metabolites; these include aliphatic alcohols, lactones, biogenic sulphides, ketones, esters, thioesters, furanones and isoprenoids (Gerber, 1983; Jachymova *et al.*, 2002). Actinomycetes also produce secondary metabolites that show bioactivities other than antibiotics such as, enzyme inhibitors, immunosuppressors, phytotoxins and pesticides (Berdy, 1995; Park *et al.*, 2002). We report here the occurrence and enzymatic activities of actinomycetes isolated from a thermal spring situated at the North Karnataka region.

## MATERIAL AND METHODS

### Collection of soil samples

The soil samples were collected from different places like, compost pit area, hair and feather dumping places and municipal wastes of Gulbarga and Bellary (Northern part of Karnataka). Top layer of the soil was removed to a depth of about 8-10 cm with a clean spade and then using clean stainless steel scoop and plastic spoon. About 50 g of soil sample was collected in sterile polythene bags and the bags were labeled with date, place of collection and the sample number.

### Isolation of Actinomycetes

The standard serial dilution plate culture method (Nakeeb and Lechevalier, 1963) was employed to isolate the pure culture of actinomycetes. Adequate serial dilution ( $10^{-1}$ – $10^{-5}$ ) was prepared from the enriched samples. 0.1 ml of the sample from the respective dilution was plated on starch casein agar. The inoculated plates were incubated at 30°C for one week. The growth of actinomycetes was observed on the medium at regular interval of 24 h. After isolation of the colonies in pure culture on the slants, they were preserved in refrigerator.

Received: Oct 17, 2012; Revised: Nov 02, 2012; Accepted: Dec 28, 2012.

\*Corresponding Author

Sreenivasa Nayaka

Department of Botany, Gulbarga University, Gulbarga-585106, Karnataka, India.

Email: [nayaka\\_s@gmail.com](mailto:nayaka_s@gmail.com)

## Morphological characterization

A morphological study including substrate mycelium, aerial mycelium, sporulation and pigmentation status of the actinomycetes (Williams and Wellington, 1980), after their adequate growth on starch casein agar was carried out as per the procedure prescribed in Goodfellow (1989) Bergey's manual of Systematic Bacteriology. The growth characters including aerial mycelium, pigmentation and growth activity of all the 67 isolates were studied. The typical colonies of actinomycetes were re-inoculated on starch casein agar in which cover slips were placed in inclined position at an angle, 45° and incubated for 3 days. After the growth of actinomycetes, the cover slip was removed and observed for the substrate mycelium and aerial mycelium (Shinobu, 1958; Kawato and Shinobu, 1959).

## Screening of Extracellular Enzymes

The selected 62 actinomycete isolates strains were screened for the production of different extracellular enzymes namely, amylase, protease, chitinase, cellulase, and keratinase. Amylase activity was assayed using agar diffusion method by growing the isolates on nutrient agar (peptone, 5 g; yeast extract, 1.5 g; beef extract, 1.5 g; NaCl, 5 g; agar, 16 g; distilled water 1000 ml; pH 7.2) supplemented with 1% iodine in 2% KI. The clear zone developed around the colony was observed and inhibition diameter was measured. Protease activity was assayed by growing the isolates on the nutrient agar medium supplemented with 1% casein as a carbon source. After incubation, the plates were flooded with 0.1 ml of hydrochloric

acid. The formation of clear zone around the colony was observed and measured. The Chitinase activity was determined by growing the isolates on colloidal chitin agar medium (CCA). After incubation the clear zone around the colonies was considered as positive for chitinase production and using scale. The cellulase activity was determined by growing the isolates on the medium containing (NaNO<sub>3</sub> 2 g; K<sub>2</sub>HPO<sub>4</sub>, 1 g; MgSO<sub>4</sub> 7H<sub>2</sub>O 0.05 g; KCl, 0.05 g; FeSO<sub>4</sub> 7H<sub>2</sub>O 0.01 g; Agar 16 g; distilled water 1000 ml; pH 6.8), supplemented with 1% CMC-Na salt as carbon source. After incubation, the plates were flooded with 5ml of 1% KI. The formation of a clear zone around the colony against reddish brown background was observed and measured using scale. The keratinase activity was assayed by growing the isolates on NA (peptone, 5 g; yeast extract, 1.5 g; beef extract, 1.5 g; NaCl, 5 g; agar, 16 g; distilled water 1000 ml; pH 7.2) medium with 4% keratin as carbon source. After the incubation, the plates were flooded with 5 ml of 1% HgCl<sub>2</sub>. The formation of a clear zone around the colony against white precipitate was seen and the diameter was measured using scale.

## RESULT

### Screening of Isolates for Extracellular Enzymes

A total of 67 thermotolerant actinomycetes were isolated (Plate-I). The qualitative activity of five different enzymes viz., amylase, protease, chitinase, cellulase and keratinase were determined in the petriplate using SCA and NA medium. Out of 57 actinomycete isolates, 50 isolates showed amylase activity, 54 isolates showed protease, 50 isolates showed chitinase, 52 isolates showed cellulase and 41 isolates showed keratinase activity (Table-1).

Table 1. Enzyme activity in thermotolerant actinomycetes isolates from soils, composts pits of Gulbarga and Bellary District

Organisms	Zone of digestion of substrate (mm)				
	Amylase	Protease	Chitinase	Cellulase	Keratinase
VSAC-1	44.7	22.3	34.3	19.0	16.3
VSAC-2	31.6	31.0	24.3	20.0	15.3
VSAC-3	47.6	15.3	17.0	32.3	22.0
VSAC-4	34.3	18.6	24.3	0	19.3
VSAC-5	59.6	39.6	29.6	35.0	44.3
VSAC-6	29.3	41.6	32.0	25.6	0
VSAC-7	39.6	22.3	45.6	28.3	0
VSAC-8	26.3	28.6	49.0	0	0
VSAC-9	21.3	50.6	32.6	19.3	0
VSAC-10	35.6	26.3	28.6	45.6	33.0
VSAC-11	41.6	29.0	20.0	32.0	18.0
<b>VSAC-12</b>	<b>81.3</b>	<b>74.3</b>	<b>78.0</b>	<b>69.0</b>	<b>82.6</b>
VSAC-13	63.6	53.2	68.4	56.0	63.0
VSAC-14	41.3	29.6	32.3	19.3	21.0
VSAC-15	32.3	24.3	0	21.0	15.0
VSAC-16	24.6	38.3	41.0	0	26.0
VSAC-17	75.3	41.1	18.0	0	0
VSAC-18	0	18.3	21.6	26.3	0
VSAC-19	0	22.0	35.6	19.0	0
VSAC-20	0	26.3	16.0	45.6	32.0
VSAC-21	32.3	54.3	48.0	47.3	44.0
VSAC-22	14.3	0	22.0	32.0	51.0
VSAC-23	27.6	0	19.3	24.0	0
VSAC-24	54.6	36.3	41.3	41.0	19.0
VSAC-25	41.3	31.6	54.0	51.6	49.0
VSAC-26	48.3	28.3	43.3	39.6	28.0
VSAC-27	37.6	42.3	27.0	32.5	15.0
VSAC-28	18.6	34.3	20.6	0	0
VSAC-29	21.3	18.0	38.3	45.0	32.6
VSAC-30	0	14.6	0	25.6	0

VSAC-31	61.6	39.6	49.0	19.3	44.0
VSAC-32	22.6	25.6	35.6	36.3	31.0
VSAC-33	29.3	21.3	29.3	34.3	22.0
VSAC-34	59.6	51.6	16.0	28.0	48.0
VSAC-35	46.6	19.6	19.6	50.3	0
VSAC-36	29.3	23.0	36.3	29.3	0
VSAC-37	35.3	31.6	51.6	49.6	39.0
VSAC-38	34.6	27.3	0	21.0	19.3
VSAC-39	29.6	19.5	22.0	35.3	25.9
VSAC-40	41.6	16.3	19.0	45.0	32.0
VSAC-41	35.3	27.6	26.3	27.3	32.6
VSAC-42	39.6	0	21.0	41.0	26.3
VSAC-43	0	18.3	32.0	21.9	0
VSAC-44	61.6	51.3	54.5	51.0	53.0
VSAC-45	59.3	44.3	19.0	21.3	0
VSAC-46	28.3	39.0	0	0	19.0
VSAC-47	22.6	25.3	49.6	50.3	28.3
VSAC-48	44.3	18.6	0	29.3	0
VSAC-49	64.8	12.0	0	19.0	32.0
VSAC-50	37.3	21.3	32.0	25.3	19.6
VSAC-51	0	41.9	45.0	39.3	24.0
VSAC-52	44.6	53.6	39.0	49.6	18.0
VSAC-53	49.6	38.6	18.3	26.3	14.0
VSAC-54	58.3	51.0	50.0	52.0	59.0
VSAC-55	27.6	39.3	0	22.0	0
VSAC-56	44.6	34.3	16.6	32.0	16.2
VSAC-57	0	19.6	25.3	29.3	21.0

Isolates on Starch Casein Agar Media, Mean of 3 values.

Maximum amylase activity was seen in isolate VSAC-12 followed by VSAC-13, VSAC-17, VSAC-24, VSAC-31, VSAC-34, VSAC-44, VSAC-45, VSAC-49 and VSAC-54 and lowest activity in isolate VSAC-22. No activity was observed in VSAC-18, VSAC-19, VSAC-20, VSAC-30, VSAC-43, VSAC-51 and VSAC-57 isolates. Highest protease activity was observed in isolate VSAC-12 followed by VSAC-13, VSAC-21, VSAC-34, VSAC-44, VSAC-52 and VSAC-54 and lowest activity in VSAC-30. No activity was observed in VSAC-22, VSAC-23 and VSAC-42 isolate. Similarly, the highest chitinase activity was seen in isolate VSAC-12 followed by VSAC-13, VSAC-25, VSAC-37, VSAC-44 and VSAC-54 and lowest in VSAC-20. However, no activity was observed in VSAC-15, VSAC-30, VSAC-38, VSAC-46, VSAC-48 and VSAC-55 isolates. The cellulase activity was maximum in isolate VSAC-12 followed by VSAC-13, VSAC-25, VSAC-35, VSAC-44 and VSAC-47. No activity was recorded in isolates VSAC-4, VSAC-8, VSAC-16, VSAC-17, VSAC-28 and VSAC-46. The keratinase activity was maximum in isolate VSAC-12 followed by VSAC-13, VSAC-22, VSAC-44 and VSAC-54. No activity was recorded in isolates, VSAC-6, VSAC-7, VSAC-8, VSAC-9, VSAC-16, VSAC-18, VSAC-19, VSAC-23, VSAC-28, VSAC-30, VSAC-35, VSAC-36, VSAC-43, VSAC-45, VSAC-48 and VSAC-55. The VSAC-12 actinomycetes for further study in our laboratory.




Fig 1. Keratinolytic actinomycete isolates from soils, composts pits of Gulbarga and Bellary District: (A) B-29; (B) C-11; (C) C-35.

## DISCUSSION

The biological activity of Indian subcontinent is one of the richest (second highest population in the world) owing to its vast geographical area, varied topology, climate, and the juxtaposition of several bio geographical regions. India is recognized as one of the twelve mega diversity regions in the world because of its richness in overall species diversity. Thus, India's contribution to the global diversity is around 8% (Manish kapoor and Rakesh kumar, 2004). It is important to study microbial diversity not only for basic scientific research, but also to understand the link between diversity and community structure and function. Human influences such as, pollution, agriculture and chemical application could adversely effect microbial diversity and perhaps also above and below ground ecosystem functioning (Jennifer *et al.*, 2004). Actinomycetes, at one time considered being microbiological curiosities of intensive search for the sources of new, biologically active compounds. The demonstrated ability of these organisms to produce useful antibiotics and to carry out other transformations of commercial interest has focused attention on factors bearing on their isolation from their natural habitat, the soil (Hesseltine *et al.*, 1954).

In the present study, different, compost pit areas, hair and feather dumping places and municipal wastes of Northern part of Karnataka were studied for the diversity of actinomycetes. The distribution of actinomycetes was observed to be predominant ( $2.4-7.80 \times 10^5$  cfu/g) in black soil when compared to red soil ( $1.90 - 3.55 \times 10^3$  cfu/g). The soils obtained from the alkaline habitat showed considerably very high predominance of actinomycetes than forest and salty soils. Abussaud and Saadoun (1991) were also reported  $10^5-10^7$  cfu/g of actinomycetes in dry soils of Jordan. A  $10^3-10^6$  cfu/g of actinomycetes were reported in acidic soils by Khan and Williams, (1975). Balagurunathan *et al.*, (1996) while studying diversity of soil actinomycetes from south India and south China recorded progressive increase in the number of actinomycetes from forest to vegetable farm lands, composts, municipal wastes which is in confirmation with observations made in this study. Soil is the most common habitat of actinomycetes, these bacteria can be found

in all kinds of soils (Kutzner, 1986 and Williams *et al.*, 1983). *Streptomyces* were found (Lacey and Crook, 1988), in composts and fooder, especially in self-heated hay or grain. During the early stages of composting or self-heating, mesophilic species were present, but these were replaced by thermotolerant species like, *Streptomyces albus* or *Streptomyces griseus*, and with increasing temperature, the real thermophilic species take their place (Goodfellow and Simpson, 1987). It is well known that bacteria, fungi and actinomycetes play an important role in decomposition of complex organic substrate found in nature. Bacterial and fungal growth rate is faster than actinomycetes, so they bring about faster degradation of these compounds. The complex organic compounds such as, chitin and keratin are not easily degradable by bacteria and fungi. Although slow growing actinomycetes decompose recalcitrant organic compounds. It is also seen that the percentage of biochemically active actinomycetes possessing amylase, protease, chitinase, cellulase and keratinase. A number of workers have been reported the presence of actinomycete isolates from soil with different enzymatic activity. Amylase activity has been reported by (Abraham and Hen, (1964): Chaphalkar and Dey (1993): Nawami, *et al.*, (2003) and Ammar *et al.*, (2002). Desai and Dhala, 1969: Ellaiah and Srinivasulu, *et al.*, 1996: Kulkarni and Deshmukh, 1999) have detected protease activity, and Shejul and Kapadnis, 1998: Nawami, *et al.*, 2003: Hiroshi Tsujibo, *et al.*, (2003) have detected chitinase activity. Cellulase activity reported by (Stutzenberger, 1972: Ali and Roymon, 1984: Shejul and Kapadnis, 1998: George *et al.*, (2001). Keratinase activity has been reported by Nickerson and Durand, 1963: Nadanishi and Yamamoto, 1974: Bockle, *et al.*, 1995: Muhin and Hadi, 2002: Suneetha and Lakshmi, (2003).

## REFERENCES

- [1] Abraham, T.A. and Hen, L.J. 1964. Activity of actinomycetes from rhizosphere and non-rhizosphere soils of corn and soybean in four physiological tests. *Can. J. Microbiol*, 10: 280-289.
- [2] Abussaud, M and Saadoun, I. 1991. *Streptomyces* flora of some Jordan Valley soils, characteristics and seasonal distribution. *Dirasat*, 18B: 66-75.
- [3] Abussaud, M. and Saadoun, I. 1991. *Streptomyces* flora of some Jordan Valley soils, characteristics and seasonal distribution. *Dirasat*. 18, 66-75.
- [4] Ali, S.S. and Royman, M.C. 1984. Antimiosis and biodegradability of some *Streptomyces* species from Raichur. *Ind. Microbiol*, 24: 124-126.
- [5] Ali, S.S. and Royman, M.C. 1984. Antimiosis and biodegrade ability of some *Streptomyces* species from Raichur. *Ind. Microbiol*, 24: 124-126.
- [6] Ammar, Y.B., Matsubara, T., Ito, K., Iizuka, M., Limpaseni, T., Pongsawasdi, P. and Minamiura, N. 2002. New action pattern of a maltose-forming alpha-amylase from *Streptomyces* sp. and its possible application in bakery. *J. Biochem. Mol. Biol*, 35(6): 568-75.
- [7] Balagurunathan, R., Xu, L. and Jiang, C. 1996. Diversity of soil actinomycetes from South India and South China. *Actinomycetes*, 4(3): 89-94.
- [8] Berdy, J. 1995. Are actinomycetes exhausted as a source of secondary metabolites. *Biotechnologia*, 8: 13-34.
- [9] Bockle, B., Galunsky, B. and Muller, R. 1995. Characterization of a keratinolytic serine proteinase from *Streptomyces pactum* DSM 40530. *Appl. Environ. Microbiol*, 61: 3705-3710.
- [10] Chaphalkar, S.R. and Dey, S. 1993. Studies of industrially important *Streptomyces* with special reference to extracellular protease. Ph.D. Thesis. University of Poona, Poona.
- [11] Desai, A.J. and Dhala, S.A. 1969. Purification and properties of proteolytic enzymes from thermophilic actinomycetes. *Journal of Bacteriology*, 100: 149-155.
- [12] Dhevendaran, K. and Annie, K. 1999. Antibiotic and L-asparaginase activity of streptomycetes isolated from fish, shellfish and sediment of Veli estuarine lake along kerala coast. *Indian. J. Mar. Sci*, 28: 335-337.
- [13] Ellaiah, P. and Srinivasulu, B. 1996. Production of extracellular protease by *Streptomyces fradiae*. *Hindustan Antibiot*, 38(4): 41-7.
- [14] Environment. Proceeding of international conference of SAARC Countries. Microbiologists Society.
- [15] George, S.P., Ahmad, A. and Rao, M.B. 2001. Studies on carboxymethyl cellulose produced by an alkalothermophilic actinomycete. *Bioresour. Technol*, 77(2): 25-29.
- [16] Gerber, N.N., 1983. Volatile substances from actinomycetes; their role in the odour pollution of water. *Water Sci. Technol*, 15(6-7): 115-125.
- [17] Goodfellow, M. 1983. Ecology of Actinomycetes. *Annu. Rev. Microbiol*, 37: 189-216.
- [18] Goodfellow, M. 1989. Super genic classification of actinomycetes. In *Bergey's Manual of Systematic Bacteriology* ed. Williams, S.T., Sharpe, M.E., Holt, J.G, 4: 2333-2339.
- [19] Goodfellow, M. and O'Donnell, A.G. 1994. Chemical Methods in Prokaryotic Systematics. John Wiley and Sons, New York.
- [20] Goodfellow, M. and Simpson, K.E. 1987. Ecology of *Streptomyces*. *Frontiers in Appl. Microbiol*, 2: 97-125.
- [21] Gray, T.R.G. and Williams, S.T. 1975. A soil microbes, (Eds). Edinburgh, Oliver and Boyd. Academic Press. London.
- [22] Hesseltine, C.W., Benedict, R.G. and Pridham, T.G. 1954. Useful criteria for species differentiation in the genus *Streptomyces*. *Ann New York Acad. Sci*, 60: 136-151.
- [23] Hiroshi Tsujibo, Takahiro Kubota, Mitsugu Yamamoto, Katsushiro Miyamoto, and Yoshihiko Inamori. 2003. Characterization of chitinase genes from an alkalophilic actinomycete, *Nocardiopsis prasina* OPC-131. *Applied and Environmental Microbiology*, 69(2): 894-900.
- [24] Jachymova, J., Vatruba, J., Viden. And Renzanka, T. 2002. Identification of *Strptomycetes* oder spectrum. *Folia. Microbial*, 47: 37-41.
- [25] Jennifer, L.K., Lee, A.B., Miranda, H., Peter, M., John, N.K., Hung Lee. and Jack, T.T. 2004. Methods of studying soil microbial diversity. *J. Microbiol. Meth*, 58: 169-188.
- [26] Jensen, D., 1995. Animal Welfare Information Center, USDA,

ARS Beltsville, MD. Personal communication.

- [27] Kawato, M. and Shinobu, R. 1959. On *streptomyces herbaricolor* nov. sp. supplement: A simple technique for the microscopical observation. Memoirs of the Osaka University Liberal Arts of Education, 8: 114.
- [28] Khan, M.R. and Williams, S.T. 1975. Studies on the ecology of actinomycetes in soil-VIII. Distribution and characteristics of acidophilic actinomycetes. *Soil Biol. Biochem*, 7: 345-348.
- [29] Kulkarni, S.W. and Deshmukh, A.M. 1999. Studies on soil actinomycetes of Solapur district. Ph.D. Thesis. Shivaji University, Kolhapur.
- [30] Kutzner, K.J. 1986. The family *Streptomycetaceae*. In: Starr, M.P., Stolp, H. Trper, H.G., Balows, A. and Schlegel, H.G. (Eds.). The prokaryotes, A Handbook on Habitats, Isolation, and Identification of Bacteria. Springer, 2: 2028-2090.
- [31] Lacey, J. 1973. Actinomycetes in soils, composts and fodders. In, Actinomycetes, Characteristics and practical importance (Skyes, G. and Skinner, F.A. Eds.) Academic Press, London. p.231-251.
- [32] Lacey, J. and Crook, B. 1988. Fungal and actinomycete spores as pollutants of the workplace and occupational allergens. *Ann. Occu. Hyg.* 32, 515-533.
- [33] Lechevalier, H.A. and Lechevalier, M.P. 1967. Biology of actinomycetes. *Ann. Rev. Microbiol*, 21: 71-100.
- [34] Manish Kapoor and Rakesh Kumar Jain. 2004. Microbial diversity-Exploring theunexplored. IMTECH - communication. 08/2004.
- [35] Muhin T.M. and Hadi, R.B. 2002. Degradation of keratin substrates by fungi isolated from sewage sludge. *J. Mycopathologia*, 4: 185-9.
- [36] Nadanishi, T. and Yamamoto, T. 1974. *Agricultural and Biological chemistry*, 38: 2391.
- [37] Nawami, N.N., Kapadnis, B.P., Das, A.P., Rao, A.S. and Mahajan, S.K. 2003. Purification and characterization of thermophilic and acidophilic chitinase from *Microbiospora* sp. V2. *J. Appl. Microbiol*, 93: 965-969.
- [38] Nawami, N.N., Kapadnis, B.P., Das, A.P., Rao, A.S. and Mahajan, S.K. 2003. Purification and characterization of thermophilic and acidophilic chitinase from *Microbiospora* sp. V2. *J. Appl. Microbiol*, 93: 965-969.
- [39] Nickerson, W.J. and Durand, S.C. 1963. Keratinase li. Properties of the crystalline enzyme. *Biochemica and Biophysica Acta*, 77:87-99.
- [40] Park, S.M., Yang, H.Y., Ching, K.C., Rhim, H., Park, J.H. and Kim. 2002. Stress- induced aggregation profiles of GST- $\alpha$ -Synuclein fusion proteins: Role of the C-terminal acidic tail of  $\alpha$ -synuclein in protein thermosolubility and stability. *Biochemistry*, 41: 4137-4146.
- [41] Seng.C.N., Kim, Y.S., Baik, K.S., Lee, S.D., Hah, Y.C., Kim, S.B and Goodfellow, M. 1999. Mycolic acid- containing actinomycetes associated with activated sludge foam. *J.Microbiology*, 37: 66-72.
- [42] Shejul, M.S. and Kapadnis, B.P. 1998. Studies on Heterotrophic Filamentous prokaryotes from aquatic habitat. Ph.D Thesis. University of Poona, Poona.
- [43] Shejul, M.S. and Kapadnis, B.P. 1998. Studies on Heterotrophic Filamentous prokaryotes from aquatic habitat. Ph.D Thesis. University of Poona, Poona.
- [44] Shinobu, R. 1958. Physiological and cultural study for the identification of soil Actinomycetes species. Memoirs of the Osaka University of the Liberal arts and selection. *Nat. Sci*, 7: 1-6.
- [45] Stutzenberge, F.J. 1972. Cellulolytic activity of *Thermomonospora curvate*: Optimal conditions, partial purification and product of cellulase. *Appi Microbioi*, 24: 83-90.
- [46] Suneetha, V. and Lakshmi, V.V. 2003. Fermentative production of kerazyme with isolates from Tirumala Hills. In: Biotechnology in Agriculture, Industry
- [47] Williams, S.T. and Wellington, E.M.H. 1980. Micromorphology and fine structure of actinomycetes. In: Goodfellow, M., Board, R.G. (Eds.), Microbial classification and identification. Academic Press, New York, London. p.139-165.
- [48] Williams, S.T., Goodfellow, M., Wellington, E.M.H., Vickers, J.C., Alderson, G. and Sneath, P.H.A. 1983. A probability matrix for identification of some *Streptomycetes*. *J. Gen. Microbiol*, 129: 1815-1830.