

Research Article

Antioxidant enzymatic effects of diethylene glycol dibenzoate on zebrafish (*Danio rerio*)

K. Sasikumar*, O. S. Sethu Raman, D. Senthil Kumar

Department of Zoology, Kandaswami Kandari's College, Velur, Namakkal-638182, Tamil Nadu, India

(Received: August 13, 2025; Revised: January 06, 2026; Accepted: January 08, 2026; Published: January 20, 2026)

*Corresponding author: K. Sasikumar, E-mail: drsasi03@gmail.com

Abstract

Endocrine disrupting chemicals (EDCs) are known to disrupt normal metabolism and can influence the incidence of obesity in animals and humans. EDCs can exert adverse effects at low concentrations, often in a non-monotonic dose-related fashion. Among EDCs, diethylene glycol dibenzoate (DGB), an approved alternative to phthalates in the production of plastic and latex products, is less abundant and its effects are almost completely unknown. Potential EDCs can disrupt the normal hormonal level by inhibiting or stimulating the production of hormones or changing the way in which the hormones are transported to target tissues. In specific, the effect of DGB has expanded to include the deregulation of the antioxidant defense (AD) system of organisms dealing with oxidative stress and several enzymes metabolism. The present study focused on the changes elicited by DGB is on the enzyme activity catalase and superoxide dismutase of the various organs (muscle, gill, and liver) of zebrafish. The observation registered in this study reflects that antioxidant enzyme activities were significantly enhanced in all the tissues (muscle, gill, and liver) when compared to control (untreated DGB). This in turn induces some enzymatic variations that lead to the changes in hormonal secretion which is responsible for functioning of targeted organs and its generative activities. This could be due to the detoxification mechanism exhibited by the zebrafish on exposure of DGB. This finding provides a support for the hypothesis that DGB may be the environmental contaminant with stress property and in need to control of its exposure immediately and completely. Also this finding may help to initiate and follow the innovative methods, to control the use of DGB in our day today life.

Keywords: Diethylene glycol dibenzoate, Zebrafish, Catalase, Superoxide dismutase, Antioxidant defense

Introduction

Today a variety of endocrine disrupting chemicals (EDCs) are recognized in the group of metabolic disruptors that includes obesogens, a wide range of environmental contaminants capable of altering energy balance regulation leading to obesity (Grün & Blumberg, 2006). Potential EDCs can disrupt the normal hormonal level by inhibiting or stimulating the production of hormones or changing the way in which the hormones are transported to target tissues (Heindel *et al.*, 2017). Recently, obesogen list has expanded to include contaminants that deregulate lipid metabolism (Chamorro-García & Blumberg, 2014). Information on harmful effect of DGB which is an approved alternative to phthalates in the processing of plastic and latex (Kermanshahi pour *et al.*, 2009) is very limited. However, a recent study demonstrated that exposure to DGB leads to the stimulation of crucial lipolytic genes via PPAR α mediated pathway. These warrant further research and increase the interest toward the identification of new mechanisms for EDC metabolic alterations and cause "transgenerational effects" by a single prenatal or early life exposure (Maradonna & Carnevali, 2018).

The antioxidant defense (AD) system of organisms provides a means of dealing with oxidative stress and includes several enzymes and vitamins (Filho, 1996; Rudneva, 1997; Kelly *et al.*, 1998; Marcon & Filho, 1999). A primary role of the AD system is protecting cellular components from ROS damage (Kelly *et al.*, 1998). The antioxidant enzymes, superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase, act to remove oxygen

radicals produced within the cell (Filho *et al.*, 1993; Michiels *et al.*, 1994; Filho, 1996; Kelly *et al.*, 1998). Superoxide dismutase occurs in two forms: 1) a cytosolic form that has copper and zinc in its active site (CuZnSOD) and 2) a mitochondrial form that has manganese in its active site (MnSOD) (McCord & Fridovich, 1969). Both forms of SOD protect the cell from potential ROS damage by converting superoxide anions to H₂O₂ and H₂O (McCord & Fridovich, 1969).

Antioxidant system is located within different cellular compartments. These enzymes are found virtually in all tissue of vertebrates, but show in general high activity in the liver. A major organ of xenobiotics uptake and enzymatic transformation of ROS (Lemaire *et al.*, 1992) eventually leak into blood. Some of these enzymes, like aminotransferase and phosphatase group, can constitute good molecular bioindicators for oxidative stress and can also indicate the magnitude of response in populations chronically exposed to contaminants such as metals and other xenobiotics (Livingstone *et al.*, 1994). Keeping this in view, the aim of this study was to assay the enzyme activity of CAT, SOD of the muscle, gill and liver of zebrafish due to exposure of DGB.

Materials and methods

Adult female zebrafish (*Danio rerio*, wild-type strain) were maintained in 100 L aquaria with oxygenated water under controlled conditions (28.0 \pm 0.5 °C under a 14/10 h of light/dark period) in accordance with protocols and procedures approved by the University of Calgary Animal

Care Committee. Fish were fed four times per day, twice with commercial adult zebrafish complete diet (Zeigler Bros., Inc.) and twice with *Artemia salina*. For the DGB trial, a total of fishes were equally distributed into three aquaria (one control and two DGB experimental groups) with a total of 10 fish in each one and treated groups. At the end of the 30th day, ten zebrafish from each of the treatments and control were collected. The Gill, liver and muscle of zebrafish were dissected and subjected to enzyme assay, SOD and CAT. About 0.4 mg of tissue sample was homogenized in 800 µL Tris-Hcl buffer (100 mM, pH-7.4) using a glass homogenizer. Then the homogenate was centrifuged at 10000 rpm for 15 min at 4 °C, and the supernatant was collected for antioxidant enzymes. Catalase (EC 1.11.1.6) activity was assessed following the Aebi method (Aebi, 1984) and SOD activity was assayed based on a modified method of Marklund and Marklund (1974).

Statistical analysis

Results of the experiment were expressed as mean and standard error of mean of different groups. The differences between the mean values were evaluated by ANOVA (16.0). The values for P<0.001 were considered significant. Accordingly, a statistical software package (SPSS) was used.

Results and discussion

Effect of DGB exposure to zebrafish for CAT assay

The enzyme activities of various tissues were assayed in zebrafish exposed to DGP. The data displayed in the CAT activities were significantly decreased (p<0.05) in DGB treated in vivo tissues compared with control (Table 1).

Effect of DGB exposure to zebrafish for SOD assay

The enzyme activity of various tissues was assayed in zebrafish exposed to DGP. The data displayed in the SOD activities were significantly increased (p<0.05) in DGB treated in vivo tissues compared with control (Table 2). SOD activity was elevated in brains of carp and the neotropical fish *Hoplias malabaricus* induced by heavy metals, synthetic organic pollutants, and biotoxins (Da Silva *et al.*, 2011; Xing *et al.*, 2012). Therefore, increased SOD activity following DBP exposure, may be attributed to the cells adaptive defense mechanism to eliminate surplus O[•]. CAT is a key enzymatic antioxidant system, which can eliminate

H₂O₂ produced from ROS, catalyzed by SOD, and thus alleviate cell damage (Zhao *et al.*, 2014).

SOD and CAT play important roles in protecting the cell against the potentially toxic effects of experimental pollutants (Kuthan *et al.*, 1986). SOD, catalyzes the dismutation of the superoxide ion (O²⁻) to hydrogen peroxide and oxygen molecule during energy processes. The reaction diminishes the caustic oxidative processes in cells. The level of antioxidant enzymes have been extensively used as an early warning indicator of lake pollution (Lin & Hwang, 1998). The present result is in good accord with the findings of Neeraj Kumar *et al.* (2011) who have confirmed that activities of anti-oxidative enzymes was significantly (P<0.01) influenced by endosulfan in a dose dependent manner in *Tilapia Oreochromis mossambicus*. They have noticed significant (P<0.01) increase in the activity of CAT, SOD and GST in gill and liver of *Tilapia*. Superoxide dismutase is an antioxidant enzyme involved in the elimination of ROS (reactive oxygen species). CAT is one among the key antioxidant enzymes that catalyses the removal of H₂O₂ formed during the reaction catalyzed by SOD (Arun & Subramanian, 2002). Biomarkers such as protein level, enzyme activity and DNA can be used to measure the interaction between biological systems and chemical, physical or biological environmental agents (Hernández-Moreno *et al.*, 2010). The liver is the primary organ for detoxification of xenobiotics and excretion of toxic substances in fish (El-Naggar *et al.*, 2009). The important function of the liver is to clean of any pollutant from the blood, but these pollutants could subsequently lead to structural damage in liver (Pathan *et al.*, 2010) as seen in the present study. Liver antioxidant enzymes are used to determine if the liver is functioning normally or if it has an injury or disease. Studies showed that enzymatic techniques are inexpensive and reliable to determine the toxicities of pollutants on marine animals in the living environment (Hegazi *et al.*, 2010). Superoxide dismutase is an antioxidant enzyme involved in the elimination of ROS (reactive oxygen species). The antioxidant enzymes that make up the antioxidant defense system are expected to be intrinsically linked and dependent upon the activity of one another. Therefore, one could expect to see correlative changes in the activity of SOD and CAT (Filho *et al.*, 1993).

Oxidative stress occurs if the activity of the antioxidant defense systems such as SOD, CAT and GPx (glutathione peroxidase) enzymes change by environmental pollution induces the production of reactive oxygen species (Li *et al.*, 2011). SOD is a vital antioxidant defense enzyme in nearly

Table 1: Changes in the CAT of the various tissues of zebrafish exposed to diethylene glycol dibenzoate

DGB treatment	Muscle protein (U/mg)	Gill protein (U/mg)	Liver protein (U/mg)
Control	4.240±0.065 ^c	7.123±0.017 ^c	9.160±0.028 ^c
1 ppm	3.160±0.030 ^b	6.183±0.014 ^b	4.086±0.014 ^b
10 ppm	2.250±0.016 ^a	3.213±0.010 ^a	2.195±0.010 ^a

***Significant at P<0.001. In a column, figures having dissimilar letters differ significantly according to Duncan New Multiple Range Test (DMRT). Values are expressed as mean±standard error

Table 2: Changes in the SOD of the various tissues of exposed to zebrafish exposed to diethylene glycol dibenzoate

DGP Treatment	Muscle protein (U/mg)	Gill protein (U/mg)	Liver protein (U/mg)
Control	10.183±0.045 ^c	8.233±0.056 ^c	4.170±0.026 ^c
1 ppm	15.213±0.020 ^b	20.213±0.037 ^b	6.736±0.020 ^b
10 ppm	19.356±0.024 ^a	23.913±0.361 ^a	7.193±0.038 ^a

***Significant at P<0.001. In a column, figures having dissimilar letters differ significantly according to Duncan New Multiple Range Test (DMRT). Values are expressed as mean±standard error

all cells to catalyze the dismutation of superoxide into oxygen thus protecting the cell from superoxide toxicity. SOD enzyme is more sensitive to the lethal effects of superoxide generating chemicals (Gardner *et al.*, 1995). CAT is found in highest levels in the liver as a result of breaking down toxins present in the blood and processing metabolic products for degradation (Aebi, 1984; Chelikani *et al.*, 2004). The other study indicated that CAT and SOD activities were significantly higher in liver than in brain and gill of fish exposed to pesticide (Li *et al.*, 2011). The present results agree with earlier reports of increased antioxidant enzymes in fish exposed to environmental pollutants.

Conclusion

The diethylene glycol dibenzoate (DGB) was approved by European Chemical Agency as an alternative to phthalates in the processing of plastic. The alteration in the enzyme activity observed in the present study could be due to the formation of reactive oxygen species such as hydrogen peroxide, superoxide anion (O_2^-), hydroxyl radicals (OH \cdot), which has to be neutralized by antioxidant enzymes. We can conclude that DGP may not provide a safe substitute and more studies on DGP safety will be needed on safety of DGP in fish and other organisms. Our study provides a framework for better understanding of the mechanisms underlying adverse health impact of stress EDCs. DGB is characterized as a high solvating plasticizer intended for the use in the manufacturing of PVC, vinyl flooring, adhesives, latex caulks, sealants and elastomers. These publications suggest that while DGB does not last long in the organism, the intermediates of its detoxification do, and the results do not support the use of DGB as an environmentally safer alternative to phthalates.

References

Aebi, H. E. (1984). Catalase. In H. U. Bergmeyer (Ed.), *Methods of Enzymatic Analysis* (Vol. 3, pp. 273-286). John Wiley & Sons, Inc.

Arun, S., & Subramanian, P. (2002). Antioxidant enzymes in freshwater prawn *Macrobrachium malcolmsonii* during embryonic and larval development. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 121(3), 273-277. [https://doi.org/10.1016/S0305-0491\(98\)10100-1](https://doi.org/10.1016/S0305-0491(98)10100-1)

Chamorro-García, R., & Blumberg, B. (2014). Transgenerational effects of obesogens and the obesity epidemic. *Current Opinion in Pharmacology*, 19, 153-158. <https://doi.org/10.1016/j.coph.2014.10.010>

Chelikani, P., Fita, I., & Loewen, P. C. (2004). Diversity of structures and properties among catalases. *Cellular and Molecular Life Sciences*, 61, 192-208. <https://doi.org/10.1007/s00018-003-3206-5>

da Silva, C. A., Oba, E. T., Ramsdorf, W. A., Magalhães, V. F., Cestari, M. M., Ribeiro, C. A. O., de Assis, H. C. S. (2011). First report about saxitoxins in freshwater fish *Hoplias malabaricus* through trophic exposure. *Toxicon*, 57(1), 141-147. <https://doi.org/10.1016/j.toxicon.2010.10.015>

El-Naggar, A. M., Mahmoud, S. A., & Tayel, S. I. (2009). Bioaccumulation of some heavy metals and histopathological alterations in liver of *Oreochromis niloticus* in relation to water quality at different localities along the River

Nile, Egypt. *World Journal of Fish and Marine Sciences*, 1(2), 105-114.

Filho, D. W. (1996). Fish antioxidant defenses - a comparative approach. *Brazilian Journal of Medical and Biological Research*, 29(12), 1735-1742.

Filho, D. W., Giulivi, C., & Boveris, A. (1993). Antioxidant defenses in marine fish: I. Teleosts. *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology*, 106(2), 409-413. [https://doi.org/10.1016/0742-8413\(93\)90154-D](https://doi.org/10.1016/0742-8413(93)90154-D)

Gardner, P. R., Raineri, I., Epstein, L. B., & White, C. W. (1995). Superoxide radical and iron modulate aconitase activity in mammalian cells. *Journal of Biological Chemistry*, 270(22), 13399-13405. <https://doi.org/10.1074/jbc.270.22.13399>

Grün, F., & Blumberg B. (2006). Environmental obesogens: Organotins and endocrine disruption via nuclear receptor signaling. *Endocrinology*, 147(6), 50-55. <https://doi.org/10.1210/en.2005-1129>

Hegazi, M. M., Attia, Z. I., & Ashour, O. A. (2010). Oxidative stress and antioxidant enzymes in liver and white muscle of Nile tilapia juveniles in chronic ammonia exposure. *Aquatic Toxicology*, 99(2), 118-125. <https://doi.org/10.1016/j.aquatox.2010.04.007>

Heindel, J. J., Blumberg, B., Cave, M., Machtinger, R., Mantovani, A., Mendez, M. A., Nadal, A., Palanza, P., Panzica, G., Sargis, R., Vandenberg, L. N., & vom Saal, F. (2017). Metabolism disrupting chemicals and metabolic disorders. *Reproductive Toxicology*, 68, 3-33. <https://doi.org/10.1016/j.reprotox.2016.10.001>

Hernández-Moreno, D., Soler, F., Míguez, M. P., & Pérez-López, M. (2010). Brain acetyl cholinesterase, malondialdehyde and reduced glutathione as biomarkers of continuous exposure of tench, *Tinca tinca*, to carbofuran or deltamethrin. *Science of The Total Environment*, 408(21), 4976-4983. <https://doi.org/10.1016/j.scitotenv.2010.07.044>

Kelly, K. A., Havrilla, C. M., Brady, T. C., Abramo, K. H., & Levin, E. D. (1998). Oxidative stress in toxicology: established mammalian and emerging piscine model systems. *Environmental Health Perspectives*, 106(7), 375-384. <https://doi.org/10.1289/ehp.98106375>

Kermanshahi pour, A., Cooper, D. G., Mamer, O. A., Maric, M., Nicell, J. A. (2009). Mechanisms of biodegradation of dibenzoate plasticizers. *Chemosphere*, 77(2), 258-263. <https://doi.org/10.1016/j.chemosphere.2009.06.048>

Kuthan, H., Haussmann, H. J., & Werringloer, J. (1986). A spectrometric assay for superoxide dismutase activities in crude tissue fractions. *Biochemical Journal*, 237(1), 175-180. <https://doi.org/10.1042/bj2370175>

Lemaire, P., Berhaut, S., Lemaire-Gony, S., & Lafaurie, M. (1992). Ultrastructural changes induced by benzo[a]pyrene in sea bass (*Dicentrarchus labrax*) liver and intestine: Important of intoxication route. *Environmental Research*, 57(1), 59-72. [https://doi.org/10.1016/S0013-9351\(05\)80019-2](https://doi.org/10.1016/S0013-9351(05)80019-2)

Li, Z.-H., Zlabek, V., Velisek, J., Grabic, R., Machova, J., Kolarova, J., Li, P., & Randak, T. (2011). Acute toxicity of carbamazepine to juvenile rainbow trout (*Oncorhynchus mykiss*): Effects on antioxidant responses, hematological parameters and hepatic EROD. *Ecotoxicology and Environmental Safety*, 74(3), 319-327. <https://doi.org/10.1016/j.ecoenv.2010.09.008>

Lin, H. C., & Hwang, P. P. (1998). Acute and chronic effects of gallium chloride (GaCl $_3$) on tilapia (*Oreochromis mossambicus*) larvae. *Bulletin of Environmental Contamination and Toxicology*, 60, 931-935. <https://doi.org/10.1007/s001289900717>

Livingstone, D. R., Forlin, L., & George, S. G. (1994). *Molecular biomarkers and toxic consequences of impact by organic*

- pollution in aquatic organism. D. Wsutcliffe, editor water quality and stress indicators in marine and fresh water ecosystem UK* (pp. 154-171). Freshwater Biological Association.
- Maradonna, F., & Carnevali, L. (2018). Lipid metabolism alteration by endocrine disruptors in animal models: An overview. *Frontiers in Endocrinology*, 9, 654. <https://doi.org/10.3389/fendo.2018.00654>
- Marcon, J. M., & Filho, D. W. (1999). Antioxidant processes of the wild tambaqui, *Colossoma macropomum* (Osteichthyes, Serrasalminidae) from the Amazon. *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology*, 123(3), 257-263. [https://doi.org/10.1016/S0742-8413\(99\)00030-4](https://doi.org/10.1016/S0742-8413(99)00030-4)
- Marklund, S., & Marklund, G. (1974). Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *European Journal of Biochemistry*, 47(3), 469-474. <https://doi.org/10.1111/j.1432-1033.1974.tb03714.x>
- McCord, J. M., & Fridovich, I. (1969). Superoxide Dismutase: An enzymatic function for erythrocyte hemoglobin. *Journal of Biological Chemistry*, 244(22), 6049-6055. [https://doi.org/10.1016/S0021-9258\(18\)63504-5](https://doi.org/10.1016/S0021-9258(18)63504-5)
- Michiels, C., Raes, M., Toussaint, O., & Remacle, J. (1994). Importance of SE-glutathione peroxidase, catalase, and Cu/Zn-SOD for cell survival against oxidative stress. *Free Radical Biology and Medicine*, 17(3), 235-248. [https://doi.org/10.1016/0891-5849\(94\)90079-5](https://doi.org/10.1016/0891-5849(94)90079-5)
- Neeraj Kumar, Antony Jesu Prabhu, P., Pal, A. K., Remya, S., Md. Aklakur, Rana, R. S., Subodh Gupta, Raman, R. P., & Jadhao, S. B. (2011). Anti-oxidative and immuno-hematological status of Tilapia (*Oreochromis mossambicus*) during acute toxicity test of endosulfan. *Pesticide Biochemistry and Physiology*, 99(1), 45-52. <https://doi.org/10.1016/j.pestbp.2010.10.003>
- Pathan, T., Shinde, S. E., Thete, P., & Sonawane, L. D. (2010). Histopathology of liver and kidney of *Rasbora daniconius* exposed to study mill effluent. *Research Journal of Biological Sciences*, 5(5), 389-394.
- Rudneva, I. I. (1997). Blood antioxidant system of Black Sea elasmobranch and teleosts. *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology*, 118(2), 255-260. [https://doi.org/10.1016/S0742-8413\(97\)00111-4](https://doi.org/10.1016/S0742-8413(97)00111-4)
- Xing, H., Li, S., Wang, Z., Gao, X., Xu, S., & Wang, X. (2012). Histopathological changes and antioxidant response in brain and kidney of common carp exposed to atrazine and chlorpyrifos. *Chemosphere*, 88(4), 377-383. <https://doi.org/10.1016/j.chemosphere.2012.02.049>
- Zhao, X., Gao, Y., & Qi, M. (2014). Toxicity of phthalate esters exposure to carp (*Cyprinus carpio*) and antioxidant response by biomarker. *Ecotoxicology*, 23, 626-632. <https://doi.org/10.1007/s10646-014-1194-x>