

Diversity of Fresh Water Hyphomycetes from Buldhana District (M.S.), India

L. C. Nemade¹, V. R. Patil¹, M.S. Patil¹ and S.A. Chaudhar²

¹Sbri V. S. Naik Arts, Commerce & Science College, Raver District- Jalgaon. 425508 (M.S.), India

²D. A.G.D. Bendale Mahila Mahavidyalaya, Jalgaon (M.S.), India

*Corresponding author, Email:

Keywords

Hyphomycetes
Freshwater fungi
India

Abstract

The present study deals with five species of freshwater hyphomycetes collected in foam samples from Purna river of Bhuldhana District. All the five species are being reported for the first time from Bhuldhana District.

1. Introduction

The occurrence of water-borne fungi has been reported from various parts of Europe, Africa, America, Asia and Australia (Ingold, 1975; Subramanian, 1971). To India, the aquatic hyphomycetes were studied by Ingold and Webster (1973), Subramanian and Bhat (1981), Sridhar *et al.* (1992), Galiath and Manoharachary (1987), Agrawal *et al.* (1991), Sati and Tiwari (1997), Sati *et al.* (2002). In Maharashtra, these fungi were recorded by Thakur (1977), Patil and Kapadnis (1980), Patil (1998), Borse and Patil (2006), Borse and Patil (2007). The present study reports five species of freshwater hyphomycetes from Bhuldhana District over a period of one year (2008-2009).

2. Materials and Methods

Samples of foam were collected from the river Purna of Bhuldhana District.

Foam analysis

In general, the foam formed by the movement of the water against natural barriers like stones, twigs and logs, especially in lotic systems, constitutes a natural trap for the conidia of aquatic hyphomycetes. Foam samples were collected at morning and evening time. Samples were made with a ladle and placed in clean wide mouthed plastic bottles and kept for 24 hours to enable the foam to dissolve. It was preserved by adding

FAA. Then samples were brought to the laboratory and observed under research microscope for the presence of conidia of hyphomycetes.

The permanent slides were prepared as suggested by Kohlmeyer & Kohlmeyer (1996). Identification of the freshwater hyphomycetes were confirmed with the help of Nilsson (1964), Ingold (1975), Marvonova (1997), and Cai *et al.* (2003) and other relevant literature. Reports of fungi from India and Maharashtra were confirmed with the help of Kamat *et al.* (1971), Bhide *et al.* (1987), Mahabale (1987), Bilgrami *et al.* (1979, 1981, 1991), Sridhar *et al.* (1992), Sarbhoy *et al.* (1986, 1996), Jamaluddin *et al.* (2004) and other relevant literature.

Taxonomic account

Anguillospora crassa Ingold (Plate 1, Fig. 1)


Conidia: hyaline, S or L shaped, 120- 200 μ m long and 15- 20 μ m wide in the middle region, tapering to 8- 10 μ m; hyaline, branched, the main body globose or ovoid at the ends.

Habitat: Conidia in foam samples, Purna river.

Distribution in India: Maharashtra (Thakur, 1977), Karnataka (Sridhar and Kaveriappa, 1986, 1992), Kumaun Himalaya (Sati and Tiwari, 1990).

Remark: The measurements and descriptions of conidia are completely agree with that of *Anguillospora crassa*. Therefore, it is assigned to that species.

Plate 1. Fig. 1...*Anguillospora crassa* Ingold, Fig.2. *Anguillospora longissima* (Sacc. And Sydow) Ingold, Fig.3.*Campylospora chaetocladia* Ranzoni, Fig.4. *Alatospora acuminata* Ingold, Fig.5.*Isthmotricladia gombakiensis* Nawawi.



Anguillospora longissima (Sacc. And Sydow) Ingold (Plate 1, Fig. 2.)

Conidia: unbranched, elongated, 8- 12 septate, sigmoid with curvature in more than one plane, 200- 280 X 2.5- 3.5 μm .

Habitat: Conidia in foam samples, Purna river.

Distribution in India: Maharashtra (Thakur, 1977, Talde, 1983), Western Ghat (Subramanian and Bhat, 1981), Kerala (Sridhar and Kaveriappa, 1985), Karnataka (Sridhar and Kaveriappa, 1982, 1988; Ramesh and Vijaykumar 2000; Western Ghat, Rajashekhar and Kaveriappa, 2003).

Remark: The measurements and descriptions of conidia are completely agree with that of *Anguillospora longissima* (Sacc. And Sydow). Therefore, it is assigned to that species.

Campylospora chaetocladia Ranzoni (Plate 1, Fig.3)

Conidia: composed of two parts, proximal half triangular, 3-4 septate, 8- 12.5 μm high, 10- 12 μm wide at the base, distal half allantoids, 3- 4 celled, 9- 13 μm long, 3.5- 5 μm wide. Appendages arising from end cells, setae like, 30- 40 μm long.

Habitat: Conidia in foam samples, Purna river.

Distribution in India: Karnataka (Sridhar and Kaveriappa, 1982, 1986, 1988, 1989, 1992), Kerala (Sridhar and Kaveriappa, 1985), Andhra Pradesh (Gaillah and Manoharachary, 1987), Maharashtra (Borse and Patil, 2006).

Remark: It has been reported for the first time from Purna river.

Alatospora acuminata Ingold (Plate 1, Fig.4)

Conidia: typically tetradiate, axis long-fusoid, arcuate or bent at branch insertion or nearly straight, 30-70 X 1.0-2.5 μm . With up to six septa; branches 12-35 X 1.0-2.2 μm , base

decurent to narrow, inserted near the middle or in the lower half of the axis; with up to four septa.

Habitat: Conidia in foam sample, Purna river.

Distribution In India: Kerala (Sridhar and Kaveriappa, 1985); Karnataka (Sridhar and Kaveriappa, 1982, 1986, 1989); Kumaun Himalaya (Sati and Tiwari, 1990); Maharashtra (Patil, 1998).

Remark: It has being reported for the first time from Buldhana District.

Isthmotricladia gombakiensis Nawawi (Plate 1, Fig.5)

Conidia: hyaline, main axis 20- 27 X 2- 3 μm , 1-3 septate, mature conidia consisting of four to six arms, the arms are fusiform, 74- 102 μm long, 4- 5.5 μm wide, 9- 15 septate, tapering to 1- 1.5 μm at the apex by very narrow isthmus, 2- 4.5 μm long, 1- 1.5 μm wide.

Habitat: Conidia in foam samples, Purna river.

Distribution in India: Karnataka (Sridhar and Kaveriappa, 1984, 1982; Ramesh and Vijaykumar, 2000), Andhra Pradesh (Sarma and Manoharachary, 1989), Western Ghats, (Rajashekhar and Kaveriappa, 2003), Maharashtra (Borse and Patil, 2006).

Remark: The measurements and descriptions of conidia are completely agree with that of *Isthmotricladia gombakiensis* Nawawi (1975).

Acknowledgements

The authors are thankful to Dr. Suresh Pawar, Principal, Shri V. S. Naik College, Raver Dist. Jalgoan (M.S.) for laboratory and library facilities.

References

Agarwal, G. P., Hasija, S. K., Agarwal, P. and Pandey, A. K. (1991). Fungi associated with

- submerged decaying leaves and twigs from Jabalpur. *Proc. Nat. Acad. Sci. India*, **61**:121-125.
- Bhide, V.P., Pande, Alka., Sathe, A.V., Rao, V.G. and Patwardhan, P.G. (1987). *Fungi of Maharashtra (Sup-I)*, MACS Res. Institute Publication, Pune, Maharashtra, pp.1-146.
- Bilgrami, K. S., Jamaluddin, S. and Rizwi, M. A. (1979). *Fungi of India*. Part-I. Today and Tomorrow's Print. and Pub., New Delhi, pp.467.
- Bilgrami, K. S., Jamaluddin, S. and Rizwi, M. A. (1981). *Fungi of India*. Part-II. Today and Tomorrow's Print. and Pub., New Delhi, pp.268.
- Bilgrami, K. S., Jamaluddin, S. and Rizwi, M. A. (1991). *Fungi of India*. Today and Tomorrow's Print. and Pub., New Delhi, pp.798.
- Borse, B. D. and Patil, R. S., (2007). Aquatic fungi from North Maharashtra-I. *Bioinfolet*, **4**:101-104.
- Borse, B. D. and Patil, S. Y. (2006). Aquatic fungi from North Maharashtra-IV: *BRI's, J. Adv. Sci. and Tech.*, **9**:91-95.
- Cai, L., Zhang, K. Q., Mc Kenzie, E. H. C. and Hyde, K.D. (2003). New species of *Dictyosporium* and *Digitodesmium* from submerged wood in Yunnan, China. *Sydowia*, **55**:129-135.
- Galiah, K, and Manoharachary, C. (1987). Studies on conidial fungi of a stream from Andhra Pradesh. *Indian Phytopath.*, **40**:466-473.
- Ingold, C. T. (1975). "An illustrated guide to Aquatic and water-Borne Hyphomycetes." Freshwater Biological Associ. Sci. Publ. **30**:PP-96.
- Ingold, C. T. and Webster, J. (1973). Some aquatic hyphomycetes from India. *Kavaka*, **1**:5-9.
- Jamaluddin, S., Goswami, M.G. and Ojha, B. M. (2004). *Fungi of India (1989-2001)*, Scientific Publishers (India) Jodhpur, pp.308.
- Kohlmeyer, B. & Kohlmeyer, J. (1996). How to prepare truly permanent microscopic slides. *Mycologist*, **10**: 107-108.
- Kamat, M. N., Patwardhan, P. G., Rao, V. G., and Sathe, A. V. (1971). *Fungi of Maharashtra*. Bulletin No.-I, M.P. Agril. Uni. Pub., Rahuri (M.S.), pp.124.
- Mahabale, T.S. (1987). *Botany and Flora of Maharashtra*. Gazetteer of India, M.S. Gazetteers, Govt. of M.S., pp.169-222.
- Marvanova, L. (1997). Freshwater hyphomycetes : a survey with remarks on tropical taxa In: *Tropical Mycology* (eds. Janardhanan, Rajendran, Natrajan and Hawksworth), Oxford and IBH Publishing Co. Pvt. Ltd., Calcutta, pp. 169-226.
- Nawawi, A. (1975). *Triscelosporus acuminatus* sp. nov. *Trans. Br. Mycol. Soc.*, **64**:345-348.
- Nilsson, S. (1964). *Freshwater hyphomycetes: Taxonomy, Morphology and Ecology*. *Symb. Bot. Ups.*, **18**:1-130.
- Patil, N.N. (1998). Aquatic hyphomycetes of Mahabaleshwar. *Geobios New Reports*, **17**:90.
- Patil, S.D. and Kapadnis, B. P. (1980). Stream spora of Maharashtra. *MVMP*, **14**: 59-64.
- Rajashekhkar, M. and Kaveriappa, M. (2003) Diversity of aquatic hyphomycetes in the aquatic ecosystem of the Western Ghats of India. *Hydrobiologia*, **501**: 167-177.
- Ramesh. Ch. and Vijaykumar, S. (2000). Seasonal occurrence of water borne fungi in Panda stream, Uttara Kannada Region, Karnataka. In: *Ecology of fungi* (eds. Bhat and Raghukumar), Gao University Press, Gao, India, pp.21-27.
- Sarbhoj, A. K., Agarwal, D.K. and Varshney, J. L. (1986). *Fungi of India*. Association Publishing Company, New Delhi, pp.274.
- Sarbhoj, A. K., Agarwal, D.K. and Varshney, J. L. (1996). *Fungi of India (1982-1992)*, CBS Publishers and Distributors, New Delhi, pp.276.
- Sarma, T.K. and Manoharachary, C. (1989). Numerical studies on conidial fungi from a stream of Andhra Pradesh. *Indian Phytopath.*, **42**:596-598.
- Sati, S. C. and Tiwari, N. (1990). Freshwater hyphomycetes from Jagashwar stream, Kumaun Himalaya India. *Nat. Acad. Lett.*, **13**:7-9.
- Sati, S. C., Tiwari, N. and Belwal, M. (2002). Conidial aquatic fungi of Nanital, Kumaun Himalaya, India. *Mycotaxon*, **81**:445-455.
- Sridhar, K. R. and Kaveriappa, K. M. (1982). Aquatic fungi on the Western Ghats forest in Karnataka. *Indian Phytopath.*, **35**:293-296.
- Sridhar, K. R. and Kaveriappa, K. M. (1984). Aquatic hyphomycetes of the Western Ghat forests in Karnataka. *Indian Phytopath.*, **37**: 546-548.
- Sridhar, K. R. and Kaveriappa, K. M. (1985). Water-Borne fungi of Kunthi River in Silent Valley-Kerala. *Indian Phytopath.*, **38**:371-372.
- Sridhar, K. R. and Kaveriappa, K. M. (1986). New records of aquatic hyphomycetes. *Indian Phytopath.*, **39**:131-132.
- Sridhar, K. R. and Kaveriappa, K. M. (1988). Survival of water-Borne fungi-imperfecti under non-aquatic conditions. *Proc. Ind. Nat. Sci. Acad.*, **54**:295-297.
- Sridhar, K. R. and Kaveriappa, K. M. (1989). Notes on aquatic hyphomycetes of mountain streams in western ghats region India. *Feddees Reportorium*, **100**:187-189.
- Sridhar, K. R. and Kaveriappa, K. M. (1992). Aquatic hyphomycetes of Western ghats streams, India. *Sydowia*: **44**
- Sridhar, K. R., Chandrashekar, K. R. and Kaveriappa, K. M. (1992). Research on the Indian subcontinents. In: *The Ecology of aquatic*

- hyphomycetes* (Eds. Barlocher), Spinger-Varlag., Heidelberg Press, New York, pp.182-211.
- Subramanian, C. V. (1971). *Hyphomycetes: An account of Indian species, except Cercosporae*. I. C.A.R. Publ., pp.930.
- Subramanian, C. V. and Bhat, D. J. (1981). Conidia from freshwater foam samples from the Western Ghats, South India. *Kavaka*, **9**: 45-62.
- Thakur, S. B. (1977). Survival of some aquatic Hyphomycetes under dry condition. *Mycologia*, **69**:843-845.

Please Cite This Article As:

L. C. Nemade, V. R. Patil, M.S. Patil and S.A. Chaudhari. 2010. Diversity of Fresh Water Hyphomycetes from Buldhana District (M.S.), India. *J. Ecobiotechnol.* 2(6):17-20.