



REGULAR ARTICLE

A SIMPLE AND RAPID METHOD FOR DNA EXTRACTION FROM LEAVES OF TOMATO, TOBACCO AND RAPE SEED

Praveen Mamidala^{1,2#}, Rama Swamy Nanna²

¹Department of Biotechnology, Vaagdevi College of Engineering, Warangal – 506005, India

[#]Department of Plant Sciences, University of Hyderabad, Hyderabad – 500046, India

²Department of Biotechnology, Kakatiya University, Warangal, A.P- 506009, India

SUMMARY

We have optimized a simple and rapid method for isolation of high quality genomic DNA from leaves of tomato, tobacco and rape seed. This protocol significantly minimizes time and the use of laboratory materials. The extracted DNA was suitable for PCR analysis. This method requires less than 1 mg of leaf tissue and is useful for transgene detection, genetic maps and other DNA based molecular analyses.

Keywords: Tomato, Tobacco, Rape Seed, DNA extraction, PCR analysis.

P. Mamidala and R.S. Nanna. A Simple and Rapid Method For DNA Extraction From Leaves of Tomato, Tobacco and Rape Seed. J Phytol1 (2009) 388-390

*Corresponding Author, Email: p.mamidala@yahoo.com

1. Introduction

Extraction of DNA from plants is prerequisite when hundreds of samples need to be analyzed rapidly for PCR analysis, genome mapping and molecular markers development. However the extraction of high-quality DNA from plants is tedious, time consuming and expensive because of different manipulation steps (1-5). In order to overcome these problems, we have optimized a simple and rapid method for high-quality DNA isolation from leaf tissues. This protocol was used as part of molecular cloning of antiporter gene conferring resistance to salt stress to minimize the time and the use of laboratory materials.

2. Materials and Methods

Plant Material

Transgenic plants were developed through *A.tumefaciens* mediated genetic transformation harboring the vacuolar antiporter (AtNHX1) gene in tomato, tobacco and rape seed (6-8). The leaves from these plants growing in the green house were used for DNA extraction to assay transgene possession.

Reagents and Chemicals:

- DEB (DNA Extraction Buffer : 1.0M Tris HCl, pH 8.0 and 0.5 M EDTA, pH 8.0;10% (w/v) SDS; 5M NaCl and 0.2% Mercaptoethanol)

- 5M Potassium acetate

DNA Isolation Protocol

- A piece of fresh leaf (approx 3-5 mg) was placed in a centrifuge tube/ependorf
- Add 400µL of DEB and leaf tissue was crushed with glass rod against the tube walls.
- To this add 50µL of 5M Potassium acetate
- Centrifuge at 16K rpm for 30 seconds
- Carefully transfer 10µL of supernatant to an eppendorf.
- To this add 990µL of sterile nanopure water

Directly this DNA was used as template for Polymerase Chain Reaction (PCR) amplification.

PCR amplification and DNA analysis

0.5µL of DNA extraction solution was added to 24.5µL of PCR mixture containing 1.5 µL 50mM MgCl₂, 2.5 µL of FP and RP (10 pmol/ µL), 2.5 µL PCR-buffer (Invitrogen), 0.5 µL 10mM dNTPs, 0.2 µL Taq Polymerase (Invitrogen), and 14.8 µL sterile double distilled water. PCR aliquot (25 µL) was centrifuged for a short spin. A thermocycler gradient (MJ Research Gradient Cycler) was used for thermal cycling as follows: 4 min at 94°C, followed by 3 cycles (6 times) of 50 Seconds at 94°C, 45 min at 50°C, 1 min at 72°C; Step II for 29 min; 10 min at 72°C; 10 min at 4°C. Primer sequences for AtNHX1 gene were as follows:

FP: 5' TTGGGTGATTATCTTGCTATTGG 3'

RP: 5' CGGCCCTTGTAACCTTGTTGTA 3'

PCR products (5-20 µL) were analysed with 1.7% agarose gel electrophoresis, stained with ethidium bromide, and visualized with UV transilluminator.


3. Results and Discussion

DNA isolated from fresh leaves was evaluated by NanoDrop ND-1000 UV-Vis Spectrophotometer. This procedure yielded 392ng/µl of high molecular weight DNA. A ratio of absorbance at 260nm to that of 280nm of 1.80-1.81 indicated the purity of DNA obtained (Table 1). PCR products were electrophoresed on agarose and amplified fragment of 800 bp was observed (Fig. 1). Unlike other high quality plant genomic DNA extraction protocols (1,3,4,5), this method does not include the use of liquid nitrogen or freeze drying for initial grinding of the leaf tissue. This protocol reduces the times of chloroform extraction by directly using Potassium acetate to remove contaminating proteins in the same step.

Table 1. Quantified DNA after isolation from leaves of transgenic plants using simple and rapid DNA extraction method

| Transgenic Plant | DNA yield (ng/µg of sample) | A260/A280 |
|-----------------------|-----------------------------|-----------|
| Tomato | 354 | 1.81 |
| Tobacco | 392 | 1.82 |
| <i>Brassica napus</i> | 348 | 1.80 |

Fig. 1: A representative picture showing the amplified products of AtNHX1 (800bp) in DNA extracted samples of transgenic plants. Lane 1 (DNA Ladder); Lane 2&9 (positive control – plasmid carrying AtNHX1); Lane 3-4 (Tomato), Lane 5-6 (Brassica napus), Lane 7-8 (Tobacco)



1 2 3 4 5 6 7 8 9

These alterations significantly minimize the time and the use of laboratory materials. One person can process as many as hundred DNA samples within one hour. The extracted DNA could be amplified by means of PCR both before and after storage at 4°C. The present protocol for DNA extraction is stable and suitable for PCR and other molecular techniques.

Acknowledgements

PM gratefully acknowledges UGC, New Delhi for financial assistance in the form of **Dr D S Kothari Post Doctoral Fellowship 2009** and Management of Vaagdevi College of Engineering for permitting to avail the same.

References

1. Murray M.G., Thompson W.F., 1980. Rapid isolation of high molecular weight DNA. *Nucleic Acids Res.* 8:4321-4325
2. Rogers S.O., Bendich A.J., 1985. Extraction of DNA from milligram amounts of fresh herbarium and mummified plant tissues. *Plant Mol Biol*, 5:69-76
3. Doyle J.J., Doyle J.L., 1987. A rapid DNA isolation procedure from small quantities of fresh leaf tissues. *Phytochem Bull.* 19:11-15
4. Doyle J.J., Doyle J.L., 1990. Isolation of plant DNA from fresh tissue. *Focus* 12:13-15
5. Huang J., Ge X., Sun M., 2000. Modified CTAB protocol using a silica matrix for isolation of plant genomic DNA. *BioTechniques*, 28:432-434.
6. Zhang H.X., Blumwald E., 2001. Transgenic salt-tolerant tomato plants accumulate salt in foliage but not in fruit. *Nat. Biotechnol* 19: 765-768.
7. Zhang H.X., Hodson J.N., Williams J.P., Blumwald E., 2001. Engineering salt-tolerant *Brassica* plants: Characterization of yield and seed oil quality in transgenic plants with increased vacuolar sodium accumulation. *Proc. Natl. Acad. Sci. USA* 98: 12832-12836.
8. Praveen M., 2006. Engineering abiotic stress resistance in tomato using cultivated tomato PhD thesis, Kakatiya University, Andhra Pradesh, India