



REGULAR ARTICLE

INFLUENCE OF LONG-TERM USE OF ORGANIC AND INORGANIC NUTRIENTS ON HLB DISEASE OF WHEAT UNDER RICE-WHEAT CROPPING PATTERN

Deepak Bhandari*

National Wheat Research Center, Nepal Agricultural Research Council, Nepal

SUMMARY

Helminthosporium leaf blight (HLB) is one of the most destructive diseases of wheat in terai (plain) region of Nepal. This study was conducted to determine the effect of long-term application of various levels of Nitrogen (N), Phosphorus (P), Potassium (K) and organic nutrients on HLB disease of wheat. The experiment was super imposed on long-term fertility experiment conducted in RCB design with three replications under rice-wheat cropping pattern at National wheat research program (NWRP), Bhairahawa, Nepal. The treatments include various combinations of 100 kg Nitrogen/ha, 0 and 60 kg Phosphorus/ha and 0, 50 and 100 kg Potash /ha with or without Sesbania (*Sesbania cannabina*) and Farm Yard Manure (FYM). Area under disease progress curve was calculated using the disease scores and yield parameters were recorded. Among the inorganic nutrients, long-term deficiency of potash in the field markedly enhances the HLB disease. Application of phosphorus in various doses increased the HLB severity in absence of potash. Regular application of at least 50 kg potash/ha in wheat noticeably reduced the HLB disease. Regular use of FYM 10 ton/ha for long term considerably reduced the HLB disease but long term use of Sesbania as green manure hadn't any impact on HLB severity. Inclusion of potash in nutrients considerably increased the yield and thousand-grain weight (TGW) of wheat. In long term, balanced use of inorganic and organic nutrients with special attention on regular inclusion of potash and FYM in the nutrient combinations is vital in suppression of HLB disease under rice-wheat cropping pattern.

Key words: AUDPC, FYM, Helminthosporium leaf blight, Phosphorus, Potassium, Sesbania

Deepak Bhandari. Influence of Long-term Use of Organic and Inorganic Nutrients on HLB Disease of Wheat under Rice-wheat Cropping Pattern. J Phytol 2/2 (2010) 73-79.

*Corresponding Author, Email: deepak359@hotmail.com

1. Introduction

Rice -Wheat is the most important cropping pattern in terai (plain) areas of Nepal. HLB is one of the most destructive diseases of wheat in terai regions of Nepal. HLB generally occurs with complex infection of Spot blotch (caused by *Cochliobolus sativus* (Ito & Kurib.) Drechsler ex. Dastur (anamorph: *Bipolaris sorokiniana*) and Tan spot (caused by *Pyrenophora tritici repentis* (Died.) Drechsler (anamorph: *Drechslera tritici repentis*) in plain areas of Nepal (Sharma *et al.*, 2003). HLB prevails every year in the major wheat growing areas of the country due to the lack of durable and highly resistant cultivars, use of unbalanced

fertilizers and poor management conditions. The losses due to the disease are remarkable in many regions of indo-gangetic plains including Nepal (Duveiller and Gilchrist, 1993; Duveiller, 2002). The yield losses due to HLB in farmers' field of Nepal range from 20 - 25% (Saari, 1997; Duveiller, 2002). The pathogens are weak hemi bio-trophs, so cultural practices prominently influence the severity of disease (Ruckstuhl, 1997; Naitao & Yousan, 1997; Bocus and Shroyer, 1998; Mathieoson *et al.*, 1990). Disease severity is increased under stressed plant growth conditions caused by low or unbalanced soil nutrients level, potash deficiency, poor

management practices and high temperature and hot wind during the later stage of the crop (Ruckstuhl, 1997; Naitao and Yousan, 1997; Bhatta *et al*, 1997; Sharma *et al*, 2005). Resistant genotypes, seed treatment and foliar sprays with triazole fungicides are effective against the disease (Bhatta *et al*, 1997; Naitao and Yousan, 1997); however, except resistant genotypes, the other methods are not economically practicable for poor farmers of Nepal. Among several factors of cultural management, the role of plant nutrients is supreme in the suppression of HLB disease. Application of potash fertilizer reduces the HLB incidence significantly in both short and long term fertility conditions (Regmi *et al*, 2002; Sharma *et al*, 2005). Nepalese farmers use unbalanced and less amount of fertilizers in wheat, which results in lower yield due to the combined effect of nutrient deficiency and higher HLB disease.

Determination of role of various combinations of major inorganic and organic nutrients on HLB severity in long run could help to develop a durable and cost effective technology for HLB management. Therefore, this study was conducted to find out the effect of long-term use of various levels of nitrogen, phosphorus, potash, green manure and FYM on HLB disease of wheat.

2. Materials and Methods

The experiment was super imposed on long-term fertility experiment under rice-wheat cropping system conducted at National Wheat Research Program, Bhairahawa in the year 2003-04. The long-term fertility experiment has been continuously conducted since 1988 with slight modification of treatments in 1995. The design was RCB with three replications.

Genotype Nepal 297 was used in both the years. The final treatments after modification in 1995 include application of various combinations of 100 kg Nitrogen/ha, 0 and 60 kg Phosphorus/ha and 0, 50 and 100 kg Potash /ha with or without Sesbania and FYM. Sesbania was grown for 60 days and mixed on field before rice planting whereas, FYM 10 ton /ha was applied to wheat only. Half of the nitrogen and full dose of phosphorus and potash was applied as basal and remaining half dose of nitrogen was applied after 25 days of seeding as top dress. Ten main tillers in each plot were randomly selected and tagged prior to scoring disease intensity. Severity of disease was recorded by visually assessing the percent diseased leaf area on the flag leaf and penultimate leaf of tagged tillers. Three scores were recorded at 7 days intervals after the heading stage of the crop. Area Under Disease Progress Curve (AUDPC) was calculated following the formula given by Das *et al*, 1992. Yield & thousand-grain weight were recorded for analysis of yield features.

3. Results and Discussion

Analysis of variance

Analysis of variance disclosed that the treatments are differed significantly ($P < 0.01$) for HLB AUDPC, yield and thousand-grain weight (TGW) in both years (Table: 1). FYM had significant ($P < 0.01$) effect on suppression of HLB severity in both the years. Application of potash significantly ($P < 0.01$) reduced the HLB severity. Application of phosphorus in absence of potash significantly ($P < 0.01$) increased the disease severity. The effect of Sesbania before rice was non significant for HLB AUDPC.

Table 1. Combined mean HLB AUDPC on different treatments of long term soil fertility experiment conducted at Bhairahawa during 2003 and 2004

S.No.	Treatments	Combined mean AUDPC of 2003 & 2004 [†]
1	Sesbania before rice and 100:60:100 kg NPK /ha	483C
2	Sesbania before rice and 100:60:0 kg NPK /ha	1027A
3	Sesbania before rice and 100:60:50 kg NPK /ha	571C
4	NPK 100:60:0 kg/ha	1052A
5	NPK 100:60:50 kg/ha	516C
6	NPK 100:60:100 kg/ha	536C
7	FYM 10 t/ha in wheat and 100:60:0 kg NPK /ha	601C
8	FYM 10 t/ha in wheat and 100:60:50 kg NPK /ha	437C
9	FYM 10 t/ha in wheat and 100:60:100 kg NPK /ha	453C
10	FYM 10 t/ha in wheat and 100:0:0 kg NPK /ha	463C
11	Sesbania before rice and 100:0:0 kg NPK /ha	759B
12	NPK 100:50:0 kg /ha	1035A
13	NPK 100:100:0 kg/ha	1120A
14	NPK 100:0:0 kg/ha	840B


[†]Means in a column followed by the same letter do not differ significantly at P = 0.05 according to Duncan's multiple range tests

Effects of Potash on HLB

Application of potash significantly reduced HLB severity on wheat under long-term rice - wheat cropping pattern (Table-1). Significantly lower disease in the treatments 100:60:100 and 100:60:50 kg NPK/ha than the treatment 100:60:0 kg NPK/ha signify the impact of potash fertilizer on suppression of HLB disease under long-term fertility conditions (Figure-1). This result supports the outcomes of various similar previous studies (Sharma *et al*, 2005, Regmi *et al*, 2002). Thin cell walls, weakened stalks and stems, smaller and shorter roots, and accumulation of unused nitrogen caused by potassium deficiency enhance the disease infection (Anonymous, 1998). Similarly, treatment 100:60:100 kg NPK/ha and 100:60:50 kg NPK/ha were statistically at par which suggest that the use of 50 kg of potash /ha is sufficient to suppress the incidence of HLB under long-term fertility conditions (Figure-1).

Likewise, the non-significant differences for HLB AUDPC among treatments Sesbania + 100:60:50, FYM + 100:60:50 and 100:60:50 indicate that application of balance dose of inorganic fertilizers is sufficient to reduce the HLB severity.

Figure1. Effect of potash on HLB of wheat under long-term fertility (R-W) conditions during 2003 & 2004




Effects of Phosphorus on HLB

The effect of Phosphorus was adverse on HLB disease in absence of potash (Table: 1). HLB incidence was higher in treatments 100:100:0, 100:60:0 and 100:50:0 in both the years. Significantly higher AUDPC in treatments having only N, P than the treatments having NPK and only N indicate the adverse effect of P on HLB disease (figure-2); however, the reasons behind this is not clear and need further investigation. This result agrees with the results of Gharti *et al*, 2002.

Similarly, significantly less HLB in the treatments Sesbania + 100:0:0 and only 100:0:0 NPK kg/ha than treatments 100:50:0,

100:60:0 and 100:100:0 NPK kg/ha suggests that for the suppression of HLB disease, application of nitrogen only should be prioritized over application of nitrogen and phosphorus. Likewise, statistically at par HLB incidence in treatments 100:50:0, 100:100:0 and 100:60:0 point out that the levels of phosphorus in absence of K haven't any effect on HLB severity. This result supports the previous findings that disease caused by *Cochliobolus sativus* is not affected by the addition of phosphorus (Bailey *et al*, 1980).

Figure 2. Effect of phosphorus on HLB of wheat under long-term fertility (R-W) conditions during 2003 & 2004



Effects of FYM on HLB


The significant and lowest AUDPC in the treatments having FYM and FYM + Potash proves the paramount role of these two nutrients on HLB suppression (Table-1). The least HLB disease in treatment having FYM 10 tons/ha + Potash may be partially due to the reduction of primary inoculums present in the soil, because application of cow dung, urine and high levels of organic matter inhibit the conidial germination of *B. sorokiniana* (Akhtar *et al*, 2006).

Likewise, the non-significant differences for HLB severity among 100:60:100 NPK kg/ha, 100: 60:50 NPK kg/ha and FYM 10 t/ha + 100:60:0 NPK kg/ha point out that application of FYM 10 tons/ha could suppress the HLB severity equal to the application of 50-100 kg potash/ha. This result agrees with similar previous reports, which advocated that organic materials reduce disease incidence caused by many plant pathogenic bacteria, fungi, and

nematodes (Abawi and Widmer, 2000; Conn and Lazarovits, 2000; Lazarovits *et al.*, 2001).

Significantly less disease in FYM 10 t/ha + 100:0:0 NPK kg/ha than the treatments Sesbania + 100:0:0 NPK kg/ha and only 100:0:0 NPK/ha confirms the sole effect of FYM on HLB suppression (figure- 3). Similarly, significantly lower disease in FYM 10 ton/ha + 100:60:0 NPK/ha as compare to treatments Sesbania before rice + 100:60:0 NPK kg/ha and only 100:60:0 NPK kg/ha suggest the importance of FYM in HLB management under different nutrient combinations.

Figure 3. Effect of FYM on HLB of wheat under long-term fertility (R-W) conditions at Bhairahawa during 2003 & 2004



Effects of Sesbania on HLB

There wasn't any impact of long-term application of Sesbania before rice on HLB disease of wheat in any nutrient combinations (figure-4). Sesbania hadn't any effect on HLB severity when it was added on the treatments containing all the three major elements such as 100:60:50 NPK kg/ha and 100:60:100 NPK kg/ha. Similarly, the addition of Sesbania on the treatment having nitrogen and phosphorus only (100:60:0 NPK kg/ha) also hadn't any influence on HLB severity. Present results do not support the previous conclusions of Nikonorova, (1997) and Davis *et al.*, (1996), in which green manuring decreased the infection level of some soil borne diseases. However, based on present results and previous conclusions, it can be suggested that green manures may have less impact on foliar diseases than soil borne diseases.

Likewise, the higher and at par levels of HLB AUDPC in treatments Sesbania +100:0:0

NPK kg/ha and only 100:0:0 NPK kg/ha suggest the lack of influence of Sesbania on HLB severity when it is applied in the field having nitrogen only. The results also supports that application of higher amount of only Nitrogen either in organic or inorganic form raised the severity of HLB because application of Nitrogen level above 80 kg /ha induces the foliar disease incidence due to denser crop canopy (Wall *et al*, 1990).

Effects of various nutrients on yield of wheat

The yield of wheat was significantly different among the treatments (table: 2); however, the variation in yield was not solely due to the levels of HLB disease. The variations in yield were due to the combined effect of levels of nutrients and levels of HLB disease. Lower wheat yield and TGW with higher HLB AUDPC (figure-5) confirms the previous reports which disclosed the negative correlations of HLB AUDPC with yield and TGW (Mahto, 1999).

Figure 4. Effect of Sesbania on HLB of wheat in different nutrient combinations under long-term fertility (R-W) conditions at Bhairahawa during 2003 & 2004




Table 2: Combined mean yield and TGW on different treatments under long-term fertility condition at Bhairahawa during 2003 and 2004

S.No.	Treatments	Combined mean yield (kg/ha)†	Combined mean TGW (gm)†
1	Sesbania before rice and 100:60:100 NPK kg/ha	2073ab	38.44a
2	Sesbania before rice and 100:60:0 NPK kg/ha	642c	21.77c
3	Sesbania before rice and 100:60:50 NPK kg/ha	2076 ab	36.54 ab
4	NPK 100:60:0 kg/ha	595c	20.98c
5	NPK 100:60:50 kg/ha	2421 ab	39.36a
6	NPK 100:60:100 kg/ha	2102 ab	35.21 ab
7	FYM 10 t/ha in wheat and 100:60:0 NPK kg/ha	1870b	32.45b
8	FYM 10 t/ha in wheat and 100:60:50 NPK kg/ha	2601a	37.47 ab
9	FYM 10 t/ha in wheat and 100:60:100 NPK kg/ha	2284 ab	39.67a
10	FYM 10 t/ha in wheat and 100:0:0 NPK kg/ha	1903b	36.13ab
11	Sesbania before rice and 100:0:0 NPK kg/ha	250c	22.04c
12	NPK 100:50:0 kg/ha	640C	20.92C
13	NPK 100:100:0 kg/ha	481C	22.75C
14	NPK 100:0:0 kg/ha	342C	23.14C


†Means in a column followed by the same letter do not differ significantly at P = 0.05 according to Duncan’s multiple range tests

Treatments containing 50 kg potash per ha had significantly higher yield and TGW

than the treatments that lack the potash. Treatments with only inorganic nitrogen and

phosphorus had significantly lower yield and significantly higher HLB AUDPC than treatments having inorganic nitrogen, phosphorus and potash. This result verifies the role of potash on wheat yield and HLB disease, which has been already explained by several previous workers (Sharma *et al*, 2005, Regmi *et al*, 2002, Gharti *et al*, 2002).

Figure 5. Relationship of HLB AUDPC with wheat yield and TGW in different nutrient combinations under long-term fertility (R-W) conditions at Bhairahawa during 2003 & 2004



4. Conclusion

In long term, balanced use of inorganic and organic nutrients with special attention on regular inclusion of potash and FYM in the nutrient combinations is vital in suppression of HLB disease under rice-wheat cropping pattern. Long term application of only nitrogen and phosphorus fertilizers boosts the HLB disease severity, where as application of 50 kg potash/ha in addition of nitrogen and phosphorus fertilizers considerably reduces the HLB disease severity. Similarly, Long-term use of FYM effectively suppresses HLB disease in wheat; where as long term applications of Sesbania couldn't suppress the intensity of HLB disease.

Acknowledgement

I would like to thank E. Duveiller (CIMMYT, SARO) Kathmandu for their support, encouragement and guidance. The support of A. P. Regmi and NWRP is duly acknowledged.

References

- Abawi, G.S., T.L. Widmer. 2000. Impact of soil health management practices on soil borne pathogens, nematodes and root diseases of vegetable crops. *Appl. Soil Ecol.* 15: 37-47.
- Akhtar, N., M. F. Begum, S. Alam and M. S. Alam. 2006. Inhibitory effect of different plant extracts, cow dung and cow urine on conidial germination of *Bipolaris sorokiniana*. *Journal of bio-science.*14: 87-92
- Anonymous, 1998. *Better Crops.* Vol. 82, No. 3, pp 37
- Bailey, L.D., H. Ukrainetz, and D.R. Walker. 1980. Effect of phosphorus placement on crop uptake and yield. *Proc. Western Canada Phosphate Symposium. Alberta Soil Science Workshop.* pp. 200-229. Calgary, Alberta.
- Bhatta, M. R., D. R. Pokhrel, R. N. Devkota, H. J. Dubin, A. Mudvari, H. P. Bimb, B. R. Thapa, B. P. Sah and D. Bhandari. 1997. Breeding for resistance to Helminthosporium Blights in Nepal: Strategies and Genetic Gains. In: E. Duveiller, H. J. Dubin, J. Reeves and A. McNab (eds.), *Helminthosporium Blight of wheat: Spot blotch and Tan spot.* CIMMYT, Mexico, D. F. pp: 188 - 195
- Bockus, W.W. and J.P. Shroyer.1998. The impact of reduced tillage on soil borne plant pathogens. *Annual Review of Phytopathology.* 36:485-500
- Conn, K.L. and G. Lazarovits. 2000. Soil factors influencing the efficacy of liquid swine manure added to soil to kill *Verticillium dahliae*. *Can. J. Plant Pathol.* 22:400-406.
- Das, M.K., S. Rajaram, C.C. Mundt and W.E. Kronstad. 1992. Inheritance of slow rusting resistance to leaf rust in wheat. *Crop Science.* 32:1452-1456
- Davis, J.R., O.C. Huisman, D.T. Westermann, S.L. Hafez, D.O.Everson, L.H. Sorensen and A.T. Schneider. 1996. Effects of green manures on *Verticillium wilt* of potato. *Phytopathology.* 86:444 - 453
- Duveiller, E. and L. Gilchrist 1993. Production constraints due to *Bipolaris sorokiniana* in wheat. Current situation and future prospects. In: D.A.Saunders

- and G.P.Hettel (eds), Wheat in heat stress environments: Irrigated dry areas and rice wheat farming system. Mexico, CIMMYT, D.F. pp: 343-352
- Duveiller, E., 2002. Helminthosporium Blight of wheat: Challenges and strategies for a better disease control. In: Advances of Wheat breeding in China. Proceedings of the first National Wheat Breeding Conference, 10-12 May, 2002, Jinan, Shandong, People Republic of China, China Science and Technology Press. pp: 57-66
- Gharti, D.B., M.P. Sah, S.K.Gami, and K.P. Bhurer. 2002. Effect of various organic and inorganic nutrients on foliar blight of wheat. In Proceedings of Wheat Research Papers presented at 25th National Winter Crops Research Workshop, Volume 1: Wheat. pp: 142- 147
- Lazarovits, G., M.Tenuta, K.L. Conn, 2001. Organic amendments as a disease control strategy for soil borne disease of high-value agricultural crops. *Australians. Plant Pathol.* 30:111-117
- Mahto, B. N. 1999. Management of *Helminthosporium* leaf blight of wheat in Nepal. *Indian Phytopath.* 52(4): 408 - 413
- Mathieson, J.T., C.M. Rush, D. Brodovsky, L.E. Clark and O.R. Jones.1990. Effects of tillage on common root rot of wheat in Texas. *Plant Disease.* 74:1006-1008
- Naitao, Chang and Wu yousan 1997. Incidence and current management of spot blotch of wheat in China. In: E. Duveiller, H.J. Dubin, J. Reeves and A. McNab (eds). *Helminthosporium blight of Wheat: Spot blotch and Tan spot.* CIMMYT, Mexico, D.F. pp: 119-125
- Nikonorova, A.K. 1997. Phytosanitary effect of combined application of green manure and antagonistic bacterium *Bacillus subtilis* on *B. sorokiniana* In: E. Duveiller, H.J. Dubin, J. Reeves and A. McNab (eds), *Helminthosporium blight of wheat: Spot blotch and Tan spot.* CIMMYT, Mexico, D.F. pp: 359-363
- Regmi, A.P., J.K.Ladha, E.M. Pasuquin, H. Pathak, P.R. Hobbs, L.L. Shrestha, D.B. Gharti and E. Duveiller. 2002. The role of potassium in sustaining yields in a long term rice-wheat experiment in the Indo-Gangetic plains of Nepal. *Biol. Fertil. Soils.* 36: 240 - 247
- Ruckstuhl, M. 1997. Population structure and epidemiology of *B.sorokiniana* in the rice wheat cropping pattern in Nepal. In: E. Duveiller, H.J. Dubin, J. Reeves and A. McNab (eds), *Helminthosporium blight of wheat: Spot blotch and Tan spot.* CIMMYT, Mexico, D.F. pp: 88 -106
- Saari, E. E. 1997. Leaf blight disease and associated soil borne fungal pathogens of wheat in South and South East Asia. In: E. Duveiller, H.J. Dubin, J. Reeves and A. McNab (eds), *Helminthosporium blight of wheat: spot blotch and tan spot.* CIMMYT, Mexico.D.F. pp: 37-51
- Sharma, R.C., S.M. Shrestha, E. Duveiller. 2003. Incidence of *Bipolaris sorokiniana* and *Pyrenophora tritici repentis* on wheat in the lowlands of Nepal. In: Rasmusseen, J.B., Friesen, T.L., Ali, S. (eds.), *Proceedings of the fourth International wheat Tan spot and spot blotch workshop, North Dakota state university, Fargo, USA,* pp. 122-127
- Sharma, S., E. Duveiller, R. Basnet, C. B. Karki and R. C. Sharma. 2005. Effect of Potash fertilization on *Helminthosporium* leaf blight severity in wheat, and associated increase in grain yield and kernel weight. *Field Crop Research.* 93: 142-150
- Wall, P.C., P. R. Hobbs, D. A. Saunders, K. D. Sayre and D.G.Tanner. 1990. Wheat crop management in the warmer areas: A review of issues and advances. In: D.A. Saunders (ed), *Wheat for the non traditional warm areas.* CIMMYT, Mexico, D.F. pp: 225-241.