

# Validation of antibacterial activity of Saponin against diarrheagenic *E.coli* isolated from leaves and bark of *Acacia arabica*

\*Deboshree Biswas and M.G Roymon

Department of Microbiology and Biotechnology, ST. Thomas College, Ruabanda, Bhilai, India.

## Abstract

The leaf and bark extract of *Acacia arabica* constitute large amount of saponin (sapogenin aglycon) along with other polyphenolic compounds. The isolated sapogenin aglycon was tested against *E.coli* O22 associated with diarrheal infections in domesticated animals along with *E.coli* MTCC 723 strain. Both well-diffusion and broth micro dilution technique was used to detect antibacterial activity. Synergistic interaction was observed on combined administration of saponin with Chloramphenicol to inhibit tested organism.

**Keywords:** Sapogenin aglycon, Broth micro dilution technique

## INTRODUCTION

Bioactive compounds are the safest and more effective alternative than synthetically produced antimicrobial agents (Banso, 2009, Tang *et al.*, 2010). In recent years, the use of plants and plant based products gain much importance for treatment of various infections (Bagyalakshmi *et al.*, 2009). The use of plant derived bioactive analogues serves as the safest agent to cure bacterial infections associated with multi drug resistant strains. Saponins are the glycosides present in almost all plants belonging to family Leguminosae. It is also present in other plants belonging to different families. The present study deals with antibacterial and synergistic activity of sapogenin aglycon isolated from *Acacia arabica* against *E.coli*.

## MATERIALS AND METHODS

### Culture collection and identification

*E.coli* O22 was obtained from stool sample of diarrheagenic calf and standard strain *E.coli* MTCC 723 was purchased from IMTECH, Chandigarh

### Collection and preparation of crude extract

Leaves and bark of *Acacia arabica* was collected during November in specimen voucher and thoroughly washed and shade dried. All plant materials were separately pulverized for preparation of extracts. Crude methanolic extract was prepared by the method recommended by Johnson *et al.*, (2008).

### Separation and purification of saponin

Crude methanolic extract of leaf and bark was fractioned using column chromatography as per method recommended by Saleem *et al.* (2009) Sundaram and Mitra, (2006). Purified bioactive compounds were obtained by subjecting the fractions to Preparative HPLC (LC-18, Shimadzu, Japan) with C-18 column (20mm x 250mm). The mobile phase used was acetonitrile: water (7:3) at a flow rate of 10mL/min. The injection volume of fractions of leaf and bark extract was of 10  $\mu$ l. The isolated saponin was evaporated to dryness under vacuum for detection of antibacterial activity (Kyung *et al.*, 2007). The mass spectrum of isolated saponin was analyzed using LC analysis Perkin Elmer Series (Japan) through positive electrospray ionization mass spectrometer (ESI-MS). The isocratic mobile phase was 100% methanol with flow rate of 10mL/min (Theerasin and Baker, 2009).

### Antibacterial activity

Antibacterial activity was determined by well diffusion method (Cock, 2008) and broth microdilution method (Kashikar & George, 2006, Mann *et al.*, 2009) with some modifications.

### Synergistic activity

*In vitro* synergistic activity between saponin and chloramphenicol was performed using well diffusion method (Odunbaku *et al.*, 2008).

## RESULTS AND DISCUSSION

*Acacia arabica* is used widely for treatment of various infections including treatment of diarrhea and dysentery (Chopra *et al.*, 1992). There are reports on antibacterial activity of leaf and bark extract of *Acacia arabica* on *E.coli* (Dabur *et al.*, 2007). However, a very limited work was carried out in field of ethnoveterinary medicines to cure diarrheal infections in animals. Lans *et al.*, (2007) documented ethnoveterinary medicine (EVM) is the scientific term used for treatment of animals by using plants. In addition, very limited work was carried out on concurrent administration of both herbal drugs and conventional antibiotics to cure infections. The present study was adopted to assess antibacterial activity of leaf and

Received: Jan 12, 2012; Revised: Feb 24, 2012; Accepted: March 15, 2012.

\*Corresponding Author

Deboshree Biswas

Department of Microbiology & Biotechnology, ST. Thomas College, Ruabanda, Bhilai, India.

Tel: +91-9691357003; Fax: +91-788-2272270

Email: [biswasdeboshree19@gmail.com](mailto:biswasdeboshree19@gmail.com)

bark extract derived steroidal saponin aglycon using well diffusion and broth micro dilution technique against diarrheagenic *E.coli* responsible for infection in animals. Among both methods broth microdilution method was the most reliable method for accurate quantification of antibacterial activity of saponin. In addition, another aim of the study was to investigate interaction of saponin with convectional antibiotics for more effective inhibition of tested organism. Table: 1 represents antibacterial activity of compound by well diffusion method on the basis of diameter of zone of inhibition and Table: 2 represent the activity of compound on the basis of MIC performed by broth microdilution assay using INT dye. The findings on antibacterial effectiveness of saponin is in agreement with the findings of other researchers (Arabski *et al.*, 2012, Hassan *et al.*, 2010) who documented effectiveness of saponin against *E.coli*.




Fig 1. Preparative HPLC Chromatogram of leaf extract fraction




Fig 2. Preparative HPLC Chromatogram of bark extract fraction




Fig 3. HPLC Chromatogram of Standard Saponin




Fig 4. Mass spectrum of leaf extract derived Saponin




Fig 5. Mass spectrum of bark extract derived Saponin




Fig 6. Mass spectrum of Standard Saponin

Table 1. Mean IZD± SEM (in mm) of saponin from leaf and bark extracts of *Acacia arabica*

<i>E.coli</i>	Leaf extract derived saponin	Bark extract derived saponin	Chloramphenicol
O22	8.6±0.6	8	0
MTCC 723	6	4	17.6± 0.6

Table 2. MIC of leaf & bark extract derived Saponin (in µg)

<i>E.coli</i>	Leaf extract derived saponin	Bark extract derived saponin
O22	8.3 ±1.6	1.26±0.3
MTCC 723	6.6±0.6	1.57±0.3

Table 3. Mean IZD± SEM (in mm) of combined effect of leaf & bark extract derived Saponin with Chloramphenicol

<i>E.coli</i>	Leaf extract derived saponin + Chloramphenicol	Bark extract derived saponin + Chloramphenicol
O22	16.6±0.6	11.6±0.6
MTCC 723	12± 1.1	26.6±0.6

The effect of combination of saponin with Chloramphenicol was presented in Table: 3. Combination of Chloramphenicol with leaf extract derived saponin showed synergistic effect on *E.coli* O22. However, on combination of Chloramphenicol to bark extract derived saponin additive effect on the same strain was noticed (Table: 3). In addition, strongest *in vitro* synergistic interaction was observed on *E.coli* MTCC 723 on combined administration of Chloramphenicol and leaf and bark extract derived saponin. Kim *et al.*, (1987) documented synergistic interaction between ginseng saponin and antibiotics to inhibit various pathogens which supports our findings on interaction

REFERENCES

[1] Arabski, M.; We,gierek-Ciuk, A.; Czerwonka, G.; Lankoff, A. and Kaca, W. 2012. Effects of saponins against clinical *E. coli* strains and eukaryotic cell line. *Journal of Biomedicine and Biotechnology*, 1:1-6.

- [2] Banso, A. 2008. Phytochemical and antibacterial investigation of bark extracts of *Acacia nilotica*. *Journal of Medicinal Plants Research*, 3: 82-85.
- [3] Bagyalakshmi, B.; Sridhar, D. and Ponmurugan, P. 2009. Antimicrobial activity of important Indian medicinal plants against pyogenic infections. *Journal of Phytology*, 1: 391–396.
- [4] Chopra, R.N.; Nayer, S.L. and Chopra I.C. 1992. *Glossary of Indian Medicinal Plants*. 3rd Edition, Publication and Information Directorial (CSIR), New Delhi India, pp. 2.
- [5] Cock, I.E. 2008. Antimicrobial activity of *Aloe barbadensis* miller leaf gel components. *The Internet Journal of Microbiology*, 4 :2
- [6] Dabur, R.; Gupta, A.T.; Mandal, K.; Singh, D.D.; Bajpai, V.; Gurav, A.M. and Lavekar, G. S. 2007. Antimicrobial activity of some Indian medicinal plants. *African Journal of Traditional Complementary and Alternative Medicines*, 4:313–318.
- [7] Hassan, S.M; Haq, A.U.; Byrd, J.A.; Berhow, M.A.; Cartwright, A.L. and Bailey, C.A. 2010. Haemolytic and antimicrobial activities of saponin-rich extracts from guar meal. *Food Chemistry*, 119: 600-605.
- [8] Johnson, M.; Maridass, M. and Irudayaraj, V. 2008. Preliminary phytochemical and antibacterial Studies on *Passiflora edulis*. *Ethnobotanical Leaflets*, 12: 425-432
- [9] Kashikar, N.D. and George, I. 2006. Antibacterial activity of *Cissus quadrangularis* Linn. *Indian Journal of Pharmaceutical Sciences*, 68: 245-247.
- [10] Kim, H.S.; Han, S.S.; Oh, K.W.; Jeong, T.S. and Nam, K.Y. (1987). Effects of ginseng saponin on antimicrobial activities of some antibiotics. *Korean Journal of Mycology*, 15: 87-91.
- [11] Kyung, K.H.; Woo, Y.H.; Kim, D.S.; Park, H.J. and Kim, Y.S. 2007. Antimicrobial activity of an edible wild plant, *Apiifolia Virgin's Bower (Clematis apiifolia DC)*. *Food Science Biotechnology*, 16: 1051 – 1054.
- [12] Lans, C.; Turner, N.; Khan, T.; Brauer, G. and Boepple, W. 2007. Ethnoveterinary medicines used for ruminants in British Columbia, Canada. *Journal of Ethnobiology and Ethnomedicine*, 3: 1-22.
- [13] Mann, A.; Amupitan, J.O.; Oyewale, A. O.; Okogun, J. I. and Ibrahim, K. 2009. Antibacterial activity of terpenoidal fractions from *Anogeissus leiocarpus* and *Terminalia avicennioides* against community acquired infections. *African Journal of Pharmacy and Pharmacology*, 3: 22-25
- [14] Odunbaku, O.A.; Ilusanya, O.A. & Akasora, K.S. 2008. Antibacterial activity of ethanolic leaf extract of *Ficus exasperata* on *Escherichia coli* & *Staphylococcus aureus*. *Scientific Research & Assay*, 3: 562-564.
- [15] Saleem, A.; Engstrom, M.; Wurster, S., Juha-Matti Savolab and Pihlaja, K. 2002. Interaction of folk medicinal plant extracts with human  $\alpha$ 2-adrenoceptor subtypes. *Medicinal Plants of Pakistan* 3: 332-338.
- [16] Sundaram, R. and Mitra, S.K. 2007. Antioxidant activity of ethyl acetate soluble fraction of *Acacia arabica* bark in rats. *Indian Journal of Pharmacology*, 39: 33-38.
- [17] Tang, J.; Meng, X.; Liu, H.; Zhao, J. ; Zhou, L.; Qiu, M.; Zhang, X.; Yu, Z. and Yang, F. 2010. Antimicrobial activity of sphingolipids isolated from the stems of cucumber (*Cucumis sativus* L.). *Molecules*, 15: 9288-9297.
- [18] Theerasin, S. and Baker, A.T. 2009. Analysis and identification of phenolic compounds in *Dioscorea hispida* Dennst. *Asian Journal of Food and Agro-Industry*, 2: 547- 560.