

# Influence of indole-3-butyric acid and triazole compounds on the photosynthetic pigments and biochemical constituents of *Withania somnifera* (L.) Dunal

P. Sakthivel, R. Sridharan\*

Department of Botany, Plant Growth Regulation Lab, Annamalai University, Annamalai Nagar- 608 002, Chidambaram, Tamil Nadu, India

Received: 27.08.2015

Accepted: 16.09.2015

Published: 16.09.2015

## \*Address for

### Correspondence:

R. Sridharan, Department of Botany, Plant Growth Regulation Lab, Annamalai University, Annamalai Nagar- 608 002, Chidambaram, Tamil Nadu, India.  
Tel.: +91-04144-238887,  
Fax: +91-04144-238080.  
E-mail: sridharanbot@gmail.com

## ABSTRACT

*Withania somnifera* (L.) Dunal is popularly known as ashwagandha and also called Indian Ginseng and Winter Cherry. It is an important medicinal plant used in Indian System of Medicines like Ayurveda, Siddha, and Unani. The root of this plant has nutritional and medicinal values out of which health foods and herbal tonic are prepared. The root has high demand in pharmaceutical and nutraceutical industries so that it is essential to increase the root yield. This is new phenomenon to increase root yield as well as improve the biochemical constituents by using plant growth regulators. In the present investigation, the influence of indole-3-butyric acid (IBA) and triazole compounds viz., triadimefon (TDM) and propiconazole (PCZ) on the photosynthetic pigments and biochemical constituents of ashwagandha. Plants were treated with IBA 2.5 mg/L, TDM 20 mg/L, and PCZ 20 mg/L separately by soil drenching on 50, 90, and 130 days after sowing (DAS). Plants were analyzed randomly on 60, 100, and 140 DAS and its parameters like, photosynthetic pigments (total chlorophyll, carotenoids, anthocyanin, and xanthophylls) in leaf and biochemical constituents (starch, protein and amino acids) in leaf, stem and root organs of ashwagandha. It was determined that total chlorophyll, carotenoids, anthocyanin, and xanthophylls, starch, protein and amino acids content were increased in all the treatments. Among the treatments, triazole compounds showed beneficial due to the enhanced the photosynthetic pigments and increased biochemical contents higher level than followed by IBA treatment. From our results, it can be concluded that the triazole shows great significance application at low concentration could be a potential agronomical tool for successfully cultivation of this medicinally important root crops. Triazole compounds enhanced the photo-assimilate to shifting partition from leaves to roots and also alter mineral uptake and plant nutrition, This characters' can be employed to satisfy needs of enhanced the photosynthetic pigments and biochemical constituents in ashwagandha.

**KEY WORDS:** Ashwagandha, medicinal plant, photosynthetic pigments, biochemical constituents, indole-3-butyric acid, triazole

## INTRODUCTION

Medicinal plants have been considered since time immemorial as an alternative source of a wide range of chemical compounds including pharmaceuticals, flavors, fragrance, colors, and insecticides. These compounds are collectively known as secondary metabolites in medicinal and aromatic plants and used as the remedy for health problems (Hussain, 1991; Okigbo *et al.*, 2009). India has been known

to be a rich repository of medicinal plants. Several scientific studies conducted throughout the world have revealed and confirmed the dramatic medicinal properties by their inherent nature of containing various phytochemicals such as flavonoids, carotenoids, and alkaloids. In India's progress in improving the cultivation of traditional medicinal crops is a matter of pride for the nation. However, compared to other crops, little attention seems to have been given to medicinal plants which occupy a unique place in the

Indian socio-economy (Jakhar *et al.*, 2003). Therefore, many scientists have repeatedly advocated to undertake applied and fundamental research work on cultivation of medicinal plants, several of which may attain the position of important cash crops for Indian farmers in future (Singh and Tyagi, 2004). Ashwagandha (*Withania somnifera* (L.) Dunal, belongs to the family Solanaceae is commonly known as Indian Ginseng and Winter Cherry. It is the most important medicinal plant used in Indian System of Medicines (Ayurveda, Unani, and Siddha) and in fact, it is mentioned as an official plant drug in the Indian Pharmacopocia (Indian Pharmacopocia, 1985). It is described as an herbal tonic and health food in Vedas and considered as “Indian Ginseng.” In “Ayurveda” the root drugs are mostly used as an alternative empirical therapy for the treatment and clinical management of male infertility (Gupta *et al.*, 2013; Singh *et al.*, 2010; Kokate *et al.*, 2005; Puri, 2003; Kapoor, 2001). The plant has also been used as an anticancer, anti-stress, anti-inflammatory and immunomodulator (Rasool and Varalakshmi, 2006; Seartezini and Sparoni, 2000; Agarwal *et al.*, 1999) and natural antioxidant properties (Bhattacharya *et al.*, 1997; Palash *et al.*, 2010). Since there is a high demand for the medicinal plants in the pharmaceutical industry, it is essential to increase the commercial cultivation, Furthermore, to ensure proper supply of medicinal plants to the drug industries. It has become necessary that these plants be propagated properly and cultivated scientifically, as well as quality and quantity (Farooqi and Sreeramu, 2004).

This is a new phenomenon to increase plant productivity scientifically as well as improve the biochemical constituents by using synthetic chemical compounds. Plant growth regulators are widely used to modify canopy structure, yield and stress tolerance in many crop plants. Manipulating the crop morphology by using plant growth regulators also increases the utilization of solar radiation and alter assimilates distribution in favor of yield increments. Plant growth retardants are synthetic compounds, which represent the commercially most important group of plant growth regulators. In addition to other agronomic tools, synthetic plant growth regulators are increasingly used to modify growth, development and stress behavior and the qualitative and quantitative yield of crop plants. Over the past few years, several triazole derivatives, collectively described as sterol biosynthesis inhibitors, have been developed and used as fungicides, and they also have plant growth regulating properties. Triadimefon (TDM) (bayleton) propiconazole (PCZ)

(banner), paclobutrazole (Bonzi), and uniconazole (sumagic) are used as growth regulators or retardants. However, all of these products can exhibit both fungicidal and growth regulating properties to varying degrees (Fletcher *et al.*, 2000). The plant growth regulating properties of triazoles are mediated by their ability to alter the balance of important plant hormones including gibberellic acids, abscisic acid (ABA), and cytokinins (Fletcher and Hofstra, 1988). The plant hormones are organic substances in low concentrations regulates the growth and development. These substances belong to different classes have different physiological role in plants to modify, regulate and development. The naturally occurring plant growth substances include auxins, gibberellins, cytokinins, abscisic acid, and ethylene (Kakimoto, 2003). Triazoles and IBA have been successfully used to increase the growth and root yield of medicinal plants. Hence, the present study becomes effectual to evaluate the effect of triazole and IBA compounds on the photosynthetic pigments (total chlorophyll, carotenoids anthocyanin, and xanthophylls) and biochemical constituents (starch, protein, and amino acids) content of *W. somnifera*.

## MATERIALS AND METHODS

### Pot Culture Experiments and Plant Treatments

The seeds of ashwagandha (*W. somnifera* L.) variety “Jawahar Asgandh-20” were obtained from Horticultural College and Research Institute, Department of Medicinal and Aromatic Crops, Tamil Nadu Agricultural University, Coimbatore, India. Indole-3-butyric acid (IBA) was obtained from Sigma Chemicals, Bangalore. Bayleton™ (TDM) was obtained from Bayer India Ltd., Mumbai. Banner™ (PCZ) was obtained from Rallis India Ltd., Mumbai and they were used for this research study.

The pot culture experiments were conducted at the Botanical Garden and in Plant Growth Regulation Laboratory, Annamalai University during the months of January-June, 2012. The pots were filled with the mixture of Red soil + Sand + Farm yard manure in the ratio of 1:1:1 and each pot were filled with the 15 kg mixture. 25 seeds were sown in each pot and finally 3 seedlings were maintained under shade net. The experiment was conducted in completely randomized block design (CRBD) with six replications. Fertilizer was not used throughout the experiments. In preliminary experiments, 0.5, 1.5, 2.0, 2.5, 3.0, and 3.5 mg/L of IBA, 5, 10, 15, 20, 25, and 30 mg/L

of TDM, and PCZ were used for the treatment and to determine the optimum concentration. 2.5 mg/L of IBA, 20 mg/L of TDM and 20 mg/L of PCZ were found to increase the root dry weight significantly and in higher concentration they slightly decreased the growth and dry weight. Hence, 2.5 mg/L IBA, 20 mg/L TDM and 20 mg/L PCZ concentrations were used to determine the effect of these chemicals on the growth of *W. somnifera*. Each pot was treated with one liter of respective treatment (2.5 mg/L IBA, 20 mg/L TDM and 20 mg/L PCZ) separately and control plant was treated with one liter of tap water. Each pot has three plants and the treatments were given on 50<sup>th</sup>, 90<sup>th</sup>, and 130<sup>th</sup> days after sowing (DAS) by soil drenching. The Electrical conductivity of the soil was 0.21/dsm and pH was 7.5 after the treatments. The average temperature was 32/26°C (maximum and minimum) and relative humidity varied between 60 and 75% during the experimental period. Plants were harvested on 60<sup>th</sup>, 100<sup>th</sup> and 140<sup>th</sup> DAS and they were used for the determination of photosynthetic pigments and biochemical constituents of *W. somnifera*.

### Pigments Analysis

Total chlorophyll and carotenoid contents were extracted from the leaves and estimated by the method of Arnon (1949). Anthocyanin content was estimated following the method of Beggs and Wellmann (1985) and xanthophylls content was estimated by the method of Neogy *et al.* (2001). The results were expressed in milligram per gram fresh weight.

### Biochemical Analysis

Starch content was extracted from the leaf, stem and root and estimated by the method of Clegg (1956), protein content was estimated according to Bradford (1976) and amino acid content was estimated followed by the method of Moore and Stein (1948). The results were expressed in milligrams per gram fresh weight.

### Statistical Analysis

The experiment was conducted by CRBD. Statistical analysis was performed using one-way analysis of variance followed by Duncan's multiple range test. The values were expressed in mean  $\pm$  standard deviation for six samples in each group  $P \geq 0.5$  values were considered as significant.

## RESULTS AND DISCUSSION

### Influence of IBA, TDM, and PCZ on the Photosynthetic Pigments Content of Ashwagandha

Triazole and IBA treatments significantly enhanced the total chlorophyll content in leaves of *W. somnifera*. Among the triazole, PCZ, and TDM treatments increased the chlorophyll content to a higher level when compared to IBA treated plants [Table 1]. TDM and hexaconazole treatments increased the chlorophyll content in radish (Sridharan *et al.*, 2006b). TDM treatment increased the chlorophyll content in *Cucumis sativus* seedlings (Feng *et al.*, 2003) and in banana (Galal *et al.*, 2011). Paclobutrazol treated barley seedlings were increased total chlorophyll content (Sunitha *et al.*, 2004), carrot (Gopi *et al.*, 2007) and tomato (Still and Pill, 2004). Paclobutrazol treated leaves were dark green due to high chlorophyll a and b content in potato (Tekalign *et al.*, 2005). Sebastian *et al.* (2002) reported enhanced chlorophyll synthesis in *Dianthis caryophyllus* treated with paclobutrazol. Similar results were observed in PCZ treated *Amorphophallus campanulatus* (Gopi *et al.*, 2005). The increased chlorophyll content with the PCZ and TDM treatment may be due to the ability of triazole to enhance cytokinin production, which stimulates the chlorophyll biosynthesis in *W. somnifera* leaves. IBA treatment significantly enhanced the chlorophyll a & b and total chlorophyll content in grapevine cuttings (Kaur *et al.*, 2002). The total chlorophyll content increased in IBA treated *Berberis thunbergii* (Pacholczak, 2006) and also in *Pisum sativum* (El-Shraiy and Hegazi, 2009). The effect of IBA on the chlorophyll concentration has been mentioned by several workers (Ludwig-Muller, 2000; El-Wahed *et al.*, 2006).

The amount of carotenoid content of the leaves of *W. somnifera* increased with all the stages of growth. Triazole and IBA treatments increased significantly the carotenoid content when compared to control plant. Among the treatments, PCZ and TDM increased the carotenoid content larger extent when compared to IBA [Table 1]. TDM treatment induced higher level of carotenoid content in cucumber seedlings (Feng *et al.*, 2003), cowpea (Gopi *et al.*, 1999), *Catharanthus roseus* (Jaleel *et al.*, 2008) and also in maize plants (Kaya *et al.*, 2006). Paclobutrazol treatment increased the carotenoid content in *Raphanus sativus* (Sankari *et al.*, 2006). Similar results were observed in carrot (Gopi *et al.*, 2007), *Solenostemon rotundifolius* (Kishorekumar *et al.*, 2007) and barley seedlings (Sarkar *et al.*, 2004). Sunitha *et al.* (2004) reported that the barley seedlings treated with paclobutrazol appeared greener and thicker due to increased pigment contents. Similar results

**Table 1: Influence of IBA, TDM and PCZ on the total chlorophyll, carotenoid, anthocyanin and xanthophyll contents of ashwagandha leaf**

Photosynthetic pigments	Growth stages (DAS)	Control	IBA	TDM	PCZ
Total chlorophyll (mg/g FW)	60	1.261±0.096 <sup>a</sup>	1.4167±0.107 <sup>b</sup>	1.516±0.116 <sup>bc</sup>	1.558±0.118 <sup>c</sup>
	100	1.630±0.125 <sup>a</sup>	1.863±0.143 <sup>b</sup>	2.00±0.152 <sup>bc</sup>	2.068±0.158 <sup>c</sup>
	140	1.900±0.143 <sup>a</sup>	2.190±0.170 <sup>b</sup>	2.395±0.183 <sup>bc</sup>	2.450±0.188 <sup>c</sup>
Carotenoids (mg/g FW)	60	1.086±0.080 <sup>a</sup>	1.213±0.092 <sup>b</sup>	1.305±0.098 <sup>bc</sup>	1.3333±0.100 <sup>c</sup>
	100	1.390±0.107 <sup>a</sup>	1.580±0.120 <sup>b</sup>	1.748±0.132 <sup>bc</sup>	1.791±0.136 <sup>c</sup>
	140	1.655±0.125 <sup>a</sup>	1.860±0.143 <sup>b</sup>	1.935±0.147 <sup>b</sup>	1.978±0.149 <sup>b</sup>
Anthocyanin (mg/g FW)	60	0.930±0.067 <sup>a</sup>	1.030±0.076 <sup>b</sup>	1.100±0.085 <sup>bc</sup>	1.130±0.085 <sup>c</sup>
	100	1.210±0.094 <sup>a</sup>	1.370±0.102 <sup>b</sup>	1.501±0.114 <sup>c</sup>	1.541±0.118 <sup>c</sup>
	140	1.498±0.114 <sup>a</sup>	1.748±0.132 <sup>b</sup>	1.900±0.143 <sup>bc</sup>	1.930±0.143 <sup>c</sup>
Xynthophyll (mg/g FW)	60	0.693±0.053 <sup>a</sup>	0.760±0.058 <sup>ab</sup>	0.821±0.060 <sup>bc</sup>	0.840±0.067 <sup>c</sup>
	100	0.921±0.069 <sup>a</sup>	1.031±0.078 <sup>b</sup>	1.120±0.085 <sup>bc</sup>	1.140±0.085 <sup>c</sup>
	140	1.296±0.098 <sup>a</sup>	1.466±0.112 <sup>b</sup>	1.600±0.120 <sup>bc</sup>	1.631±0.123 <sup>c</sup>

Expressed values are mean±SD of six replicates in each group. Values that are not sharing a common superscript (a, b, c) differ significantly at  $P \leq 0.05$ . IBA: Indole-3-butyric acid, TDM: Triadimefon, PCZ: Propiconazole, DAS: Days after sowing, SD: Standard deviation

were observed in growth regulators treated *Catharanthus* plants (Jaleel *et al.*, 2006). Uniconazole and paclobutrazol treatment induced higher level of carotenoid content in wheat seedlings (Fletcher and Hofstra, 1988; Berova *et al.*, 2003). Increased level of uniconazole treatment and thus increased zeatin might be responsible for the increased synthesis of carotenoid in the plants (Grossmann *et al.*, 1994). Triazole increase the active oxygen species, thus delaying the senescence of wheat and prolonging the duration of flag leaf photosynthesis *Triticum aestivum* and *Didymella exitialis* (Bertelsen *et al.*, 2001; Cromey *et al.*, 2004). IBA increased the vegetative growth and pigments concentration in Maize (Kaya *et al.*, 2006). Similar result was also observed in grapevine cuttings (Sukhwant *et al.*, 2002). Growth hormones have been shown to play an important role in regulating the amount and distribution of assimilates in plants (Galston and Davies, 1969). The increased photosynthetic content in leaves increased in IBA treated cutting that these might have altered the synthesis and translocation of assimilates (Kaur *et al.*, 2002). IBA increased photosynthetic pigments in garlic and these results are agreement with Bideshki and Arvin (2013), onion (Amin *et al.*, 2007) and pea (El-Shraiy and Hegazi, 2009). The IBA increasing photosynthetic pigments increased growth parameters and bulb yield in garlic and increasing in growth parameters and yield was reported at onion (Amin *et al.*, 2006; 2007).

PCZ, TDM, and IBA treatments increased the anthocyanin content of ashwagandha leaves. Among the treatments, PCZ and TDM increased larger extent when compared to IBA [Table 1]. TDM treatment increased the chlorophyll and anthocyanin content in radish (Lichtenthaler, 1979; Sridharan *et al.*, 2006b). Triazole greatly increases anthocyanin accumulation in carrot through tissue culture (Husen and Dougall, 1992). Tetraconazole increased the anthocyanin content in maize (Angela *et al.*, 1997). Similar

results were observed in hexaconazole and paclobutrazol treated in carrot (Gopi *et al.*, 2007). Paclobutrazol treated plant leaves were dark green due to high chlorophyll content in *D. caryophyllus* (Sebastian *et al.*, 2002) and in potato (Tekalign and Hammes, 2004; Tekalign *et al.*, 2005). Triazole induced a transient raise in abscisic acid content in bean (Asare-Boamah and Fletcher, 1986). The increase in ABA content induced by triazole might be the cause for the increased anthocyanin content of ashwagandha.

Triazole and IBA treated plants showed increased xanthophyll content at all stages of growth. Among those, PCZ increased it to a higher level than TDM when compared to IBA treated *W. somnifera* leaves [Table 1]. Triazole compounds increased photosynthetic pigments and induced a variety of morphological, physiological, biochemical responses in *A. campanulatus* (Gopi *et al.*, 2005). Triazole treatment increased the xanthophyll content to a higher level in cucumber (Feng *et al.*, 2003) and carrot (Gopi *et al.*, 2007). TDM treatment increased the chlorophyll, carotenoid and xanthophyll content in the leaves of barley (Forster, 1978). The unsaturated C<sub>40</sub> hydrocarbons not only give color to fruits and flowers but also have multiple functions in photosynthesis. They participate in light harvesting in photosynthetic membranes and protect the photosynthetic apparatus from excessive light energy by quenching triplet chlorophylls and singlet oxygen (Siefermann-Harms, 1987). PBZ increased the chlorophyll content to a larger extent when compared to the control plants in *rosea* and *alba* varieties of *C. roseus* (Jaleel *et al.*, 2008).

Auxins are another group of endogenous plant growth substances. The major site of cytokinin biosynthesis in higher plants is the root, and then transported to the aerial portions of the plant through the xylem. These hormones have potent effects on plant physiology and

are intimately involved in the regulation of cell division, apical dominance, chloroplast development, anthocyanin production and maintenance of the source-sink relationship (Hutchinson and Kieber, 2002). Growth hormones have been shown to play an increasingly important role in regulating the amount and distribution of assimilates in plants (Galston and Davies, 1969). IBA might have altered the synthesis and translocation of assimilates. This may be due to alterations in the rate of photosynthesis and outflow of assimilates from leaves. So, triazole and IBA treated ashwagandha plants increased the chlorophyll, anthocyanin, and xanthophylls content.

### Influence of IBA, TDM, and PCZ on the Biochemical Contents of Ashwagandha

In ashwagandha plants, the starch contents were increased with all stages of growth in the control and treated plants. The starch contents increased in all the plant parts by the triazole and IBA treatments. Among the treatments, PCZ and TDM caused a profound effect in the increase of starch content when compared to IBA. Among the plant parts, roots had higher level accumulation of biochemical contents when compared to leaves and stem in PCZ treatments [Table 2]. Triazole exhibited a higher level of accumulation of starch content in root at all stages of growth when compared to IBA treated plants. Triazole compounds are known to alter the carbohydrate status in various plants like sweet orange (Vu and Yelenosky, 1992) and potato (Kapur *et al.*, 1993). TDM and paclobutrazol increased the nonstructural carbohydrates in mature tuber of *Dioscorea* (Jaleel *et al.*, 2007) and *S. rotundifolius* (Kishorekumar *et al.*, 2007). Triazole compounds are known for their altering effect of carbohydrate metabolism in many plants (Fletcher *et al.*, 2000). The increased starch content in triazole treated plants may be due to decrease in starch hydrolysis as reported in treated Bean (Steffens *et al.*, 1983; Upadhyaya *et al.*, 1986). Starch is the predominant carbohydrate reserve in many plants and is found in both

the photosynthetic tissues (Slattery *et al.*, 2000). Triazole inhibited the gibberellins biosynthesis while increasing the cytokinins. The increased cytokinin content induced by TDM and PCZ might have increased the starch content in roots of ashwagandha. IBA and NAA concentration increased the major reserve compounds such as starch and soluble carbohydrates content in *Aechmea blanchetiana* (Chu *et al.*, 2010). IBA treatment significantly increased the total carbohydrate content in *P. sativum* (El-Shraiy and Hegazi, 2009). Similar results have observed in *Zea mays* (Amin *et al.*, 2006). IBA caused a gradual increase in the total carbohydrate in *Thunbergia grandiflora* and *Beaumontia grandiflora* (Hussein, 2008, 2003).

In all the treatments significantly increased the protein content in the leaves and in roots of *W. somnifera*. Among the treatments, PCZ and TDM caused higher level of accumulation of protein content of leaf and root at all stages of growth when compared to IBA. Among the plant parts storage roots had higher content when compared to leaves and stem [Figure 1]. The biochemical content increased, while protein, amino acid, were increased due to TDM treatment and lead to early sprouting in *Dioscorea rotundata* (Jaleel *et al.*, 2007). TDM treatment increased the protein content in cucumber seedling (Feng *et al.*, 2003). Difenconazole caused higher level of protein accumulation in all parts of *Mentha piperita* (Kavina *et al.*, 2011). TDM and PCZ treatment increased the protein content in *R. sativus* (Sridharan *et al.*, 2006a). Ketaconazole treatment increased the protein content in cowpea (Gopi *et al.*, 1999; Somasundaram *et al.*, 2005). TDM treatment increased the protein content in *C. roseus* (Jaleel *et al.*, 2008). Esashi and Leopold (1968) correlated the increased soluble protein synthesis with the enhanced cytokinin content of the tubers of *Begonia evansiana*. The increased cytokinin content induced by TDM and PCZ treatments might have increased the protein content in the leaf, stem and root of *W. somnifera*. The protein content was significantly higher in both IBA and NAA treated

Table 2: Influence of IBA, TDM and PCZ on the starch content of ashwagandha

Plant parts	Growth stages (DAS)	Control	IBA	TDM	PCZ
Starch (mg/g fr. wt.)					
Leaf	60	13.661±1.039 <sup>a</sup>	14.731±1.124 <sup>ab</sup>	15.320±1.167 <sup>c</sup>	15.430±1.175 <sup>c</sup>
	100	15.006±1.148 <sup>a</sup>	16.495±1.262 <sup>ab</sup>	17.190±1.316 <sup>c</sup>	17.310±1.325 <sup>c</sup>
	140	16.370±1.247 <sup>a</sup>	18.023±1.370 <sup>ab</sup>	19.003±1.447 <sup>c</sup>	19.123±1.455 <sup>c</sup>
Stem	60	12.038±0.918 <sup>a</sup>	12.931±0.981 <sup>ab</sup>	13.623±1.037 <sup>c</sup>	13.745±1.046 <sup>c</sup>
	100	13.711±1.050 <sup>a</sup>	14.816±1.130 <sup>ab</sup>	15.813±1.211 <sup>c</sup>	15.911±1.220 <sup>c</sup>
	140	15.103±1.151 <sup>a</sup>	16.803±1.276 <sup>b</sup>	17.713±1.348 <sup>c</sup>	17.831±1.359 <sup>c</sup>
Root	60	16.223±1.236 <sup>a</sup>	17.673±1.344 <sup>ab</sup>	18.763±1.429 <sup>c</sup>	18.820±1.433 <sup>c</sup>
	100	18.033±1.381 <sup>a</sup>	19.853±1.518 <sup>ab</sup>	21.283±1.630 <sup>c</sup>	21.511±1.645 <sup>c</sup>
	140	20.013±1.523 <sup>a</sup>	22.376±1.704 <sup>b</sup>	24.336±1.852 <sup>c</sup>	24.513±1.868 <sup>c</sup>

Expressed values are mean±SD of six replicates in each group. Values that are not sharing a common superscript (a, b, c) differ significantly at  $P \leq 0.05$ . IBA: Indole-3-butyric acid, TDM: Triadimefon, PCZ: Propiconazole, DAS: Days after sowing, SD: Standard deviation

*Aechmea blanchetiana* (Chu *et al.*, 2010). IBA treatment also increases the proteins in maize plants (El-Wahed *et al.*, 2006), *Dalbergia sissoo* (Husen, 2008) and grapevine (Kaur *et al.*, 2002).

In ashwagandha plants, the amino acid content increased with the age in control and treated plants in all growth stages. In root and leaves, a maximum increase was noted on 140 DAS in triazole when compared to IBA treatments. Among the organs, leaf tissue accumulated higher level of amino acids than stem and root tissue of the *W.somnifera* with

triazole treatments [Figure 2]. Similarly, TDM treatment increased the amino acid content in *Catharanthus* in all parts at all growth stage (Jaleel *et al.*, 2007). Hexaconazole and paclobutrazol treatments increase in amino acid content was reported in carrot plants by Gopi *et al.* (2007) and TDM and PCZ treated in *R. sativus* (Sridharan *et al.*, 2006a). Uniconazole and ABA treatment were found to increase the concentration of amino acid, protein and proline in *Phaseolus vulgaris* (Mackay *et al.*, 1990). Similarly, Penconazole induced a moderate increase in amino acid in higher plants (Radice and Pesci, 1991). TDM increased




Figure 1: (a-c) Influence of indole-3-butyric acid, triadimefon and propiconazole on the protein content of ashwagandha




Figure 2: (a-c) Influence of indole-3-butyric acid, triadimefon, and propiconazole on the amino acid content of ashwagandha

the amino acid content in soybean (Panneerselvam *et al.*, 1998) and radish (Muthukumarasamy *et al.*, 2000). IBA treated pea plant significantly increased the amino acid content (El-Shraiy and Hegazi, 2009) and garlic (Bideshki and Arvin, 2013).

## CONCLUSION

All the treatments significantly increased photosynthetic pigments and biochemical constituents of leaf, stem and root in *W. somnifera*. Among the treatments, PCZ and TDM caused a pronounced effects on increased the photosynthetic pigments (total chlorophyll, carotenoids anthocyanin, and xanthophylls) and biochemical constituents (starch, protein, and amino acids) content when compared to IBA. Triazole treated plants typically appear dark greener and this has been correlated with higher level of chlorophyll content in ashwagandha leaves. Triazole compounds Viz., PCZ and TDM act as growth regulators or retardants to enhanced the biochemical constituents such as., starch, protein and amino acids contents, in starch more in roots, which is helpful to satisfy the needs of improve the root growth and enhance the biochemicals of medicinally important root crops of ashwagandha.

## REFERENCES

- Agarwal R, Diwanay S, Patki P, Patwardhan B. Studies on immunomodulatory activity of *Withania somnifera* (Ashwagandha) extracts in experimental immune inflammation. *J Ethnopharmacol* 1999;67:27-35.
- Amin AA, Rashad M, Sh EL, Gharib FA. Physiological response of maize plants (*Zea mays* L.) to foliar application of morphactin CF and indole-3-butyric acid. *J Biol Sci* 2006;6:547-54.
- Amin AA, Sh EL, Rashad M, El-Abagy HM. Physiological effect of indole-3- butyric acid and salicylic acid on growth, yield and chemical constituents of Onion plants. *J Appl Sci Res* 2007;3:1554-63.
- Angela R, Gandolfina F, Franco G, Chiara T. Effects of a Triazolic fungicide on maize plant metabolism; modifications of transcript abundance in resistance-related pathways. *Plant Sci* 1997;130:51-62.
- Arnon DI. Copper enzymes in isolated chloroplasts, polyphenol oxidase in *Beta vulgaris* L. *Plant Physiol* 1947;24:1-15.
- Asare-Boamah NK, Fletcher RA. Protection of bean seedlings against heat and chilling injury by triadimefon. *Physiol Plant* 1986;67:353-8.
- Beggs CJ, Wellmann E. Analysis of light controlled anthocyanin synthesis in coleoptiles of *Zea mays* L. The role of UV-B blue red and far red light. *Photochem Photobiol* 1985;41:401-6.
- Berova M, Zlatev Z, Stoeva N. Effect of paclobutrazol on wheat seedlings under low temperature stress. *Bulg J Plant Physiol* 2003;28:75-84.
- Bertelsen JR, De-Neergaard E, Petersen VS. Fungicidal effects of azoxystrobin and epoxiconazole on Phyllosphere fungi, senescence and yield of winter wheat. *Plant Pathol* 2001;50:190-205.
- Bhattacharya SK, Satyan KS, Ghosal KS. Antioxidant activity of glycowithanolides from *Withania somnifera*. *Indian J Exp Biol* 1997;35:236-9.
- Bideshki A, Arvin MJ. Interactive effects of methyl jasmonate (MJ) and indole-3-butyric acid (IBA) on growth and biochemical parameters, bulb and allicin yield of Garlic (*Allium sativum* L.) under drought stress in Iran. *Int J Agric Res Rev* 2013;3:349-60.
- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248-54.
- Chu EP, Tavares AR, Kanashiro S, Giampaoli P, Yokota ES. Effects of auxins on soluble carbohydrates, starch and soluble protein content in *Aechmea blanchetiana* (Bromeliaceae) cultured *in vitro*. *Sci Hortic* 2010;125:451-5.
- Clegg KM. The application of the anthrone reagent to the estimation of starch in cereals. *J Sci Food Agric* 1956;7:40-4.
- Cromey MG, Butler RC, Mace MA, Cole AL. Effects of the fungicides azoxystrobin and tebuconazole on *Didymella exitialis*, leaf senescence and grain yield in wheat. *Crop Prot* 2004;23:1019-30.
- El-Shraiy AM, Hegazi AM. Effect of acetylsalicylic acid, indole-3-butyric acid and gibberellic acid on plant growth and yield of Pea (*Pisum sativum* L.). *Aust J Basic Appl Sci* 2009;3:3514-23.
- El-Wahed A, Amin AA, El-Sh RM. Physiological effect of some bioregulators on vegetative growth, yield and chemical constituents of yellow Maize plants. *World J Agric Sci* 2006;2:149-55.
- Esashi Y, Leopold AC. Regulation of tuber development in *Begonia eransiana* by cytokinin. In: Wightman F, Setterfield G, editors. *Biochemistry and Physiology of Plant Growth Substances*. Ottawa: The Runge; 1968. p. 923-42.
- Farooqi AA, Shreeramu BS. *Cultivation of Medicinal and Aromatic Crops*. Hyderabad, India: University Press (India) Ltd.; 2004.
- Feng Z, Guo A, Feng Z. Amelioration of chilling stress by triadimefon in cucumber seedlings. *Plant Growth Regul* 2003;39:277-83.
- Fletcher RA, Gilley A, Davis TD, Sankhla N. Triazoles as plant growth regulators and stress protectants. *Hortic Rev* 2000;24:55-138.
- Fletcher RA, Hofstra G. Triazole as potential plant protectants. In: Berg D, Plempel M, editors. *Sterol Biosynthesis Inhibitors*. Cambridge, England: Ellis Horwood Ltd.; 1988. p. 321-31.

- Forster H. Mechanism of action and side effects of triadimefon and triadimenol in barley plants. 3<sup>rd</sup> International Conference Plant Pathology. Munchon; 1978. p. 365.
- Galal AA, Ibrahiem IA, Salem JM. Some morphological and physiological abnormalities in growth and development of banana triadimefon-treated cultivars. *Genet Plant Physiol* 2011;1:91-102.
- Galston AW, Davies PJ. Hormonal regulation in higher plant. *Science* 1969;163:1288-97.
- Gopi R, Jaleel CA, Sairam R, Lakshmanan GM, Gomathinayagam M, Panneerselvam R. Differential effects of hexaconazole and paclobutrazol on biomass, electrolyte leakage, lipid peroxidation and antioxidant potential of *Daucus carota* L. *Colloids Surf B Biointerfaces* 2007;60:180-6.
- Gopi R, Sridharan R, Somasundaram R, Alagu GM, Panneerselvam R. Growth and photosynthetic characteristics induced by triazoles in *Amorphophallus campanulatus* Blume. *Genet Appl Plant Physiol* 2005;31:171-80.
- Gopi R, Sujatha BM, Rajan SN, Karikalan L, Panneerselvam R. Effect of triadimefon in the sodium chloride stressed cowpea (*Vigna unguiculata* L.) seedlings. *Indian J Agric Sci* 1999;69:743-5.
- Grossmann K, Kwiatkowski K, Hauser C, Siefert F. Influence of the triazole growth retardant BAS 111W on phytohormone levels in senescing intact pods of oilseed rape. *Plant Growth Regul* 1994;14:115-8.
- Gupta A, Mahdi AA, Shukla KK, Ahmad MK, Bansal N, Sankhwar P, Sankhwar SN. Efficacy of *Withania somnifera* on seminal plasma metabolites of infertile males: A proton NMR study at 800 MHz. *J Ethnopharm* 2013;149:208-14.
- Husen A, Dougall DK. The effect of growth retardants on anthocyanin production in carrot cell suspension cultures. *Plants Cell Rep* 1992;11:304-9.
- Husen A. Clonal propagation of *Dalbergia sissoo* Roxb. and associated metabolic changes during adventitious root primordium development. *N Forest* 2008;36:13-27.
- Hussain A. Economic aspects of exploitation medicinal plants. In: Akerele O, Heywood V, Synge H, editors. *The Conservation of Medicinal Plants*. Cambridge, New York: Cambridge University Press; 1991. p. 124-38.
- Hussein MM. Effect of planting dates and indole-3-butyric acid on rooting of *Beaumontia grandiflora* Wallich. cuttings and consequent plant growth. *Arab Univ J Agric Sci Ain Shams Univ Cairo* 2003;11:765-87.
- Hussein MM. Studies on the rooting and the consequent plant growth on the stem cutting of *Thunbergia grandiflora* (Roxb. ex Rottl.) Roxb. 2-Effect of indole-3-butyric acid. *World J Agric Sci* 2008;4:811-7.
- Hutchinson CE, Kieber JJ. Cytokinin signaling in *Arabidopsis*. *Plant Cell* 2002;14:S47-59.
- Indian Pharmacopocia, Ashwagandha. In: IP 1985, Appendix 3.3:10:69, Ministry of Health and Family Welfare, Government of India. New Delhi, India: Controller of Publications.
- Jakhar ML, Singh K, Kakralya BL. Prospective strategy for biodiversity conservation and development in medicinal and aromatic plants. In: Sharma HS, Khan TI, editors. *Environmental Conservation, Depleting Resources and Sustainable Development*. Jaipur, India: Aavishkar Publishers; 2003. p. 145-156.
- Jaleel CA, Gopi R, Manivannan P, Kishorekumar A, Gomathinayagam M, Panneerselvam R. Changes in biochemical constituents and induction of early sprouting by triadimefon treatment in white yam (*Dioscorea rotundata* Poir.) tubers during storage. *J Zhejiang Univ Sci B* 2007;8:283-8.
- Jaleel CA, Gopi R, Manivannan P, Kishorekumar A, Sankar B, Panneerselvam R. Paclobutrazol influences on vegetative growth and floral characteristics of *Catharanthus roseus* (L.) G. Don. *Indian J Appl Pure Biol* 2006;21:369-72.
- Jaleel CA, Gopi R, Panneerselvam R. Growth and photosynthetic pigments responses of two varieties of *Catharanthus roseus* to triadimefon treatment. *C R Biol* 2008;331:272-7.
- Kakimoto T. Perception and signal transduction of cytokinin. *Ann Rev Plant Biol* 2003;54:605-27.
- Kapoor LD. *Handbook of Ayurvedic Medicinal Plants*. London, UK: CRC Press; 2001. p. 337-8.
- Kapur KK, Narula SL, Upadhyaya MD. Modification of physiological and biochemical responses in potato with CCC and triazole, *Proceedings of the National Symposium Plant Physiological*. Lucknow, India: NBRL; 1993. p. 361-9.
- Kaur SS, Cheema B, Chhabra R, Talwar KK. Chemical induction of physiological changes during adventitious root formation and bud break in grapevine cuttings. *Plant Growth Regul* 2002;37:63-8.
- Kavina J, Gopi R, Panneerselvam R. Traditional and nontraditional plant growth regulators alter the growth and photosynthetic pigments in *Mentha piperita* Linn. *Int J Environ Sci* 2011;1:124.
- Kaya C, Levent Tuna A, Alves AA. Gibberellic acid improves water deficit tolerance in maize plants. *Acta Physiol Plant* 2006;28:331-7.
- Kishorekumar A, Jaleel CA, Manivannan P, Sankar B, Sridharan R, Panneerselvam R. Comparative effects of different triazole compounds on growth, photosynthetic pigments and carbohydrate metabolism of *Solenostemon rotundifolius*. *Colloids Surf B Biointerfaces* 2007;60:207-12.
- Kokate CK, Purohit AP, Gokhale SB. *Pharmacognosy*. 33<sup>rd</sup> ed. Pune: Nirali Prakashan; 2005. p. 518-20.
- Lichtenthaler HK. Effect of biocides on the development of the photosynthetic apparatus of radish seedlings grown under strong and weak high concentrations. *Z Naturforsch* 1979;34:939-40.
- Ludwig-Muller J. Indole-3-butyric acid in plant growth and development. *Plant Growth Regul* 2000;32:219-30.



- Mackay CE, Hall JC, Hofstra G, Fletcher RA. Uniconazole induced changes in abscisic acid, total amino acids and proline in *Phaseolus vulgaris*. Pestic Biochem Physiol 1990;37:74-82.
- Moore S, Stein WH. Photometric method for use in the chromatography of amino acids. J Biol Chem 1948;176:367-88.
- Muthukumarasamy M, Gupta SD, Panneerselvam R. Influence of triadimefon on the metabolism of NaCl stressed radish. Biol Plant 2000;43:67-72.
- Neogy M, Datta JK, Mukherji S, Roy AK. Effect of aluminium on pigment content, hill activity and seed yield in mung bean. Indian J Plant Physiol 2001;6:381-5.
- Okigbo RA, Anuagasi CL, Amadi CL. Advances in selected medicinal and aromatic plants indigenous to Africa. J Med Plant Res 2009;3:86-95.
- Pacholczak A. The effect of shading of stock plants on rhizogenesis in stem cuttings of *Berberis thunbergii* 'Red Rocket'. Physiol Plant 2006;28:567-75.
- Palash M, Mitali G, Kumar MT, Prasad DA. Pharmacognostic and free-radical scavenging activity in the different part of ashwagandha *Withania somnifera* (L.Dunal). Int J Drug Dev Res 2010;2:830-43.
- Panneerselvam R, Muthukumarasamy M, Rajan SN. Amelioration of NaCl stress by triadimefon in soybean seedlings. Biol Plant 1998;41:133-7.
- Puri HS. Rasayana - Ayurvedic Herbs for Longevity and Rejuvenation. London: Taylor and Francis; 2003.
- Radice M, Pesci P. Effect of triazole fungicides on the membrane permeability and on FC-induced H<sup>+</sup>- extrusion in higher plants. Plant Sci 1991;74:81-8.
- Rasool M, Varalakshmi P. Immunomodulatory role of *Withania somnifera* root powder on experimental induced inflammation: *In vivo* and *in vitro* study. Vascul Pharmacol 2006;44:406-10.
- Sankari S, Gopi R, Gomathinayagam M, Sridharan R, Somasundaram R, Panneerselvam R. Responses of triazoles on growth and antioxidant levels in white radish. Indian J Appl Pure Biol 2006;21:77-80.
- Sarkar S, Perras MR, Falk DE, Zhang R, Pharis RP, Fletcher RA. Relationship between gibberellins, height and stress tolerance in barley (*Hordeum vulgare* L.) seedlings. Plant Growth Regul 2004;42:125-35.
- Seartezzini P, Speroni S. Review on a some plants of India traditional medicine with antioxidant activity. J Ethnopharm 2000;71:23-47.
- Sebastian B, Alberto G, Emillio AC, Jose AF, Juan AF. Growth development and color response of potted *Dianthus caryophyllus* to paclobutrazol treatment. Sci Hortic 2002;1767:1-7.
- Sieffermann-Harms D. The light-harvesting and protective functions of carotenoids in photosynthetic membranes. Physiol Plant 1987;69:561-8.
- Singh G, Sharma PK, Dudhe R, Singh S. Biological activities of *Withania somnifera*. Ann Biol Res 2010;1:56-63.
- Singh K, Tyagi SK. Medicinal Plants-Applied Biology of Domestication and Export. Jaipur, India: Aavishkar Publishers; 2004.
- Slattery CJ, Kavakli IH, Okita TW. Engineering starch for increased quantity and quality. Trends Plant Sci 2000;5:291-8.
- Somasundaram R, Manivannan P, Kishorekumar A, Gopi R, Alagu Lakshmanan GM, Panneerselvam R. Growth and antioxidant potential induced by ketoconazole in sodium chloride stressed *Vigna unguiculata* (L.) Walp. J Theor Exp Biol 2005;2:77-82.
- Sridharan R, Kishorekumar A, Somasundaram R, Manivannan P, Alagu Lakshmanan GM, Gomathinayagam M, Panneerselvam R. Effect of triazole on growth and chlorophyll pigment in radish. Indian J Environ Ecoplan 2006b;12:57-62.
- Sridharan R, Kishorekumar A, Somasundaram R, Manivannan P, Gomathinayagam M, Panneerselvam P. Morphological and biochemical responses of radish using triazole compounds. Geobios 2006a;33:253-6.
- Steffens GL, Wang SY, Steffens CL, Brennan T. Influence of paclobutrazol (PP 333) on apple seedling growth and physiology. Proc Plant Growth Regul Soc Am 1983;10:195-205.
- Still JR, Pill WG. Growth and stress tolerance of tomato seedlings (*Lycopersicon esculentum* Mill.) in response to seed treatment with paclobutrazol. Hortic Sci 2004;79:197-203.
- Sukhwant K, Cheema SS, Chhabra BR, Talwar KK. Chemical induction of physiological changes during adventitious root formation and bud break in grapevine cuttings. Plant Growth Regul 2002;37:63-8.
- Sunitha S, Perras MR, Falk DE, Ruichuon Zhang R, Pharis P, Fletcher RA. Relationship between gibberellins, height and stress tolerance on barley seedlings. Plant Growth Regul 2004;42:125-35.
- Tekalign T, Hammes PS. Response of potato grown under non-inductive condition to paclobutrazol. Shoot growth, chlorophyll content, net photosynthesis, assimilate partitioning, tuber yield, quality and dormancy. Plant Growth Regul 2004;43:227-36.
- Tekalign T, Hammes S, Robbertse J. Paclobutrazol induced leaf stem and root anatomical modifications in potato. Hortic Sci 2005;40:1343-6.
- Upadhyaya A, Davis TD, Sankhla N. Some biochemical changes associated with paclobutrazole induced adventitious root formation on bean hypocotyls cuttings. Ann Bot 1986;57:309-15.
- Vu JC, Yelenosky G. Growth and photosynthesis of sweet orange plants treated with paclobutrazol. J Plant Growth Regul 1992;11:85-9.