

Regular Article

## A comparative kinetic study on $\beta$ -amylase and its antioxidant property in germinated and non germinated seeds of *Glycine max. L*

Chandrika C\*, Vijayashree C, Granthali, Rajath S, Nagananda G.S and S. Sundara Rajan

Department of Biochemistry, Genohelix Biolabs- A Division of CASB, Jain University, 127/2, Bull Temple Road, Chamrajpet, Bangalore-560019, Karnataka, India

\*Corresponding author Email: [chandrikac237@gmail.com](mailto:chandrikac237@gmail.com)

Enzyme activity plays a key role in each stage of plant development starting from the initial seed germination responses. A comparative analysis of beta amylase activity was performed, isolated from germinated and non germinated *Glycine max* seeds, which was partially purified by dialysis. The optimum pH and temperature for germinated seeds was found to be 4.5 and 37<sup>o</sup> C and that for non germinated seeds were 5.5 and 75<sup>o</sup> C respectively. MnCl<sub>2</sub> and CoCl<sub>2</sub> exhibited marked activating effect on the enzyme, while HgCl<sub>2</sub> was a potent inhibitor for both seeds. The K<sub>m</sub> and V<sub>max</sub> value for  $\beta$ -amylase with soluble starch as substrate was found to be 3.03mg/ml and 6.6micromol/min/ml for germinated seed and for non germinated seeds it was found to be 5mg/m and 10micromole/min/ml respectively. The molecular weight of partially purified enzyme was 57 $\pm$ 1kDa on SDS PAGE with 1.918 fold purification and 27.98% yield for germinated seeds and for non germinated seeds the purification fold was 1.507 with 26.65% yield. The enzyme showed highest amount of total antioxidant activity of 304 $\mu$ g/ml in partially purified germinated seed. For ABTS free radical scavenging activity, partially purified enzyme from germinated seed showed the highest of 76.09% activity with an IC<sub>50</sub> value of 58.20 $\mu$ g/ml. Germinated seeds have highest enzymatic activity which can be considered for several industrial purposes.

**Key words:** *Glycine max*,  $\beta$ -amylase, antioxidants, enzyme activity.

Most species of the legume family (*Fabaceae*) have non-endospermic seeds. Dry legumes are good sources of protein, energy and other nutrients in developing countries. However, their use is limited because of high dietary bulk; presence of anti nutritional factors, mainly phytic acid in most of the legumes; and low protein and carbohydrate digestibility (Sumathi *et al.*, 1995; Komal and Darshan, 2000; Mubarak, 2005; Negi *et al.*, 2001). The cotyledons serve as sole food

storage organs. During embryo development the cotyledons absorb the food reserves from the endosperm completely. In most seeds, the food is stored in the cotyledons, or seed leaves, as starch. Although starch is a storage form of glucose, it cannot pass through living membranes due to the large molecular structure, so before it can be used for energy it needs to be broken down to its constituent sugars by enzymes. Enzymes are the biological catalysts that allow the chemical

reactions necessary for metabolism to occur. During germination it secretes the amylase enzyme that breaks down endosperm starch into sugars to nourish the growing seedling.

Amylase is classified into alpha, beta and gamma. Both  $\alpha$ -amylase and  $\beta$ -amylase are present in seeds.  $\beta$ -amylase is present in an inactive form prior to germination, whereas  $\alpha$ -amylase and proteases appear once germination has begun. Germination was shown to increase monosaccharide and decreased disaccharide contents of legumes due to  $\alpha$ -amylase (Akinlosottu and Akinyele, 1991). Amylase are among the most important enzymes used for several biotechnological applications particularly employed in starch processing industries for the hydrolysis of polysaccharides such as starch into simpler sugar constituents (Mubarak, 2005). It is also found useful in a wide variety of industrial applications such as production of ethanol and high-fructose corn syrup, detergents, desizing of textiles, modified starches, hydrolysis of oil-field drilling fluids and paper recycling (Sumathi et al., 1995).

Antioxidants are the substances able to prevent or inhibit oxidation processes in human body as well as in food products. The natural antioxidants are a stable part of nutrition as they occur in almost all edible plant products (Aleksandra and Tomasz, 2007). An antioxidant may play a role in antioxidation as a free radical scavenger, reducing agent, chelator and/or singlet oxygen scavenger. Studies have demonstrated that plant phenolics are a major source of natural antioxidants (Emma and Nedyalka, 2003). They can be distributed in fruits, seeds, leaves, vegetables, barks, roots, and flowers of plants (Wanasundara et al., 1997, Wang & Lin, 2000).

So, the present investigation was started with the perspective to analyze the activity of enzyme isolated from germinated and non germinated seeds and to check its

antioxidant capacity with the prospective of increasing its importance at industrial level.

### Materials and Methods

Soya Bean seeds (*Glycine max*), a common variety in India, was purchased from a local market in Bangalore and stored in dry place at room temperature for experimental set up.

### Preparation and extraction of crude enzyme

Two sets of 25g Glycine max seeds were taken and washed thoroughly under running tap water and surface sterilized by 2% savlon, and tween20 for 20 minutes. It was then rinsed with 70% ethanol for 30 seconds and 0.05%  $\text{HgCl}_2$  for 6minutes. The seeds were then rinsed with distilled water to remove the traces of  $\text{HgCl}_2$ . One set of sterilized seeds were soaked in 25mM sodium acetate buffer pH 5.5 overnight at room temperature and the other set of seeds were allowed to germinate by covering them in muslin cloth for 24 hrs. The germinated and non germinated seeds were homogenized in sodium acetate buffer using blender. The homogenate was filtered through muslin cloth. The filtrate was collected and clarified further by centrifugation at 5000rpm for 20minutes at 4°C. The supernatant (crude precipitate) was collected and stored at 4°C for further enzyme assays.

### Enzyme and Protein assay

The enzyme activity for germinated and non germinated seeds were determined by Bernfeld method (Bernfeld, 1955). The crude enzyme extract was diluted in the ratio of 1:30, using sodium acetate buffer. 0.1ml of and the diluted sample was incubated in 0.1ml of 1% soluble starch at room temperature for 5minutes. To this, 1.0ml of DNS was added and kept in boiling water bath for 20minutes. The final volume was made up to 10ml using distilled water and absorbance was read at 540 nm using

distilled water as blank. A standard curve using 1mg/ml of maltose was prepared. One unit (U/ml) of amylase activity was defined as the amount of enzyme that releases 1 $\mu$ g of maltose per minute under standard assay conditions. The amount of protein in crude enzyme extract was determined by Lowry's method (Lowry *et al.*, 1951) for both germinated and non germinated extract using bovine serum albumin as standard.

#### **Partial purification of amylase**

The crude extract of germinated and non-germinated seeds (50ml) was precipitated using 100% saturated ammonium sulphate solution. The precipitation was carried out at 4°C by constant stirring and centrifuged at 10,000rpm (9615g) for 15minutes. The precipitate was dissolved in sodium acetate buffer pH 5.5 and subjected to dialysis. Specific activity for the three fraction was estimated. The dialyzed sample was used for further studies.

#### **Starch hydrolysis test for amylase (Shelby, 1993)**

The hydrolysis of starch by partially purified amylase was studied by preparing starch agar plates. The plates were prepared by adding 1ml of 1% soluble starch mixed with 1% liquid agar and poured on a clean glass slide and allowed to solidify. After solidification, a well was created in the plate using gel punch and 40 $\mu$ l of partially purified germinated and non germinated enzyme was added. Then slides were maintained in a moist condition and incubated overnight at room temperature. After overnight incubation 1ml of iodine solution was added.

#### **TLC analysis (Bilal and Figen, 2005)**

The hydrolysis of soluble starch with dialyzed sample of both germinated and non germinated seeds were determined by thin-layer chromatography on TLC plate. 0.5ml of partially purified enzyme was incubated with

0.5ml of 2% starch for 5 minutes. The sugars released after enzymatic hydrolysis of amylase were separated on TLC using glucose, maltose and starch as standard with the solvent system of butanol: acetic acid: water (4:1:5). Spots on the TLC plate were detected using DPA (diphenylamine) spray after drying at 70°C.

#### **Determination of molecular weight of $\beta$ -amylase (Laemmli, 1970)**

The molecular weight of partially purified  $\beta$ -amylase was analyzed by SDS-PAGE (12%) from Laemmli's method. The bands were visualized by coomassie blue staining in order to determine the molecular weight of  $\beta$ -amylase using standard protein marker of 2-212kDa.

#### **Effect of pH on $\beta$ - amylase activity**

$\beta$ -Amylase activity from the dialyzed samples were determined using 1% soluble starch as substrate at a pH range from 4.5 to 7.8 using 25mM sodium acetate buffer. The optimum pH for enzyme activity was determined using DNS reagent.

#### **Effect of temperature on $\beta$ -amylase activity**

A temperature gradient was employed in order to determine the  $\beta$ -amylase activity. Buffer, substrate solution, and dialyzed samples were incubated at different temperatures between 0°C to 100°C. The optimum temperature was determined using DNS reagent for enzyme activity.

#### **Effect of metal ions on $\beta$ -amylase activity**

The effect of metal ions on the activity of  $\beta$ -amylase were determined by adding known concentrations of metal ions such as CaCl<sub>2</sub>, HgCl<sub>2</sub>, CuCl<sub>2</sub>, SDS, MgCl<sub>2</sub>, EDTA, FeCl<sub>3</sub>, MnCl<sub>2</sub>, CoCl<sub>2</sub>, ZnCl<sub>2</sub>, AlCl<sub>3</sub> and KCl. Each of these metal ions were added to the enzyme substrate reaction and incubated for 5minutes and its activation and inhibition property of each metal ion on enzyme activity was determined using DNS reagent.

### **Effect of Substrate Concentration for determination of Km and Vmax (Lineweaver and Burk, 1984)**

The Michaelis-Menten kinetic constants, Km and Vmax, for partially purified  $\beta$ -amylase was determined by using varying concentrations of soluble starch. Enzyme activity (U/ml) against different concentrations of substrates ( $\mu$ M) were measured under standard assay condition and the kinetic constants, Km and Vmax, were estimated by the method of Lineweaver and Burk plot.

### **Determination of antioxidant capacity**

**Phosphomolybdenum assay** (Prieto *et al.*, 1999)

0.1ml of crude and dialyzed enzyme samples from germinated and non germinated seeds of *Glycine max* was taken and 1ml of total antioxidant reagent (0.6M sulphuric acid, 28mM sodium phosphate and 4mM ammonium molybdate) was added and mixed. Different aliquots (20 $\mu$ g -100 $\mu$ g/ml) of standard ascorbic acid was taken and the volume was made up to 0.1ml with DMSO and 1ml of total antioxidant reagent was added and incubated in a thermal block at 95°C for 90 minutes, cooled to room temperature and the absorbance was measured at 695nm against DMSO as blank.

**ABTS (2, 2-azinobis (3-ethylbenzothiazoline-6-sulfonic acid)) radical cation decolorization assay** (Re *et al.*, 1999)

ABTS radical cation (ABTS<sup>+</sup>) was produced by reacting 7mM ABTS with 2.45mM ammonium persulfate and the mixture was allowed to stand in dark at room temperature for 12-16 hours before use. Enzymes samples of germinated and non germinated, crude and dialyzed, (2mg/ml) were taken at various concentrations (10-50 $\mu$ g/ml) and the volume was adjusted to 500 $\mu$ l with DMSO which also serves as blank. 0.3ml of ABTS solution was added, the final volume was made up to 1ml using ethanol and incubated in dark for 30 minutes at room

temperature. The absorbance was read at 745nm. Radical cat-ion decolonization activity was expressed as the inhibition percentage of cat-ions by the samples and was calculated using the formula,  
% ABTS radical scavenging activity = [(control OD - Sample OD)/ Control OD] \*100.

### **Statistical analysis:**

The experiments were conducted in triplicates. All the values obtained from the mean replicates were averaged. The data were analyzed in relation to the variance and presented as mean+/-standard error (SE). Analysis of variance was conducted by two way ANOVA and all the statistical analysis was performed at 1% significance level using IBM SPSS Statistics (version 20) by IBM.

## **Results**

### **Starch Hydrolysis test**

The crude extract of germinated and non germinated soybean seeds confirmed the presence of amylase after overnight incubation on starch agar plate, a clear zone of starch hydrolysis was observed on addition of iodine solution. The zone of clearance was comparatively high in germinated than in non germinated seed extract.

### **Partial purification of enzyme**

Crude extract was partially purified by ammonium sulphate precipitation and subjected to dialysis and purified to 1.918 fold with a yield of 27.98% for germinated seeds (Table 1) and non germinated seeds were purified to 1.507 fold with yield of 26.65% (Table 2).

### **Determination of molecular weight of $\beta$ -amylase**

The molecular weight of partially purified  $\beta$ -amylase in germinated and non germinated seeds were found to be approximately 57 $\pm$ 1kDa by SDS-PAGE on staining with coomassie brilliant blue (figure 2).

**Table 1:** Summary of purification of  $\beta$ -amylase in germinated soybean seed extract

Step	Volume (ml)	Total Activity ( $\mu\text{mol}/\text{min}$ )	Total Protein (mg)	Specific activity (units/mg of protein)	Purification fold	Yield (%)
Crude enzyme	50	23.3	15.0	1.416	1.0	100
Purified enzyme by chilled ammonium sulphate precipitation method	10	11.748	4.5	2.610	1.843	50.4
Dialysed product	8.0	6.52	2.4	2.716	1.918	27.98

**Table 2:** Summary of purification of  $\beta$ -amylase in non germinated soybean seed extract.

Step	Volume (ml)	Total Activity ( $\mu\text{mol}/\text{min}$ )	Total Protein (mg)	Specific activity (units/mg of protein)	Purification fold	Yield (%)
Crude enzyme	50	21.25	9.3	2.505	1	100
Purified enzyme by chilled ammonium sulphate precipitation method	10	10.98	3.0	3.66	1.46	51.6
Dialysed product	8.0	5.66	1.5	3.776	1.507	26.65

**Table 3:** Percentage of scavenging activity and  $\text{IC}_{50}$  value of different enzyme extracts

Extracts	ABTS% of scavenging	ABTS $\text{IC}_{50}$ ( $\mu\text{g}/\text{ml}$ )	Total antioxidant ( $\mu\text{g}/\text{ml}$ )
Ascorbic acid	52.082 <sup>c</sup>	55.66 <sup>a</sup>	-
Non germinated crude	27.97 <sup>e</sup>	160.74 <sup>e</sup>	20.02 <sup>d</sup>
Non germinated dialyzed	54.00 <sup>b</sup>	76.33 <sup>c</sup>	90.27 <sup>c</sup>
Germinated crude	43.73 <sup>d</sup>	125.46 <sup>d</sup>	239.3 <sup>b</sup>
Germinated dialyzed	76.09 <sup>a</sup>	58.20 <sup>b</sup>	304.0 <sup>a</sup>

**Note:** Mean of triplicates. Mean values with different superscripts (a, b, c, d, e, f, g, h, i, j and k) differ significantly at  $P < 0.01$  by Tukey (HSD) test.

#### TLC analysis of partially purified amylase

The partially purified enzyme was analyzed on TLC plate using maltose and glucose as standard. Both the germinated and non germinated sample fractions correspond with standard maltose which confirmed it to be  $\beta$ -amylase (figure 1).


#### Effect of pH on the $\beta$ -amylase activity

To determine the effect of pH on enzyme activity, different pH ranges- 4.5, 5.5, 6.5, 7.5, 7.8 of sodium acetate buffer was taken and soluble starch was used as substrate.  $\beta$ -amylase activity was found to be maximum at pH 5.5 in germinated seeds and

for non germinated seeds the activity was found to be maximum at pH 4.5 (figure 3). Results showed a considerable decrease in amylase activity as the pH increased.



**Figure 1:** Thin Layer Chromatography of enzyme hydrolyzed product of non-germinated (NG) and germinated (G) seed samples. (Glu- Glucose, Mal- Maltose)



**Figure 2:** Molecular weight of  $\beta$ -amylase determined on SDS-PAGE (PM-Protein marker, DNG- Dialyzed Non Germinated, DG- Dialyzed Germinated, PNG-Precipitated Non Germinated, PG- Precipitated Germinated, seed enzyme samples)

### Effect of temperature on $\beta$ -amylase activity


The enzyme was subjected to various temperatures ranging from 0°C - 100°C and its activity was analyzed. The activity of  $\beta$ -amylase in germinated and non germinated seeds were found to be maximum at temperature 37°C and 75°C (figure 4) respectively. The activity of enzyme gradually decreased with the increase in temperature.

### Effect of metal ions on $\beta$ -amylase activity


Various metal ions such as KCl,  $\text{CaCl}_2$ ,  $\text{HgCl}_2$ ,  $\text{CuCl}_2$ , SDS,  $\text{MgCl}_2$ , EDTA,  $\text{FeCl}_3$ ,  $\text{MnCl}_2$ ,  $\text{CoCl}_2$ ,  $\text{ZnCl}_2$ ,  $\text{AlCl}_3$  (at concentration of 2mM) were tested for activation or inhibition effects on the enzyme. There was a significant increase in enzyme activity in germinated seeds with  $\text{MnCl}_2$ ,  $\text{CoCl}_2$  (figure 5) and metals ions such as  $\text{CaCl}_2$ ,  $\text{CuCl}_2$ , SDS,  $\text{MgCl}_2$ ,  $\text{FeCl}_3$ ,  $\text{ZnCl}_2$ ,  $\text{AlCl}_3$ , KCl showed a negligible increase in the activity whereas  $\text{HgCl}_2$  and EDTA inhibited the activity of enzyme. In non germinated seeds the activity was inhibited by  $\text{HgCl}_2$  and metal ions such as  $\text{FeCl}_3$ ,  $\text{ZnCl}_2$ ,  $\text{MgCl}_2$ , EDTA,  $\text{CaCl}_2$  showed a slight increase in enzyme activity, whereas  $\text{CoCl}_2$ ,  $\text{AlCl}_3$  and KCl showed the highest enzyme activity which acts as a potent activator.

### Effect of substrate concentration and determination of $K_m$ and $V_{max}$

The enzymatic activity for germinated and non-germinated seed was determined using soluble starch as the substrate. The kinetic constant  $K_m$  and  $V_{max}$  was determined using Lineweaver Burke plot. The substrate concentration ( $K_m$ ) for germinated and non-germinated seeds was 3.03mg/ml and 5mg/ml respectively. The maximum velocity ( $V_{max}$ ) for germinated and non-germinated was found to be 6.6 $\mu\text{mol}/\text{min}/\text{ml}$  and 10 $\mu\text{mol}/\text{min}/\text{ml}$  respectively (figure 6 and figure 7).



**Figure 3:** Effect of pH for  $\beta$ -amylase activity in germinated and non germinated seed.



**Figure 4:** Effect of temperature for  $\beta$ -amylase activity in germinated and non germinated seed.

### Antioxidant studies

The enzyme showed highest amount of total antioxidant content of 304 $\mu$ g/ml in dialyzed enzyme sample from germinated seeds where as non germinated seeds showed 90.24 $\mu$ g/ml. ABTS free radical scavenging activity showed 76.09% which was higher with an  $IC_{50}$  value of 58.20 $\mu$ g/ml

for germinated seed enzyme when compared to non germinated seed with 54% of scavenging activity and  $IC_{50}$  value of 76.33 $\mu$ g/ml (table 3) (figure 8).

### Discussion

During seed germination various enzymes are involved in the growth of seedling such

as amylase, which is an important enzyme employed in the starch processing industries for the hydrolysis of polysaccharide (Sanni, 2000). Amylase has vast industrial applications. The enzymes in germinated and non germinated seeds are affected by various factors such as pH, temperature and metal ions. The present investigation was carried on to check if any variation in the above factors during germination and before germination. pH affect the activity of the  $\beta$ -amylase, whose activity is mostly high in acidic pH in both germinated and non germinated *Glycine max* seeds which supports the previous studies (Nitta et al., 1979; Toda and Svensson, 2000; Shen et al., 1988). The most important factor is the temperature, which is essential for all seed germination, also affects the enzyme activity. Our studies revealed that  $\beta$ -amylase are highly active at 37°C to 40°C which justifies from the work by lizotte et al, 1990, in pea epicotyls. It has also been reported that beta amylase is active at 70°C (Shen, 1988; Kirti, 2012), this agrees with the results obtained in

the present study. Beyond this range there is decline in the activity because of the Structural unfolding trasition at high temperature (Duy and fitter, 2005).

Various metal ions play a major role either as activator or an inhibitor for enzyme activity.  $MnCl_2$  and  $CoCl_2$  enhanced enzyme activity in germinated and non germinated seeds. This is in line with the findings of (Dahot et al., 2001) in *Moringa oleifera* seeds and (Dutta et al., 2006) in *Heliodiaptomus viduus*. Most of the findings have shown that  $HgCl_2$  is a potent inhibitor for enzyme activity. Lin et al., 1998 and Gupta et al., 2003, reports the inhibitory activity by  $Hg^{+}$  in microbial amylase which coincides with our present finding. As per the studies on beta amylase in plants, it shows that the molecular mass is between 50 to 60kDa (French, 1961; Thoma et al., 1972; Mikami et al., 1993; Joyce et al., 1998, Motoyasu et al., 1988), and the reports by Yoshiki Yamasaki, 2003 closely relates with our present findings but doesn't supports the findings of Gertler and Yehudith, 1965; Sarowar et al., 2009.






Figure 5: Effect of metal ions on  $\beta$ -amylase activity in germinated seeds and non germinated seed.






**Figure 6:** LB plot using soluble starch as substrate for germinated seed extract



**Figure 7:** LB plot using soluble starch as substrate for non germinated seed extract



**Figure 8:** ABTS scavenging activity of different extracts and its IC<sub>50</sub> (µg/ml) (AA- Ascorbic acid, NGC- Non Germinated Crude, NGD- Non Germinated Dialyzed, GC- Germinated Crude, GD- Germinated Dialyzed extracts).

Enzyme activity is also based on the substrate concentration ( $K_m$ ) and its maximum velocity ( $V_{max}$ ), the result obtained in the present research closely relates with the work on kinetic study by Femi-Ola and Ibikunle, 2013, and this also supports with the work by Deshwal Sapna, 2012; Sanni *et al.*, 2000. Low  $K_m$  activity indicates high affinity of the enzyme for the substrate (Hamilton *et al.*, 1998). Antioxidants has a potent capacity of inhibiting oxidation processes in human body as well as in food products, which was analyzed on our isolated enzyme from germinated and non germinated seeds, shown to be better in germinated seeds which agrees with the findings of Fernandez- Orozco *et al.*, 2008. Based on the present work, it can be speculated that  $\beta$ -amylase and antioxidants from germinated seeds of *Glycine max* can be employed for industrial purpose

### Acknowledgements

We acknowledge Dr. R. Chenraj Jain, President, Jain University Trust., Dr. N Sundararajan, Vice Chancellor, Jain University., Prof. K.S. Shantamani, Chief Mentor, Jain University and Dr. S Sundara Rajan, Director, CASB- Jain University, Bangalore for providing financial assistance, the necessary laboratory facilities and support. We also thank the staff at Genohelix Biolabs for technical support.

### References

- Akinlosottu A and Akinyele IO (1991) Effect of soaking, dehulling and fermentation on the oligosaccharides and nutrient content of cowpeas (*Vigna unguiculata*). Food Chem., 41: 43-53.
- Aleksandra Duda-Chodak and Tomasz Tarko (2007) Antioxidant properties of different fruit seeds and peels. Acta Sci. Pol, Technol. Aliment., 6(3): 29-36.
- Bilal Balkan and Figen Ertan (2005) Production and properties of  $\alpha$ -amylase from *Penicillium chrysogenum* and its application in starch hydrolysis. Prep. Biochem. Biotech., 35: 169-178.
- Brenfield P (1955) Amylases alpha and beta. In: Coloewick SP and Kalpan NO, eds. Method in Enzymology, 1: 149-151.
- Dahot, MU, Saboury AA, Ghobadi S and Moosavi-Movahedi AA (2001) Properties of the alpha amylase from *Moringa oleifera* seeds. J. Biol. Sci., 1: 747-749.
- Deshwal Sapna (2012) Effect of Substrate Concentration on the Amylase Activity in Germinating Buckwheat Seeds. Int. J. Res. Chem. Environ., 2: 148-149.
- Djordje Malenčić, Zoran Maksimović, Milan Popović. and Jegor Miladinović (2008) Polyphenol contents and antioxidant activity of soybean seed extracts. Bioresource Technol., 99: 6688-6691.
- Dubravka Štajner, Boris Popović M and Ksenija Taški (2009) Effects of  $\gamma$ -irradiation on antioxidant activity in soybean seeds, Cent. Eur. J. Biol., 4(3): 381-386.
- Dutta, T.K, Jana M, Pahari PR and Bhattacharya T (2006) The effect of temperature, pH and salt on amylase in *Heliodiaptomus viduus* (Gurney) (Crustacea: Copepoda: Calanoida). Turk. J. Zool., 30: 187-195.
- Duy C, Fitter J (2005) Thermostability of Irreversible Unfolding-Amylase Analysed by Unfolding Kinetics. J. Biological Chem., 280(45): 37360-37365.
- Emma M. Marinova, Nedyalka V. Yanishlieva (2003) Antioxidant activity and mechanism of action of some phenolic acid at an ambient and high temperatures. Food Chem., 81(2): 189-197.
- Femi-Ola Titilayo Olufunke and Ibikunle Ibidapo Azeez (2013) Purification and Characterization of Beta-Amylase of *Bacillus subtilis* Isolated from Kolanut Weevil, J. Bio. Life Sci. 4(1): 68-78.

- French D. (1961) The Enzymes, eds Boyer P. D., Lardy H., Myrbaek K. (Academic Press, Inc. New York), 2<sup>nd</sup> ed., 4:345-367.
- Gertler A and Yehudith Birk (1965) Purification and characterization of a  $\beta$ -amylase from soya beans. *Biochem J.*, 95(3): 621-627.
- Gupta, R, Gigars P, Mohapatra H, Goswami VK and Chauhan B (2003) Microbial  $\alpha$ -amylase: A biotechnological perspective. *Process. Biochem.*, 38: 1599-1616.
- Hamilton, L.M., Kelly C.T and Fogarty W.M(1998) *Carbohydrate Research.*, 314: 251-257.
- Ikram-ul-Haq, Abdullah R, Ashraf H and Shah AH (2002) Isolation and screening of fungi for the biosynthesis of alpha amylase. *Biotechnology*, 2: 61-66
- Joyce A., Gana, Newton E, Kalengamaliro, Suzanne Cunningham M, and Jeffrey J Volenec (1998) Expression of  $\beta$ -Amylase from Alfalfa Tap roots. *Plant Physiol.*, 118(4): 1495-1506.
- Kirti Rani (2012) Extraction and study of kinetic parameters of variety of sprouted pulses - amylases. *Int. J. of Pharm. & Life Sci.*, 3(8): 1893-1896.
- Komal Preet and Darshan Punia (2000) Proximate composition, phytic acid, polyphenols and digestibility (*in vitro*) of four brown cowpea varieties. *Int. J. Food Sci. Nutr.*, 51(3): 181-193.
- Laemmli U K (1970) Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. *Nature* 227: 680 - 685.
- Lin, L.L.; Chyau, C.C. and Hsu, W.H. (1998) Production and properties of a raw-starch-degrading amylase from thermophilic and alkaliphilic *Bacillus* sp. TS-23. *Biotechnol. Appl. Biochem.*, 28: 61-68.
- Lineweaver H and Burk D (1934) The determination of enzyme dissociation constants. *J. Am. Chem. Soc.*, 56 (3): 658-666.
- Lizotte PA., Henson CA., Duke SH (1990) Purification and characterization of pea epicotyl  $\beta$ -amylases, *Plant Physiol.*, 92: 615-621.
- Lowry, OH, Rosebrough NJ, Farr AL and Randall RJ (1951) Protein measurement with folin phenol reagent. *J. Biol. Chem.*, 193:265-275.
- Mikami B, Hehre EJ, Sato M, Katsube Y, Hirose M, Morita Y and Sacchettini JC (1993) The 2.0-Å resolution structure of soybean beta-amylase complexed with alpha-cyclodextrin. *Biochemistry* 32: 6836-6845.
- Motoyasu Adachi, Bunzo Mikami, Tomoyuki Katsube and Shigeru Utsumi (1998) Crystal Structure of Recombinant Soybean  $\beta$ -Amylase Complexed with  $\beta$ -Cyclodextrin, *J. Biol. Chem.*, 27: 19859-19865.
- Mubarak AE (2005) Nutritional composition and antinutritional factors of mung bean seeds (*Phaseolus aureus*) as affected by some home traditional processes. *Food Chem.*, 89 (4): 489-495.
- Negi A, Boora P and Khetarpaul N (2001) Starch and protein digestibility of newly released moth bean cultivars: Effect of soaking, dehulling, germination and pressure cooking. *Mol Nutr Food Res.*, 45 (4): 251-254.
- Nitta Y, Kinikata T and Watanabe T (1979) Kinetic study of soybean beta-amylase. The effect of pH, *Journal of biochemistry* 85(1): 41-45.
- Prieto P., M. Pineda and M. Aguilar (1999) Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Anal. Biochem.*, 269: 337 -341.
- Fernandez-Orozco, Rebeca, Frias, Juana, Zielinski, Henryk Piskula, Mariusz K., Kozłowska, Halina; Vidal-Valverde, Concepción (2008) Kinetic study of the antioxidant compounds and antioxidant capacity during germination of *Vigna radiata* cv. emmerald, *Glycine max* cv. jutro

- and *Glycine max* cv. merit. Food Chemistry, 111 (3): 622-630
- Re R., N. Pellegrini, A. Proteggente, A. Pannala, M. Yang and C. Rice-Evans (1999) Antioxidant activity applying an improved ABTS radical cation decolorization assay. Free Radic. Biol. Med., 26: 1231 -1237.
- Richardson, T. H, Tan, X, Frey, G, Callen, W, Cabell, M, Lam, D, Macomber, J, Short, J M, Robertson, D E. and Miller, C J (2002) A novel, high performance enzyme for starch liquefaction. Discovery and optimization of a low pH, thermostable alpha-amylase. J. Biol. Chem. 277(29): 26501-26507.
- Sanni, Tajudeen Morakinyo (2000) Isolation, purification and characterization of  $\beta$ -amylase from African Oil bean (*Pentaclethra mactothylla*). Ph.D thesis. Federal University of Technology, Akure.
- Sarowar Jahan MG, Shaela Pervin M, Rowshanul Habib M, Farjana Nikkon and Habibur Rahman (2009) Purification and characterization of  $\beta$ -amylase from radish (*Raphanus sativus* L.) root. J. Appl. Sci. Res. 5(12):2225-2233.
- Shelby N. Freer (1993) Purification and Characterization of the Extracellular  $\alpha$ -Amylase from *Streptococcus bovis* JB1. Appl. Environ. Microbiol., 59(5), 1398-1402.
- Shen G.J, Saha B C, Lee Y E, Bhatnagar L and Zeikus G (1988) Purification and characterization of a novel thermostable beta-amylase from *Clostridium thermosulphurogenes*. Biochem. J. 254(3): 835-840.
- Sumathi A, Malleshi NG and Rao SV (1995) Elaboration of amylase activity and changes in past viscosity of some common Indian legumes during germination. Plant Foods Hum. Nutr., 47: 341-347.
- Thoma J A, Spradlin J E and Dygert S (1972) in *The Enzymes*, ed Boyer P. D. (Academic Press, Inc. New York), 3rd Ed. 5:115-189.
- Toda R and Svensson L (2000) Partial hydrolysis of sweet potato starch with  $\beta$ -amylase. Agric. Chem., 51: 1365-1371.
- Wanasundara PK, JPD, Shahidi F and Shukla VKS (1997) Endogenous Antioxidants from Oil Seeds and Edible Oil. Marcel Dekker, Inc. 225-292.
- Wang SY & HS Lin (2000) Antioxidant activity in fruits and leaves of blackberry, raspberry, and strawberry varies with cultivar and developmental stage. J. Agri. Food Chem., 48(2): 140-146.
- Yoshiki Yamasaki (2003),  $\beta$ -Amylase in germinating millet seeds. Phytochem., 64(5): 935-939.