

Regular Article

**Evidence of *WUSCHEL* (*WOX2*) gene expression during induction of somatic embryogenesis from apical shoot buds of mature trees of *Pinus roxburghii***

Ravindra B. Malabadi<sup>1\*</sup>, K. Nataraja<sup>1</sup>, S. Vijaya Kumar<sup>2</sup>, Gangadhar S. Mulgund<sup>1</sup>

<sup>1</sup>Department of Botany, Karnatak University, Pavate nagar, Dharwad-580003, Karnataka, India

<sup>2</sup>Department of Biotechnology, Madanapalle Institute of Technology and Science, Madanapalle-517325, Chittoor District, Andhra Pradesh, India

\*Corresponding author E-mail: [mlbd712@rediffmail.com](mailto:mlbd712@rediffmail.com)

**This study highlights for the first time the expression of transcription factor, *WOX2* (*WUSCHEL* homeobox 2) in the embryogenic tissue derived from the apical shoot buds of mature trees of 3 genotypes of *P. roxburghii*. Therefore, *PrWOX2* could be used as a potential molecular signature for the identification of embryogenic cultures during the early development of somatic embryos in conifers. On the other hand, the non-embryogenic tissue of shoot bud cultures which were initiated without cold treatment (control) failed to show the expression of *WOX2*. This clearly demonstrates the role of *WOX2* in the somatic embryogenesis pathway and might be directly related to the stress conditions.**

**Key words:** Apical buds, Chir pine, meristematic tissue, pines, somatic embryogenesis,

Somatic embryogenesis is the development of embryo-like structures under *in vitro* conditions on tissues derived from somatic cells (Konar and Nataraja, 1965; Nataraja and Konar, 1970; Feher *et al.* 2003; Malabadi *et al.* 2011a, 2011b, 2011c, 2011d, 2011e, 2011f, 2011g). During cloning of mature conifers, isolated somatic cells (stem cells) under any external stress conditions of cold/heat or chemical are induced to form a somatic embryo. Embryogenic systems derived from vegetative shoot apices or secondary needles of mature pines have been well established in at least a few conifers (Litz *et al.* 1995; Malabadi and van Staden, 2003, 2005a, 2005b, 2005c, 2006; Malabadi, 2006; Malabadi and Nataraja, 2006a, 2006b; Aronen *et al.* 2007, 2008; Malabadi and Nataraja,

2007a, 2007c, 2007f, 2007g; Malabadi *et al.* 2008a, 2008b, 2008c; Park *et al.* 2009a; Malabadi and Teixeira da Silva, 2011). Furthermore, the tissue used, developmental phase of the source plant, stress source, stress chemical concentration, and duration of stress are important for induction of somatic embryos of mature conifers (Malabadi *et al.* 2004). In addition, these factors are very sensitive and affect each other. The fact indicates that each stress induces a common reaction, and the reaction is related to the plant somatic cell de-differentiation and re-differentiation to somatic embryogenesis (Ikeda-Iwai *et al.* 2003). Probably, each stress treatment induces expression of a factor that controls the start of somatic embryogenesis, through the common reaction (Ikeda-Iwai *et*

al. 2003). Since most somatic cells are not naturally embryogenic, an induction phase is required for the cells to acquire embryogenic competence (Konar and Nataraja, 1965; Feher et al. 2003; Ikeda-Iwai et al. 2003; Namasivayam, 2007; Malabadi et al. 2009a). This is in contrast to the zygote in sexual reproduction which is intrinsically embryogenic. Stress conditions, for example temperature-related, hormonal or osmotic, can influence the fate of tissue cultured plants cells i.e. induce either apoptosis or a developmental switch, and this phenomenon could be utilized for initiation of somatic embryogenesis as well (Feher et al. 2003; Ikeda-Iwai et al. 2003; Namasivayam, 2007; Malabadi et al. 2009a). The molecular bases of this metabolic and developmental switch are poorly understood despite extensive research with different tissue culture systems.

During the past 2 decades, considerable efforts have been made to identify genes with altered expression pattern during somatic embryogenesis (Zuo et al. 2002; Arroyo-Herrera et al. 2008; Cairney and Pullman, 2007; Malabadi and Nataraja, 2007g; Malabadi et al. 2009b). Most of these genes however, are up regulated only in the developmental stages, suggesting that they do not play a direct role in the vegetative-to-embryogenic transition (Zuo et al. 2002; Malabadi and Nataraja, 2007g; Malabadi et al. 2009b). Cell differentiation depends on the proper and sequential expression of key genes required for somatic embryogenesis. Several aspects of control are required for this which includes: chromatin modifications, DNA methylation, correct amount of particular transcription factors, proper nuclear arrangement, etc (Malabadi and Nataraja, 2007g; Malabadi et al. 2009b). There are different groups or class of genes expressed during induction of somatic embryogenesis. For example, carrot somatic embryogenesis receptor kinase (*SERK*) gene, whose expression appears to mark

vegetative-to-embryogenic transition; however its function remains unclear. Over expression of the *Plant Growth Activator 6* (*PGA6*) gene promotes the formation of somatic embryos from various vegetative tissues, as well as from zygotic embryos independent of any external plant hormones (Malabadi et al. 2009b). These somatic embryos following a developmental process remarkably similar to that of zygotic embryogenesis were able to germinate and grow healthy, fertile plants, suggesting that *PGA6* is involved in the maintenance of embryonic cell identity. *PGA6* was found to be identical to *WUS*, a homeodomain protein previously characterized as a key regulator for the specification of meristem cell fate. During the last few years the homeobox transcription factor *WUSCHEL* (*WUS*) has been shown to cause dedifferentiation when expressed on somatic cells followed by a production of new stem cells that can lead to somatic embryogenesis or organogenesis. The homeobox gene *WUSCHEL* (*WUS*) is required to specify stem cell identity (Baurle and Laux, 2005; Leibfried et al. 2005). *WUSCHEL* (*WUS*) was originally identified as a central regulator of shoot and floral meristems in *Arabidopsis*, where it was expressed in a small group of cells, and is required to maintain the overlying stem cells undifferentiated (Baurle and Laux, 2005; Leibfried et al. 2005). Among all the genes, *WUSCHEL* homeobox (*WOX2*) in *Picea abies* (Palovaara and Hakman, 2008; Palovaara et al. 2010) is one of the important putative marker gene expressed during the early development of somatic embryos. The transcription factor, *WOX2* (*WUSCHEL* homeobox 2), which is one of the 15 *WOX* family proteins, play an important role in controlling many aspects of somatic embryogenesis in plants (Palovaara and Hakman, 2008; Palovaara et al. 2010). In the higher plants, some genes and transcription factors related to important mechanisms of embryogenesis are plant specific. Therefore, new findings such as discovery and

identification of new transcription factors that control the expression of embryogenesis-related genes and embryogenesis-specific phytohormone response be made in a near future. In our previous study, we were able to isolate cDNA clones of genes expressed in embryogenic and non-embryogenic tissue derived from the vegetative shoot apices of mature trees of *P. roxburghii* (Malabadi and Nataraja, 2007g). Furthermore, we have also constructed cDNA library in our previous study (Malabadi and Nataraja, 2007g). This study is the continuation of our previous work and highlights for the first time the confirmation of expressed gene particularly *WUSCHEL (WOX2)* (Palovaara and Hakman, 2008; Palovaara et al. 2010) in embryogenic tissue derived from the apical shoot buds of mature trees of *P. roxburghii*.

## Materials and methods

### Initiation of embryogenic tissue

Shoot apical domes from mature trees (14- years old) of *Pinus roxburghii* of 3 genotypes (PR11, PR105, and PR521) were collected from the Western Ghat Forests, India (14° 5' to 15° 25' N latitude and 74° 45' to 76° E longitude with an average rainfall of 80 cm.). They were cleansed with 1% Citramide (Sodium hypochlorite 3.5%) for 5 min and rinsed thoroughly with sterilized distilled water. These were surface decontaminated with 70% ethanol for 5 min followed by immersion in 0.5% HgCl<sub>2</sub> for 2 min and rinsed 4-times with sterile double distilled water. Transverse-thin sections of approximately 0.5-1.0 mm thick were cut using sharp sterilized blade or scalpel from shoot apical domes (upper part with 2 to 3 sections only) for the initiation of embryogenic tissue. These shoot apical dome sections were cultured individually on full strength (Inorganic salts) DCR (Gupta and Durzan, 1985) medium containing 0.2 g l<sup>-1</sup> polyvinyl pyrrolidone (PVP), 2.0 g l<sup>-1</sup> Gellan gum (Sigma), 30 g l<sup>-1</sup> maltose (Analar, Sigma) and 0.3 % activated charcoal (Sigma)

without growth regulators. The cultures were raised in 25 mm X145 mm glass culture tubes (Borosil) containing 15 ml of medium. These cultures were incubated in dark at 4° C for 3 days. Thin apical dome sections after incubation in dark at 4° C for 3 days were subcultured on full strength DCR medium supplemented with 22.62 µM 2, 4-D, 26.85 µM NAA, and 8.87 µM BA (initiation medium) (Malabadi and Nataraja, 2006ab). The pH of the medium was adjusted to 5.8 with NaOH or HCl before Gellan gum was added. The media were then sterilized by autoclaving at 121° C and 1.05 kg/cm<sup>2</sup> for 15 min. L-glutamine and casein hydrosylate were filter sterilized and added to the media after it had cooled to below 50° C. All the cultures were maintained in the dark at 25 ± 2° C with 55-60% relative humidity.

All the cultures were maintained in the dark at 25 ± 2° C for four weeks. Embryogenic tissue was initiated according to our previous existing protocols (Malabadi et al., 2004; Malabadi and Nataraja, 2006ab; Malabadi and van Staden, 2006). For the initiation of non-embryogenic tissue, thin sections of shoot apical dome without cold-pretreatment subcultured on the initiation medium (Malabadi and Nataraja, 2006ab) were served as the control. They were also maintained in dark for four weeks.

### Maintenance of embryogenic tissue

The embryogenic tissue showing proembryonal masses was again subcultured onto maintenance medium. The full-strength DCR(Gupta and Durzan, 1985) basal medium containing 60 g l<sup>-1</sup> maltose, 2 g l<sup>-1</sup> Gellan gum supplemented with 2.26 µM 2,4-D, 2.68 µM NAA and 0.88 µM BA (maintenance medium) was used for this purpose in accordance with our existing previous protocols (Malabadi and Nataraja, 2006ab). On the maintenance medium, embryonal suspensor masses were cultured for 30 days with two subcultures. The presence of embryonal masses was

determined by morphological and microscopic observations. The non-embryogenic cultures (control) were also subcultured on the maintenance medium and maintained in dark condition.

For the following experiments of *WUSCHEL* (*WOX2*) (Palovaara and Hakman, 2008) gene expression studies, two types of plant material, 1) embryogenic tissue initiated by cold-pre treatment, and 2) control (non-embryogenic tissue) induced without cold pre-treatment have been used for the isolation of total RNA. For total RNA isolation, one gram fresh wt of embryogenic tissue and control (non-embryogenic tissue) were then placed in a cryostorage vial partially immersed in liquid nitrogen. Ten vials of plant tissues (embryogenic tissue and non-embryogenic tissue-control) were collected. Frozen tissues were stored at -70°C until further analyses were performed.

#### RNA preparation and cDNA synthesis

Frozen samples of embryogenic tissue and control (non-embryogenic tissue) derived from the apical shoot buds of *Pinus roxburghii* of 3 genotypes (PR11, PR105, and PR521) were ground in a mortar and pestle with liquid nitrogen. Total RNA was isolated according to the modified method of Chang et al. (1993). To remove residual genomic DNA, 25 µg of RNA was treated with TURBO-DNase™ (Ambion, Austin, TX, USA). cDNA was generated from 1 µg of DNase-treated RNA using the Superscript II RT system (Invitrogen, CA, USA) according to the manufacturer's protocol (Park et al. 2009). Each reaction was run in duplicate, generating two independent cDNA samples for each RNA sample isolated from 3 genotypes (PR11, PR105, and PR521) of *P. roxburghii* as reported in our previous study (Malabadi and Nataraja, 2007g).

#### Cloning of *P. roxburghii* *WOX2*

For the isolation of the *WOX2* gene, following gene specific primers were

designed based on homologous sequences publically available, including PaWOX2 (*P. abies* *WOX2*, Acc. AM286747) and PtWOX2 (*P. taeda* *WOX2*, Acc. DR693345; Cairney et al. 2006; Park et al. 2009). The gene specific primers were custom synthesized and used for the PCR amplification of the PR *WOX2* gene using cDNA as the template. Gene alignment was done using Bio Edit programme. Following PCR amplification from *P. roxburghii* cDNAs, the resulting amplicons were cloned into the TOPO-TA cloning vector (Invitrogen) (Park et al. 2009). The gene *WOX2* was confirmed by the DNA sequence analysis. (Forward: 5'-ATG GCC GAG GGT CAA TCC ACC ATG A-3'); (Reverse 3' CTT GCC AGG ATG CTG AGG GAT A-5')

#### Maturation of somatic embryos

For maturation, embryogenic tissue clumps of each of the 3 genotypes (PR11, PR105, and PR521) were partially desiccated. One gram fresh weight of tissue of each embryogenic line were transferred to sterile empty Petri dishes (60 mm diam.) containing two sterile Whatman filter paper disks (50 mm) (Schleicher and Schuell, qualitative circles). The Petri dishes were sealed with Parafilm and kept at 25 ± 2°C in the dark for 24 hr to obtain the desired extent of desiccation. After desiccation, the partially desiccated tissue of each embryogenic line was transferred to maturation medium to induce cotyledonary embryo development. The full strength DCR (Gupta and Durzan, 1985) medium with 60 g l<sup>-1</sup> maltose, 37.84 µM abscisic acid (ABA) and 5 g l<sup>-1</sup> Gellan gum (maturation medium) was tested for this purpose. All the cultures were placed in the dark at 25 ± 2°C and these maintained for 8 to 12 weeks (Malabadi, 2006b; Malabadi and Nataraja 2006a, 2006b).

#### Germination and plantlet recovery

After 6 to 8 weeks of maturation in the presence of ABA and higher concentrations of

maltose (60 g l<sup>-1</sup> maltose) the cotyledonary somatic embryos were recovered from the cultures for germination. Before germination, the cotyledonary somatic embryos of the all embryogenic lines were cold pre-treated at 2°C and kept in dark for 5 days. The germination medium (GM) used was half DCR (Gupta and Durzan, 1985) medium with 2 g l<sup>-1</sup> Gellan gum (Malabadi and Nataraja 2006a, 2006b). In the first week of germination, cultures were kept in darkness, and then transferred to diffuse light in the second week, and thereafter to a 16-hr photoperiod under a light intensity of 50 µmol m<sup>-2</sup> s<sup>-1</sup> for hardening. Somatic embryos were considered germinated as soon as radical elongation occurred and conversion to plantlets was based on the presence of epicotyls. After 4 to 6 weeks on germination medium, plantlets were transferred to vermiculite in a controlled growth room. All the experiments were repeated over 3 different and consecutive years. Data were arcsine transformed before being analyzed for significance using ANOVA (analysis of variance, p<0.05) or evaluated for independence using a Chi-square test. Further, the differences were contrasted using a Duncan's multiple range test (α=0.05) following ANOVA. All statistical analysis was performed using SPSS statistical software package version 13.0.1.1, Microsoft Windows.

### Results and discussion

In our present study, we were able to differentiate both embryogenic and non-embryogenic tissue on the basis of gene expression studies. We have established a reproducible protocol for the induction of somatic embryogenesis using apical shoot buds of mature trees of many recalcitrant pines (Malabadi and van Staden, 2003, 2005a, 2005b, 2005c, 2006; Malabadi, 2006; Malabadi and Nataraja, 2006a, 2006b; Aronen *et al.* 2007, 2008; Malabadi and Nataraja, 2007a, 2007c, 2007f, 2007g; Malabadi *et al.* 2008a, 2008b, 2008c; Park *et al.* 2009a; Malabadi and

Teixeira da Silva, 2011). This is one of the major breakthroughs in the forest biotechnology. Embryogenic tissue derived from apical shoot buds could be easily separated and identified on the basis of microscopic observation. This is the first hand information for the confirmation of induction of somatic embryogenesis from shoot buds of mature trees of recalcitrant pines. The second most important feature for the identification of somatic embryogenesis is the nature, and type of the cells which was well documented and demonstrated in our previous papers (Malabadi and van Staden, 2003, 2005a, 2005b, 2005c, 2006; Malabadi, 2006; Malabadi and Nataraja, 2006a, 2006b; Aronen *et al.* 2007, 2008; Malabadi and Nataraja, 2007a, 2007c, 2007f, 2007g; Malabadi *et al.* 2008a, 2008b, 2008c; Park *et al.* 2009a; Malabadi and Teixeira da Silva, 2011). In case of pines, the transverse thin layer of apical shoot buds showed outermost epidermis layer, then internal layer of cortex region, followed by thin cambial region and central pith or medullar region respectively. The active dividing and totipotent cells (stem cells) are positioned only at the cambial layer of the apical meristematic tissue in conifers, so that their growth and division leads to a continuous flow of progeny cells. These cambial layer cells (stem cells) under stress conditions such as cold/heat or chemical treatment undergo differentiation and leading to the embryogenic pathway in conifers. But activation of the cambial layer cells is one of the important steps for the successful induction of somatic embryogenesis in conifers. This has been achieved and successful in many recalcitrant conifers (Malabadi and van Staden, 2003, 2005a, 2005b, 2005c, 2006; Malabadi, 2006; Malabadi and Nataraja, 2006a, 2006b; Aronen *et al.* 2007, 2008; Malabadi and Nataraja, 2007a, 2007c, 2007f, 2007g; Malabadi *et al.* 2008a, 2008b, 2008c; Park *et al.* 2009a; Malabadi and Teixeira da Silva, 2011). These cells (stem cells) are very small in size

and slow in growth at the initial stages of the development. The embryogenic cells were small, richly cytoplasmic and actively dividing with a prominent nucleus. These cells were generally present in small, compact aggregates, rich in starch grains and showed competence for embryogenesis. On the other hand the rest of the layers (epidermis, cortex region and central pith or medulla) have induced nonembryogenic tissue in conifers. Microscopic observation of non-embryogenic tissue confirmed loosely arranged, elongated thin paracymatous cells. The non-embryogenic tissue is composed of large cells, vacuolated and often elongated with sparse cytoplasm and few starch grains. These cells can survive up to 6 months with multiple subcultures, ultimately leading to death due to the exudation of high phenolic compounds and other factors. Therefore, there is a clear distinction between embryogenic and non-embryogenic tissue on the basis of nature, and type of the cells without any molecular marker. A well experienced Scientist in plant tissue culture can recognize and identify the embryogenic cells very easily without the help of molecular marker. Therefore, inducing somatic embryogenesis using apical shoot buds of mature conifers is an important step in the commercial forestry. Therefore, somatic embryogenesis is still a very challenging process and considered as a fundamental science but not a technique. On the other hand plant tissue culture is an important *in vitro* technique.

For further characterization of the embryogenic tissue, we have developed a potential molecular marker on the basis of the expression of one of the important transcription factor, *WOX2* (*WUSCHEL* homeobox 2) in the embryogenic tissue of *P. roxburghii*. Transcription factor, *WOX2* (*WUSCHEL* homeobox 2) was strongly expressed in the embryogenic tissue but not

in non-embryogenic tissue (control). In our present study, embryogenic tissue was developed under cold stress conditions. On the other hand non-embryogenic tissue was induced without cold treatment. Expression of *WOX2* (*WUSCHEL* homeobox 2) in the embryogenic tissue of 3 genotypes (PR11, PR105, and PR521) of *P. roxburghii* clearly confirmed its involvement in the somatic embryogenic pathway and might be directly related to stress conditions. Therefore, stress conditions always induce somatic embryogenesis in many plant species which also directly related to the expression of genes. This is the first evidence of expression of *WOX2* in *P. roxburghii* and could be used as marker for the identification of embryogenic tissue in pines. In gymnosperms, no *WUS* homolog was found (Palovaara and Hakman, 2008) and as such it is believed that the *WOX* and *WUS* genes have not diverged in some species (Kiselev *et al.* 2009; Park *et al.* 2009). An interrogation of the *WOX* family members in gymnosperms obtained from the public databases identified that *PrWOX2* has 80% similarity to *P. abies PaWOX2* and is almost identical to other pine species sequence such as *PsWOX2* of *Pinus sylvestris* (99% homology) and *PtWOX2* of *P. taeda* (100% homology) (Park *et al.* 2009). These findings are consistent with previous findings (Malik *et al.* 2007, Palovaara and Hakman, 2008, Palovaara *et al.* 2010) which suggested that *WOX2* may be a potential marker to predict the embryogenic potential of spruce and *Brassica* cultivars (Park *et al.* 2009). Therefore, on the basis of this wonderful study by our laboratory further strengthens the concept of cloning of mature trees of pines. *PrWOX2* could be used as a potential genetic marker for the identification of the embryogenic cultures, which is just an added advantage for developing SE protocols for recalcitrant pines.

## References

- Aronen TS, Pehkonen T, Malabadi RB, Ryyanen L (2008) Somatic embryogenesis of Scots pine-advances in pine tissue culture at Metla. Vegetative propagation of conifers for enhancing landscaping and tree breeding. Proceedings of the Nordic meeting held in September 10<sup>th</sup>-11<sup>th</sup> 2008 at Punkaharju, Finland. *Working Papers of the Finnish Forest Research Institute*, **114**: 68-71.
- Aronen TS, Ryyanen L, Malabadi RB (2007) Somatic embryogenesis of Scots pine: initiation of cultures from mature tree explants and enhancement of culture system [Abstract]. In: *IUFRO Tree Biotechnology Conference*, June 3-8, 2007, Ponta Delgada, Azores, Portugal, No. SIX. 2.
- Arroyo-Herrera A, Gonzalez AK, Moo RC, Quiroz-Figueroa FR, Loyola-Vargas VM, Rodriguez-Zapata LC, Burgeff D' Hondt C, Suarez-Solis VM, Castano E (2008) Expression of *WUSCHEL* in *Coffea canephora* causes ectopic morphogenesis and increases somatic embryogenesis. *Plant Cell Tissue and Organ Culture*, **94**: 171-180.
- Baurle I and Laux T (2005) Regulation of *WUSCHEL* transcription in the stem cell niche of the *Arabidopsis* shoot meristem. *The Plant Cell*, **17**:2271-2280.
- Cairney J, Zheng L, A. Cowels A et al. 2006. Expressed sequence tags from loblolly pine embryos reveal similarities with angiosperm embryogenesis. *Plant Mol. Biol*, **62**:485-501.
- Cairney J and Pullman GS (2007) The cellular and molecular biology of conifer embryogenesis. *New Phytologist*, **176**: 511-536.
- Chang S, Puryear J and Cairney J (1993) A simple and efficient method for isolating RNA from pine trees. *Plant Molecular Biology Reports*, **11**: 113-116.
- Feher A, Pasternak TP, Dudits D (2003) Transition of somatic plant cells to an embryogenic state. *Plant Cell, Tissue and Organ Culture*, **74**: 201-228.
- Gupta PK and Durzan DJ (1985) Shoot multiplication from mature trees of Douglas fir and sugar pine. *Plant Cell Reports*, **4**: 177-179.
- Ikeda-Iwai M, Umehara M, Satoh S, Kamada H (2003) Stress-induced somatic embryogenesis in vegetative tissues of *Arabidopsis thaliana*. *Plant Journal*, **34**: 107-114.
- Kiselev KV, Turlenko AV, Zhuravlev YN (2009) PgWUS expression during somatic embryo development in a *Panax ginseng* 2c3 cell culture expressing the rolC oncogene. *Plant Growth Reg*, **59**:237-243.
- Konar RN and Nataraja K (1965) Experimental studies in *Ranunculus sceleratus* L. Development of embryos from the stem epidermis. *Phytomorphology*, **15**: 132-137.
- Leibfried A, To JPC, Busch W, Stehling S, Kehle A, Demar M, Kieber JJ and Lohman JU (2005) *WUSCHEL* controls meristem function by direct regulation of cytokinin-inducible response regulators. *Nature*, **438**/22:1172-1175.
- Litz RE, Moon PA and Chavez VM (1995) Somatic embryogenesis from leaf callus derived from mature trees of the cycad *Ceratozamia hildae* (Gymnospermae). *Plant Cell, Tissue and Organ Culture*, **40**: 25-31.
- Malabadi RB and van Staden J (2003) Somatic embryos can be induced from shoot apical domes of mature *Pinus patula* trees. *South African Journal of Botany*, **69**: 450-451.
- Malabadi RB, Choudhury H and Tandon P (2004) Initiation, maintenance and

- maturation of somatic embryos from thin apical dome sections in *Pinus kesiya* (Royle ex. Gord) promoted by partial desiccation and Gellan gum. *Scientia Horticulture*, **102**: 449-459.
- Malabadi RB and van Staden J (2005a) Somatic embryogenesis from vegetative shoot apices of mature trees of *Pinus patula*. *Tree Physiology*, **25**: 11-16.
- Malabadi RB and van Staden J (2005b) Role of antioxidants and amino acids on somatic embryogenesis of *Pinus patula*. *In Vitro Cellular and Developmental Biology- Plant*, **41**: 181-186.
- Malabadi RB and van Staden J (2005c) Storability and germination of sodium alginate encapsulated somatic embryos derived from the vegetative shoot apices of mature *Pinus patula* trees. *Plant Cell Tissue and Organ Culture*, **82**: 259-265.
- Malabadi RB and van Staden J (2006) Cold-enhanced somatic embryogenesis in *Pinus patula* is mediated by calcium. *South African Journal of Botany*, **72**: 613-618.
- Malabadi RB (2006) Effect of glutathione on maturation of somatic embryos derived from vegetative shoot apices of mature trees of *Pinus roxburghii*. *Journal of Phytological Research*, **19**: 35-38.
- Malabadi RB and Nataraja K (2006a) Cryopreservation and plant regeneration via somatic embryogenesis using shoot apical domes of mature *Pinus roxburghii* Sarg. trees. *In Vitro Cellular and Developmental Biology - Plant*, **42**: 152-159.
- Malabadi RB and Nataraja K (2006b) RAPD detect no somaclonal variation in cryopreserved cultures of *Pinus roxburghii* SARG. *Propagation of Ornamental Plants*, **6**: 114-120.
- Malabadi RB and Nataraja K (2007a) Smoke-saturated water influences somatic embryogenesis using vegetative shoot apices of mature trees of *Pinus wallichiana* A. B. Jacks. *Journal of Plant Sciences*, **2**: 45-53.
- Malabadi RB and Nataraja K (2007c) Genetic transformation of conifers: Applications in and Impacts on commercially forestry. *Transgenic Plant Journal*, **1**(2): 289-313.
- Malabadi R. B., Nataraja K (2007f) Plant regeneration via somatic embryogenesis using secondary needles of mature trees of *Pinus roxburghii* Sarg. *International Journal of Botany*, **3**: 40-47.
- Malabadi RB and Nataraja K (2007g) Isolation of cDNA clones of genes differentially expressed during somatic embryogenesis of *P. roxburghii*. *American Journal of Plant Physiology*, **2**: 333-343.
- Malabadi RB, Teixeira da Silva JA, Nataraja K (2008a) A new approach involving salicylic acid and thin cell layers for cloning mature trees of *Pinus roxburghii* (Chir Pine). *The American Journal of Plant Science and Biotechnology*, **2**(2): 56-59.
- Malabadi RB, Teixeira da Silva JA, Nataraja K (2008b) Salicylic acid induces somatic embryogenesis from mature trees of *Pinus roxburghii* (Chir pine) using TCL technology. *Tree Forestry Science and Biotechnology*, **2**(1): 34-39.
- Malabadi RB, Mulgund GS, Vijaykumar S (2009a) How somatic cells follows embryogenic pathway during cloning mature trees of conifers? *Journal of Phytological Research*, **22**(1): 53-56.
- Malabadi RB, Mulgund GS, Vijaykumar S (2009b) Expression of *WUSCHEL*-gene promoting somatic embryogenesis in plants. *Journal of Phytological Research*, **22** (1): 103-106.
- Malabadi RB and Teixeira da Silva JA (2011a) Thin cell layers: Application to

- forestry biotechnology. *Tree and Forestry Science and Biotechnology*, 5(1): 14-18.
- Malabadi RB, Teixeira da Silva JA, Mulgund GS (2011b) Induction of somatic embryogenesis in *Pinus caribaea*. *Tree and Forestry Science and Biotechnology*, 5(1), 27-32.
- Malabadi RB, Teixeira da Silva JA, Nataraja K, Vijaya Kumar S, Mulgund GS (2011c) Induction of somatic embryogenesis in mature coniferous forest trees. *Research in Biotechnology*, 2(5):08-33.
- Malabadi RB, Teixeira da Silva JA, Nataraja K, Vijaya Kumar S, Mulgund GS (2011d) Induction of somatic embryogenesis in mango (*Mangifera indica*). *International Journal of Biological Technology*, 2(2):12-18.
- Malabadi RB, Nataraja K, Vijaya Kumar S, Mulgund GS (2011e) Journey of a single cell to a plantlet *via in vitro* cloning mature trees of conifers. *Research in Biotechnology*. 2(6):01-07.
- Malabadi RB, Mulgund GS, Nataraja K, Vijaya Kumar S (2011f) Induction of somatic embryogenesis and plant regeneration in different varieties of sugarcane (*Saccharum officinarum* L.). *Research in Plant Biology*. 1(4):39-41.
- Malabadi RB, Mulgund GS, Nataraja K, Vijaya Kumar S (2011g) Induction of somatic embryogenesis in Papaya (*Carica papaya* L.). *Research in Biotechnology*, 2(5):40-55
- Malik MR, Wang F, Dirpaul JM, Zhou N, Polowick PL, Ferrie AMR, Krochko JE, (2007) Transcript profiling and identification of molecular markers for early microspore embryogenesis in *Brassica napus*. *Plant Physiol*, 144:134-154.
- Namasivayam P (2007) Acquisition of embryogenic competence during somatic embryogenesis. *Plant Cell, Tissue and Organ Culture*, 90: 1-8.
- Nataraja K and Konar RN (1970) Induction of embryoids in reproductive and vegetative tissues of *Ranunculus sceleratus* L. *in vitro*. *Acta Botanica Neerlandica*, 19: 707-716.
- Palovaara J and Hakman I (2008) Conifer WOX-related homeodomain transcription factors, developmental consideration and expression dynamic of WOX2 during *Picea abies* somatic embryogenesis. *Plant Mol. Biol*, 66:533-549.
- Palovaara J, Hallberg H, Stasolla C, Hakman I (2010) Comparative expression pattern analysis of WUSCHEL-related homeobox 2 (WOX2) and WOX8/9 in developing seeds and somatic embryos of the gymnosperm *Picea abies*. *New Phytol*. doi: 10.1111/j.1469-8137.2010.03336.x.
- Park SY, Klimaszewska K, Malabadi RB, Mansfield SD (2009) Embryogenic cultures of lodgepole pine originating from mature trees and from immature seed explants. IUFRO Tree Biotechnology Conference, June 28<sup>th</sup>-July 2<sup>nd</sup> 2009, Whistler, BC, Canada, p 60 (abstract).
- Zuo J, Niu QW, Frugis G, Chua NH (2002) The WUSCHEL gene promotes vegetative-to embryonic transition in *Arabidopsis*. *The Plant Journal*, 30(3):349-359.