

Mini Review

Arbuscular Mycorrhiza: Their distribution and association with plants in the revegetated mine spoils of India – an overview

Logaprabha V.* and Tamilselvi K. S.

Department of Botany, PSGR Krishnammal College for Women, Coimbatore, Tamil Nadu,
India

Corresponding author Email: tamilselviks@yahoo.com

Reclamation of mine spoils is receiving considerable attention in the recent years due to the ill effects of mining and dumping of the mine spoil waste leading to various environmental hazards. Mycorrhizal inoculation of mine spoils and mine spoil dumps is essential for the establishment of a diverse plant community and a better ecosystem. This review focuses on the status of AM fungi, their distribution, diversity and density in the various mine spoils and mine spoil dumps of India. Species belonging to the genus *Glomus* was found to be widely distributed among the mine spoils. Legumes were the dominant and tolerant plants of the mine spoils supporting mycorrhizal colonisation.

Keywords: AM fungi, *Glomus*, India, legumes, mine spoils.

Arbuscular mycorrhiza helps in the maintenance of plant biodiversity, microbial community, ecosystem stability and function. They play an indomitable role in the restoration of the degraded lands. India is the second largest country with a total land area of 5.7 lakh sq.kms in working mines. Mining leads to excessive soil damage by altering the natural ecosystem, ecological integrity and the mycorrhizal network prevailing in the soil. In spite of the vast mining operations and serious hazards involved due to mining, only very little research has been done on the mycorrhizal status in the mine spoils of India. No serious attempts have been made on the reclamation technology for the reclamation of mine spoils in India. Hence there is an urging need to establish a suitable mycorrhizal technology for the reclamation of mine spoils in India. This paper gives an overview of the dominant

AM Fungal species and the mycotrophic plants of the mine spoils. A suitable reclamation technology may be developed with these plant species for the speedy revegetation of the mine spoils.

Distribution of AM fungi in various mine spoils

Nine species of VAM fungi belonging to the 3 major genera *viz.*, *Acaulospora*, *Glomus* and *Gigaspora* were identified from a rehabilitated coal mine spoil at Bisrampur, Madhya Pradesh (Dugaya et al., 1996). *Acaulospora*, *Glomus* and *Gigaspora* species were commonly found, while *Scutellospora* and *Sclerocystis* species were scarce in the coal mine overburden spoils at Kusmunda in Bilaspur district of Madhya Pradesh (Chandra and Jamaluddin, 1999).

VAM fungal infection varied from plant to plant among the thirty one plant

species, including legumes, grasses and sedges found growing naturally in the gypsum mine spoil in Rajasthan. The percent root infection was higher among the plants growing in mine spoil than those found in the normal soil (Tarafdar and Rao, 1997).

Only *Glomus globiferum* was present in the recently revegetated lignite mine spoil (1-2 years old) of Neyveli, while the adjacent undisturbed area possessed 23 VAM species (*Entrophospora colombiana*, *Gigaspora albida*, *Gi. gigantea*, *Gi. margarita*, *Glomus albidum*, *G. aggregatum*, *G. ambisporum*, *G. citricolum*, *G. claroidium*, *G. clarum*, *G. constrictum*, *G. deserticola*, *G. dimorphicum*, *G. etunicatum*, *G. fecundisporum*, *G. heterosporum*, *G. intraradices*, *G. macrocarpum*, *G. maculosum*, *G. mosseae*, *G. pustulatum*, *G. tenue*, *Sclerocystis coremioides*, *Sc. papillosa*, *Scutellospora aurigloba* and *S. pakistanica*). Only *G. aggregatum* and *G. pustulatum* were present in the oldest (25 years) disturbed mine spoil site (Ganesan et al., 1990).

Glomus was the most dominant genus isolated among the various mine spoils and mine spoil dumps in India (Ganesan et al., 1990; Dugaya et al., 1996; Selvam and Mahadevan, 2002; Mukhopadhyay and Maiti, 2010).

Glomus mosseae was commonly found in various mine spoils (Ganesan et al., 1990; Selvam and Mahadevan, 2002; Mukhopadhyay and Maiti, 2010; Singh and Jamaluddin; 2011). *G. aggregatum* was present in the coal, lignite and magnesite mine spoils, while *G. ambisporum* was common in the calcite, coal and lignite mine spoils. The other dominant species colonising the mine spoils in their decreasing order of magnitude were *G. deserticola*, *G. fasciculatum*, *G. heterosporum* and *G. intraradices*. The dominant status of the genus *Glomus* over the other genera might be due to the sporogenous characteristics of the fungi. Species of *Glomus* and *Acaulospora* produce small spores in a short time compared to the large spores produced by species of *Gigaspora*

and *Scutellospora* in the same environment (Nandakwang et al., 2008).

Entrophospora colombiana was the only species in the genus *Entrophospora* to be found in the mine spoils and dumps. It was commonly present among the coal and lignite mine spoils (Ganesan et al., 1990; Mehrotra, 1998).

Some of the other species commonly found in various mine spoils and dumps were *Acaulospora scrobiculata*, *Gigaspora gigantea*, *Gi. margarita*, *Sclerocystis microcarpus*, *Sc. pachycaulis*, *Scutellospora aurigloba*, *S. erythropo* and *S. Persica* (Table 1).

Most of the plants growing in the mine spoil possessed mycorrhizal association. The intensity of infection increased with the increasing age of mine spoils. The mycorrhizal inoculum needed for the infection seem to be available from the adjoining forest areas or rehabilitated sites. *Acacia auriculiformis*, *A. nilotica*, *Azadirachta indica*, *Dalbergia sissoo*, *Dendrocalamus strictus*, *Casuarina equisetifolia*, *Cassia siamea*, *Pithecellobium dulce* and *Gmelina arborea* were some of the dominant mycotrophic plants of the mine spoils in India (Table 2).

Most of these plants belong to the order leguminosae and possess nodules in their roots which help them in fixing atmospheric nitrogen. With the help of mycorrhizal association and root nodules, these plants were able to survive efficiently in the degraded soils. Plants belonging to the family Amaranthaceae and Cyperaceae (*Achyranthes aspera*, *Alternanthera sessilis*, *A. pungens*, *Celosia argentea* *Cyperus difformis* and *C. rotundus*) which were reported to be non-mycorrhizal, were found to be mycorrhizal. The low levels of phosphorus and nitrogen in the soil might have lead to the increased probability of infection in such plants as reported by Mosse and Phillips, 1971. However, the plants of the family Cleomaceae were found to be non-mycorrhizal, irrespective of the soil conditions.

Factors affecting mycorrhizal colonisation in mine spoils

The density and distribution of AM fungi depends on the age of the revegetated site, degree of surface soil disturbance, amount of homogeneous topsoil, and

presence of susceptible root material (Loree et al., 1987; Maiti, 1997; Singh and Jamaluddin, 2006), depth of the soil (Kabir et al., 1998; Xueli et al., 2002) and the physico-chemical nature of the soil.

Table-1: AM Fungi collected from various naturally revegetated mine spoils of India

Place of Collection	Acaulospora spp.,	Entrophospora spp.,	Gigaspora spp.,	Glomus spp.,	Sclerocystis spp.,	Scutellospora spp.,	Reference
Calcite mine spoil, Thazhaiyuthu, Tirunelveli, Tamil Nadu	—	—	—	<i>G. ambisporum</i> <i>G. claroidium</i> <i>G. deserticola</i> <i>G. dimorphicum</i> <i>G. heterosporum</i> <i>G. multicale</i> <i>G. pulvinatum</i> <i>G. pustulatum</i>	<i>Sc. pachycaulis</i> <i>Sc. sinuosa</i>	<i>S. gregaria</i>	Ganesan et al., 1990
Coal mine spoils, Kothagudem, Andhra Pradesh	<i>A. foveata</i>	<i>E. colombiana</i>	—	<i>G. aggregatum</i> <i>G. ambisporum</i> <i>G. botryoides</i> <i>G. citricoloum</i> <i>G. claroidium</i> <i>G. fasciculatum</i> <i>G. heterosporum</i> <i>G. mosseae</i>	<i>Sc. microcarpus</i> <i>Sc. rubiformis</i>	—	Ganesan et al., 1990
Coal mine site, Chandrapur, Maharashtra	<i>A. scrobiculata</i>	<i>E. colombiana</i>	—	<i>G. aggregatum</i> <i>G. ambisporum</i> <i>Glomus spp.,</i>	—	<i>S. calospora</i>	Mehrotra, 1998
Lignite mine spoil, Neyveli, Tamil Nadu	—	<i>E. colombiana</i>	<i>Gi. albida</i> <i>Gi. gigantea</i> <i>Gi. margarita</i>	<i>G. albidum</i> <i>G. aggregatum</i> <i>G. ambisporum</i> <i>G. citricoloum</i> <i>G. claroidium</i> <i>G. clarum</i> <i>G. constrictum</i> <i>G. deserticola</i> <i>G. dimorphicum</i> <i>G. etunicatum</i> <i>G. fecundisporum</i> <i>G. heterosporum</i> <i>G. intraradices</i> <i>G. macrocarpum</i> <i>G. maculosum</i> <i>G. mosseae</i> <i>G. pustulatum</i> <i>G. tenuie</i>	<i>Sc. coremioides</i> <i>Sc. papillosa</i>	<i>S. aurigloba</i> <i>S. pakistanica</i>	Ganesan et al., 1990
Overburden dumps of Neyveli Lignite Corporation, Neyveli, Tamil Nadu	<i>A. gerdemanni</i>	<i>E. colombiana</i>	<i>Gi. gigantea</i> <i>Gi. margarita</i>	<i>G. fasciculatum</i> <i>G. macrocarpum</i> <i>G. mosseae</i> <i>G. vesciculiformis</i>	<i>Sc. microcarpus</i> <i>Sc. pachycaulis</i>	<i>S. corolloidea</i> <i>S. erythropa</i> <i>S. persica</i> <i>S. verrucosa</i>	Selvam and Mahadevan, 2002
Magnesite mine spoil, Salem, Tamil Nadu	<i>A. bireticulata</i>	—	—	<i>G. aggregatum</i> <i>G. fasciculatum</i> <i>G. intraradices</i>	<i>Sc. microcarpus</i> <i>Sc. dussii</i> <i>Sc. pachycaulis</i>	<i>S. aurigloba</i> <i>S. erythropa</i> <i>S. nigra</i> <i>S. persica</i>	Raman et al., 1992
Limestone mine spoil, Odisha	<i>A. denticulata</i> <i>A. scrobiculata</i> <i>A. delicata</i>	—	<i>Gi. rosea</i> , <i>Gi. gigantea</i> , <i>Gi. margarita</i>	<i>G. arboreense</i> <i>G. deserticola</i> <i>G. fasciculatum</i> <i>G. intraradices</i> <i>G. mosseae</i>	—	<i>S. heterogama</i> <i>S. persica</i> <i>S. verrucosa</i>	Singh and Jamaluddin, 2011

Table-2: Mycorrhizal status of plants occurring in various revegetated mine spoils of India

Name of the plants	C	Co	L	Li	M	AM Colonisation	Percent root colonisation
Monocot plants							
Agavaceae	-	-	+	-	-	+	10
<i>Agave angustifolia</i> ¹							
Arecaceae							
<i>Cocos nucifera</i> ¹	-	-	+	-	-	Soil only	
<i>Phoenix sylvestris</i> ²	-	-	-	-	+	+	58
Cyperaceae							
<i>Cyperus difformis</i> ²	-	-	-	-	+	+	32
<i>C. rotundus</i> ⁵	-	+	-	-	-	+	45
Poaceae							
<i>Bambusa arundinacea</i> ¹	+	-	-	-	-	+	80
<i>Chloris barbata</i> ¹	+	-	+	-	-	+	10
<i>Chrysopogon zeylanicus</i> ¹	+	-	+	-	-	+	40
<i>Cynodon dactylon</i> ^{1,5}	-	+	+	-	-	+	0,50
<i>Dactyloctenium aegyptium</i> ⁵	-	+	-	-	-	+	60
<i>Desmodium triflorum</i> ⁵	-	+	-	-	-	+	48
<i>Dendrocalamus strictus</i> ^{1,5,6}	-	+	+	-	-	+	0,30,55
<i>Dichanthium annulatum</i> ⁵	-	+	-	-	-	+	65
<i>Digitaria bicornis</i> ⁵	-	+	-	-	-	+	57
<i>Eragrostis tenella</i> ⁵	-	+	-	-	-	+	40
<i>E. unioloides</i> ⁵	-	+	-	-	-	+	50
<i>Eragrostis spp.,</i> ¹	-	-	+	-	-	+	20
<i>Grevillea pteridofolia</i> ⁵	-	+	-	-	-	+	30
<i>Heteropogon contortus</i> ⁵	-	+	-	-	-	+	30
<i>Phragmites karka</i> ¹	-	-	+	-	-	+	0
<i>Saccharum arundinaceum</i> ⁵	-	+	-	-	-	+	70
<i>S. spontaneum</i> ⁵	-	+	-	-	-	+	55
<i>Sporobolus indicus</i> ⁵	-	+	-	-	-	+	20
Dicot plants							
Amarantaceae							
<i>Achyranthes aspera</i> ^{1,5}	-	+	-	-	-	-	0,45
<i>Aerva lanata</i> ¹	+	-	-	-	-	-	0
<i>Alternanthera sessilis</i> ⁵	-	+	-	-	-	+	42
<i>A. pungens</i> ⁵	-	+	-	-	-	+	38
<i>Celosia argentea</i> ⁵	-	+	-	-	-	+	25
Anacardiaceae							
<i>Anacardium occidentale</i> ^{1,5}	-	+	+	-	-	+	0,20
Annonaceae							
<i>Annona squamosa</i> ²	-	-	-	-	+	+	56
Apocynaceae							
<i>Holarrhena antidysentrica</i> ⁵	-	+	-	-	-	+	65
Asclepiadaceae							
<i>Calotropis gigantea</i> ²	-	-	-	-	+	+	62
<i>C. procera</i> ⁵	-	+	-	-	-	+	60
Asteraceae							
<i>Ageratum conyzoides</i> ⁵	-	+	-	-	-	+	50
<i>Bidens biternata</i> ⁵	-	+	-	-	-	+	40
<i>Blumea spp.</i> ⁵	-	+	-	-	-	-	0
<i>Eclipta prostrata</i> ⁵	-	+	-	-	-	+	52
<i>Parthenium hysterophorus</i> ^{5,7}	-	+	-	-	-	+	20,20
<i>Tridax procumbens</i> ^{5,7}	-	+	-	-	-	+	45,40
<i>Vicoa indica</i> ¹	-	+	-	-	-	+	10
Capparidaceae							
<i>Cleome aspera</i> ¹	-	+	-	-	-	-	0
<i>C. chelidonii</i> ¹	+	-	-	-	-	-	0
<i>C. viscosa</i> ¹	-	+	-	-	-	-	0

Casuarinaceae							
<i>Casuarina equisetifolia</i> ^{1,2,5}	-	+	+	-	+	+	40,78,40
Ceasalpiniaceae							
<i>Cassia auriculata</i> ^{1,2}	+	-	-	-	+	+	20,56
<i>C. fistula</i> ⁵	-	+	-	-	-	+	45
<i>C. occidentalis</i> ¹	-	+	-	-	-	+	10
<i>C. siamea</i> ^{1,4,5,6}	-	+	+	-	-	+	60,‡,73,90
<i>C. tora</i> ^{1,5}	-	+	-	-	-	+	45
<i>Caesalpinia pulcherrima</i> ¹	+	-	-	-	-	+	50
<i>Delonix regia</i> ¹	+	-	+	-	-	+	80
<i>Peltophorum pterocarpum</i> ⁴	-	+	-	-	-	+	‡
<i>Tamarindus indica</i> ^{1,5}	+	+	-	-	-	+	45,50
Combretaceae							
<i>Terminalia arjuna</i> ^{5,6}	-	+	-	-	-	+	60,58
<i>T. bellerica</i> ⁵	-	+	-	-	-	+	50
Convolvulaceae							
<i>Evolvulus alsinoides</i> ⁵	-	+	-	-	-	+	60
<i>Ipomoea carnea</i> ²	-	-	-	-	+	+	48
<i>I.hispida</i> ⁵	-	+	-	-	-	-	0
<i>Merremia tridentata</i> ⁵	-	+	-	-	-	+	56
Cucurbitaceae							
<i>Citrullus colocynthis</i> ¹	+	-	-	-	-	+	80
<i>Mukia maderaspatana</i> ¹	+	-	-	-	-	+	60
Dipterocarpaceae							
<i>Shorea robusta</i> ⁵	-	+	-	-	-	+	30
Ebenaceae							
<i>Diospyros melanoxylon</i> ⁵	-	+	-	-	-	+	40
Euphorbiaceae							
<i>Croton bonplandianus</i> ¹	-	+	-	-	-	+	85
<i>Euphorbia hirta</i> ^{1,5}	-	+	-	-	-	+	72,48
<i>Phyllanthus amarus</i> ¹	-	+	-	-	-	+	40
<i>P. emblica</i> ⁵	-	+	-	-	-	+	40
<i>P. niruri</i> ⁷	-	+	-	-	-	+	30
<i>P. virgatus</i> ^{1,5}	-	+	-	-	-	+	75,60
Lamiaceae							
<i>Leucas plukenetii</i> ⁵	-	+	-	-	-	+	42
<i>Ocimum basilicum</i> ⁵	-	+	-	-	-	+	55
<i>O. gratissimum</i> ⁷	-	+	-	-	-	+	30
<i>Plectranthus caninus</i> ²	-	-	-	-	+	+	82
Lythraceae							
<i>Woodfordia fruticosa</i> ⁵	-	+	-	-	-	-	0
Malvaceae							
<i>Sida acuta</i> ¹	-	+	-	-	-	+	60
<i>S. cordata</i> ^{1,5}	-	+	-	-	-	-	0
Meliaceae							
<i>Azadirachta indica</i> ^{1,2,5}	-	+	+	-	+	+	40,64,70
<i>Melia azedarach</i> ⁵	-	+	-	-	-	+	30
Mimosaceae							
<i>Acacia auriculiformis</i> ^{1,2,4,5,6}	-	+	+	-	+	+	10,20,‡,68,89
<i>A. catechu</i> ^{4,5}	-	+	-	-	-	+	‡,43
<i>A. mearnsii</i> ²	-	-	-	-	+	+	64
<i>A. niloticas</i> ^{1,4,5,7}	-	+	+	-	-	+	0,‡,55,20
<i>Acacia spp.,³</i>	-	+	-	-	-	-	‡
<i>Albizia lebbeck</i> ¹	-	+	+	-	-	+	90
<i>A. procera</i> ⁴	-	+	-	-	-	+	‡
<i>Albizia spp.,³</i>	-	+	-	-	-	-	†
<i>Leuceana leucocephala</i> ⁶	-	-	+	-	-	+	68
<i>Pithecellobium dulce</i> ^{1,3,4}	-	+	+	-	-	+	0,‡,‡
<i>Prosopis juliflora</i> ⁵	-	+	-	-	-	+	30
<i>P. spicigera</i> ¹	-	-	+	-	-	-	-

Moraceae	-	+	-	-	-	+	65
<i>Ficus benghalensis</i> ⁵	-	+	-	-	-	+	30
<i>F. elastica</i> ⁵	-	+	-	-	-	+	20
<i>F. religiosa</i> ⁵	-	+	-	-	-	+	60
<i>Morus alba</i> ⁵							
Myrtaceae							
<i>Eucalyptus globulus</i> ^{1,2}	-	-	+	-	+	+	0,62
<i>E. hybrida</i> ^{4,5}	-	+	-	-	-	+	‡,40
<i>Syzygium cumini</i> ⁵	-	+	-	-	-	+	52
Nyctaginaceae							
<i>Boerhaavia diffusa</i> ¹	-	+	-	-	-	-	0
Oxalidaceae							
<i>Biophytum sensitivum</i> ¹	+	-	-	-	-	+	70
Papavaraceae							
<i>Argemone mexicana</i> ^{5,7}	-	+	-	-	-	-	35,40
Papilionaceae							
<i>Alysicarpus vaginalis</i> ¹	-	+	-	-	-	+	85
<i>Atylosia scarabaeoides</i> ⁵	-	+	-	-	-	+	20
<i>Butea monosperma</i> ⁵	-	+	-	-	-	+	50
<i>Clitoria</i> sp. ⁵	-	+	-	-	-	+	30
<i>Crotalaria albida</i> ⁵	-	+	-	-	-	+	10
<i>Dalbergia sissoo</i> ^{1,3,4,5,6}	+	+	+	-	-	+	15,‡,‡,58,94
<i>Indigofera linnifolia</i> ⁵	-	+	-	-	-	+	70
<i>Pongamia glabra</i> ⁵	-	+	-	-	-	+	40
<i>P. pinnata</i> ^{2,3}	-	+	-	-	+	+	48
<i>Tephrosia purpurea</i> ¹	-	+	-	-	-	+	35
<i>Zornia gibbosa</i> ⁵	-	+	-	-	-	+	58
Portulaceae							
<i>Portulaca pilosa</i> ¹	-	-	-	-	-	-	0
Proteaceae							
<i>Grevillea robusta</i> ¹	-	-	+	-	-	Soil only	
Rhamnaceae							
<i>Zizyphus glaberrima</i> ⁵	-	+	-	-	+	-	30
<i>Z. mauritiana</i> ⁷	-	-	-	-	+	+	20
Rubiaceae							
<i>Ixora pavetta</i> ²	-	-	-	-	+	+	52
<i>Spermococe hispida</i> ¹	-	+	-	-	-	+	50
Rutaceae							
<i>Aegle marmelos</i> ⁵	-	+	-	-	+	-	40
Sapotaceae							
<i>Madhuca indica</i> ⁵	-	+	-	-	+	-	56
Scrophulariaceae							
<i>Hedyotis affinis</i> ⁵	-	+	-	-	+	-	35
<i>Scoparia dulcis</i> ⁵	-	+	-	-	+	-	0
Simarubaceae							
<i>Ailanthus excelsa</i> ⁴	-	+	-	-	-	+	‡
Solanaceae							
<i>Datura metel</i> ¹	+	-	-	-	-	+	8
<i>Solanum xanthocarpum</i> ¹	+	-	-	-	-	+	8
Typhaceae							
<i>Typha angustata</i> ¹	+	-	-	-	-	+	50
Ulmaceae							
<i>Holoptelia integrifolia</i> ⁵	-	+	-	-	+	+	55
Verbenaceae							
<i>Gmelina arborea</i> ^{3,4,6}	-	+	-	-	-	+	‡,‡,38
<i>Tectona grandis</i> ¹	-	+	+	-	-	-	68,0

C-Calcite mine spoil; Co- Coal mine spoil; L-Lignite mine spoil; Li-Lime stone mine spoil;

M-Magnesite mine spoil; AM- Arbuscular Mycorrhiza. + Presence; - Absence; ‡ Not known.

¹-Ganesan *et al.*, (1990); ²- Raman *et al.*, (1992); ³- Dugaya *et al.*, (1996); ⁴- Chandra and Jamaluddin (1999);

⁵- Kumar *et al.*, (2003); ⁶-Mukhopadhyay and Maiti (2010); ⁷-Singh and Jamaluddin (2011).

Conclusion

Naturally revegetated mine spoils had lower level of mycorrhizal colonisation compared to the artificially reclaimed mine spoils. This may be due to the inoculation of AM spores to the planted species which promotes the colonisers for establishment (Singh and Jamaluddin, 2011). Enriching the soil with AM fungi by planting of legumes can be the best approach for the reclamation of mine spoils and mine spoil dumps.

References

- Chandra KK, Jamaluddin. 1999. Distribution of vesicular arbuscular mycorrhizal fungi in coal mine overburden dumps. *Ind. Phytopathol.* 52 (3): 254-258.
- Dugaya D, Williams AJ, Chandra KK, Gupta BN, Banerjee SK. 1996. Mycorrhizal development and plant growth in amended coal mine overburden. *Indian Journal of Forestry* 19(3): 222-226.
- Ganesan V, Ragupathy S, Parthipan B, Rajini Rani DB, Mahadevan A. 1990. Distribution of vesicular-arbuscular mycorrhizal fungi in coal, lignite, and calcite mine spoils of India. *Biol Fertil Soils*. 12: 131-136.
- Kabir Z, O'Halloran IP, Widden P, Hamel C. 1998. Vertical distribution of arbuscular mycorrhizal fungi under corn (*Zea mays L.*) in no-till and conventional tillage systems. *Mycorrhiza* 8(1): 53-55.
- Kumar A, Raghuwanshi R, Upadhyay RS. 2003. Vesicular arbuscular mycorrhizal association in naturally revegetated coal mine spoil. *Tropical Ecology* 44(2): 253-256.
- Loree MAG, Williams SE. 1987. Colonisation of westerngrass (*Agropyron smithii* Rydb.) by vesicular arbuscular mycorrhizal fungi during the revegetation of a surface mine. *New Phytol.* 106: 725-743.
- Mehrotra VS. 1998. Arbuscular mycorrhizal associations of plants colonizing coal mine spoil in India. *J. Agri. Sci.* 130(2): 125-133.
- Mosse B, Phillips JM. 1971. The influence of phosphate and other nutrients on the development of vesicular-arbuscular mycorrhiza in culture. *J. Gen. Microbiol.* 69: 157.
- Mukhopadhyay S, Maiti SK. 2009. Biofertiliser: VAM fungi-Future prospect for biological reclamation of mine degraded lands. *Ind. J. Environ. Protect.* 29(9): 801-808.
- Mukhopadhyay S, Maiti SK. 2010. Natural mycorrhizal colonization in tree species growing on the reclaimed coalmine overburden dumps: Case study from Jharia coalfields, India. *The Bioscan* 3: 761-770.
- Nandakwong P, Elliot S, Lumyang S. 2008. Diversity of arbuscular mycorrhizal fungi in forest restoration area of Doe Suthep-Pui National Park, North Thailand. *J. Microsc Soc Thai*. 22: 60-64.
- Raman N, Nagarajan N, Gopinathan S, Sambandan K. 1992. Mycorrhizal status of plant species colonizing a magnesite mine spoil in India. *Biol Fertil Soils*. 16: 76-78.
- Rao AV, Tarafdar JC. 1998. Selection of plant species for rehabilitation of gypsum mine spoil in arid zone. *J. Arid Env.* 39(4): 559-567.
- Selvam A, Mahadevan A. 2002. Distribution of mycorrhizas in an abandoned fly ash pond and mined sites of Neyveli Lignite Corporation, Tamil Nadu, India. *Basic and Applied Ecology* 3(3): 277-284.
- Singh A, Jamaluddin. 2006. Multiplication and trapping of vesicular arbuscular mycorrhiza fungi in soil of dumps of limestone quarries. *Mycorrhiza News* 17(4): 17-19.
- Singh AK, Jamaluddin. 2011. Status and diversity of arbuscular mycorrhizal fungi and its role in natural regeneration on limestone mined spoils. *Biodiversitas* 12(2): 107-111.
- Xueli He, Stanislav M, Steinberger Y. 2002. Spatial Distribution and Colonization of Arbuscular Mycorrhizal Fungi under the Canopies of Desert Halophytes. *Arid Land Research and Management* 16(2): 149-160.