

Review article

## Genetics of salt tolerance in rice

P. Deepa Sankar\*

Plant Biotechnology Division, School of Bio Sciences and Technology, VIT University, India

\*Corresponding author email: pdeepasankar@vit.ac.in

**Experimental procedures viz., line x tester analysis, diallel analysis and stability analysis have been utilized in rice breeding programs to study combining ability, heterosis, gene action and stress tolerance capacity of various genotypes based on yield related traits. The results have helped in development of good varieties and hybrids. Excessive work is available in the particular area of research as many plant breeders in Asia work towards improving the yield of rice. A review on gene action studies in rice has been attempted with emphasis on the research progress made in India.**

**Key Words:** Rice, gene action, additive, non additive

Rice is synonymous to food in Asia, a life giving cereal, highly respected and considered auspicious by many in the world. It is also well known for its easy digestibility, taste and nutritive value. Rice is preferred differentially based on grain quality and the nature of processing (Deepa Sankar, 2015). Widespread salinity is a major factor in world agriculture that limits crop productivity. Rice is an important crop and experimental evidence suggests that most rice varieties exhibit salt sensitivity and varietal differences tend to manifest only at moderate salt concentration, that is 50mM NaCl (Yeo and Flowers, 1984).

### **Nature of yield related traits under salt stress**

The *in situ* performance of a variety and its reduction in grain yield at defined stress levels can be followed in screening and evaluating rice breeding materials and varieties. A high percentage of spikelet sterility relates to a low level of salt tolerance in rice. Low Na<sup>+</sup>/K<sup>+</sup> ratio of ion uptake is positively correlated with a high level of salt

tolerance and can be taken into consideration as a desired characteristic while screening rice lines (Mishra *et al.*, 1997). Early studies reported that salt injury in rice was caused by both osmotic imbalance and an accumulation of the chloride ion (Akbar, 1975). Besides osmotic imbalance and ion toxicity, nutritional imbalance causes salt injury in rice. The symptoms of salt injury were observed to be stunted growth, rolling and withering of leaves, occurrence of white blotches, browning and drying of older leaves (Tagawa and Ishizaka, 1963; Akbar, 1975; Akbar, 1986).

Rice has been found to be tolerant during germination, vegetative growth and maturity but very sensitive at early seedling growth (2-3 leaf stage), pollination and fertilization stages (Pearson *et al.*, 1966; Akbar and Ponnampereuma, 1982). Krishnamurthy *et al.* (1971) observed that tillering was not influenced by salinity. Karim and Haque (1986) reported that plant height, root length, number of tillers per plant, straw yield and dry weight of root was affected by salinity during vegetative stage. Yield reduction at

maximum tillering stage due to salinity was observed by Ahmed *et al.* (1989). Decreasing

grain yield with increase in salinity was reported by Powar and Mehta (1997).

**Table 1. Nature of yield attributing genes**

| Character                                   | Gene action  | Author and year   |
|---|--|---|
| <b>Days to 50 per cent flowering</b>        | Additive<br>Non additive<br>Both additive and non additive         | Lakshmi Narayanan (2000)<br>Rogbell and Subbaraman (1997); Ghara <i>et al.</i> (2012)<br>Narayanan and Rangasamy (1991)   |
| <b>Plant Height</b>                         | Additive<br><br>Non additive<br><br>Both additive and non additive | Akbar <i>et al.</i> (1985); Mosina (1986); Rogbell (1995);<br>Lakshmi Narayanan (2000); Babu (2002)<br>Thirumeni <i>et al.</i> (2003); Malini <i>et al.</i> (2006); Ghara <i>et al.</i> (2012)<br>Narayanan and Rangasamy (1991); Rahimi <i>et al.</i> (2010) |
| <b>Productive tillers per plant</b>         | Additive<br>Non additive<br>Both additive and non additive         | Bhattacharya (1978); Babu (2002)<br>Rogbell (1995)<br>Narayanan and Rangasamy (1991)  |
| <b>Panicle length</b>                       | Additive<br>Non additive<br><br>Both additive and non additive     | Thirumeni <i>et al.</i> (2000) ; Lakshmi Narayanan (2000)<br>Thirumeni <i>et al.</i> (2003); Malini <i>et al.</i> (2006); Ghara <i>et al.</i> (2012)<br>Narayanan and Rangasamy (1991); Rahimi <i>et al.</i> (2010)   |
| <b>Grains per panicle</b>                   | Additive<br>Non additive<br><br>Both additive and non additive     | Thirumeni <i>et al.</i> (2000); Malini <i>et al.</i> (2006)<br>Rogbell and Subbaraman (1997); Ghara <i>et al.</i> (2012)<br>Narayanan and Rangasamy (1991)<br>Rahimi <i>et al.</i> (2010)   |
| <b>Spikelet fertility</b>                   | Additive<br>Non additive   | Bhattacharya (1978)<br>Akbar and Yabuno (1974); Babu (2002)<br>Thirumeni <i>et al.</i> (2003); Ghara <i>et al.</i> (2012)   |
| <b>100 grain weight</b>                     | Additive<br>Non additive<br>Both additive and non additive         | Ba Tao <i>et al.</i> (1992)<br>Lakshmi Narayanan (2000)<br>Narayanan and Rangasamy (1991)   |
| <b>Single plant yield</b>                   | Additive<br><br>Non additive<br>Both additive and non additive     | Akbar <i>et al.</i> (1985); Narayanan and Rangasamy (1991);<br>Choung <i>et al.</i> (1991)<br>Rogbell and Subbaraman (1997)<br>Yoshida <i>et al.</i> (1976); Mishra (1995)  |
| <b>Na<sup>+</sup> : K<sup>+</sup> ratio</b> | Additive<br>Non additive<br>Both additive and non additive         | Babu (2002)<br>Thirumeni (1998)<br>Gregario and Senadhira (1993); Mishra (1995); Mishra <i>et al.</i> (1996); Mishra <i>et al.</i> (2003)   |
| <b>Salinity tolerance</b>                   | Additive<br>Non additive<br>Both additive and non additive         | Jones (1986); Mishra <i>et al.</i> (2003)<br>Moeljopawiro and Ikehashi (1981); Lavrichenko (1985)<br>Akbar <i>et al.</i> (1985) ; Mosina (1986) ; Gregario and Senadhira (1993) ; Mishra <i>et al.</i> (1996)   |

For transplanted rice, initial survival is crucial in determining later performance. Seedlings that survive at this stage are expected to produce grain although spike sterility cannot be ruled out (Ansari *et al.*, 2003). Seedling survival after planting could, to some extent, be compensated for by using old seedlings (Ansari *et al.*, 1994), but initial seedling survival also reflects upon the performance of a particular line in terms of yield. In a study performed by Ansari *et al.* (2003) cultivars Ganga White, Nona Bokra, IR 6 and Shua-92 had better seedling survival, whereas IR 28 and Basmati had the lowest seedling survival. These cultivars had a more or less similar ranking in straw or grain weight.

### Gene action studies

Gene action refers to the behavior or mode of expression of genes in a genetic population. Success in breeding for quantitative traits depends upon the gene action involved for the traits concerned. Baker (1978) pointed out that the relative importance of general and specific combining ability variances could be used for predictability of gene action. Nature of gene action of various yield related traits is useful to the plant breeders in formulating the breeding procedure for obtaining maximum genetic advance in yield and yield components. Works on rice regarding nature of gene action are presented in Table 1.

### Gene action studies involving two line hybrids

Rice hybrids have already contributed towards food security in China. India has also made remarkable progress towards development and use of hybrid rice technology. But the excessive dependence on a single source of cytoplasmic male sterility (CMS) *via* WA and the cumbersome process of seed production and parental line development warrant the development of

alternate approaches to exploit hybrid vigour in rice. Two line breeding is one such possibility that emerged following the chance discovery of a photoperiod-sensitive genic male sterile plant called Nongken 58 by Prof. Shi Ming Song of China (Shi and Deng, 1986) which was found to be sterile under longer photoperiods (>14 hr) and fertile under shorter photoperiods (<13 hr) subsequently. Temperature sensitive genic male sterility (TGMS) lines were identified by Chinese and Japanese scientists. They are sterile under high temperature (>32°C) but under low temperature (< 24°C) they are fertile (Tan *et al.*, 1990; Maruyama *et al.*, 1990).

World will have to produce 60% more rice by 2030 than what it produced in 1995. Theoretically rice has great yield potential to be tapped and there are many ways to raise rice yield. Among them it seems at present that the most effective and economic way available is to develop hybrid varieties based on successful experience in China. Following the success of three-line hybrid rice in 1970's, two-line hybrid rice was successfully commercialized in 1995 with a yield advantage of 5-10 % higher than that of existing three-line hybrid rice (Yuan Longping, 2004). Hence it becomes really important to analyze genetic studies made on two line rice hybrids as well.

Chen *et al.* (1997) studied combining ability of agronomic traits in two line intersubspecific hybrid rice. They reported that yield per plant appeared to be mainly controlled by non-additive effects while plant height and heading date were mainly conditioned by additive gene effects.

A combining ability study was made to assess the nature of gene action for seven quantitative traits using three TGMS lines (Lines) and six non TGMS lines (Testers) in a L x T mating design (Rangaswamy *et al.*, 1998). TS-15 showed significant positive *gca* effects for number of filled grains per panicle and spikelet fertility. TS-16 was a good

general combiner for plant height, productive tillers per plant and panicle length. TS-18 recorded significant *gca* effect for number of filled grains per panicle, spikelet fertility and grain yield per plant. Among the testers, ADT 36 and IR-37399 recorded significant positive *gca* effects for grain yield per plant. Seven hybrids *viz.*, TS-15 x ASD-18, TS-15 x IR-55722, TS-15 x IR-59601, TS-16 x ASD-18, TS-16 x ADT-36, TS-18 x ADT-36, and TS-16 x IR-68902B exhibited significant positive *sca* effects for grain yield.

Sampoornam and Thiyagarajan (1998) studied combining ability for grain yield and related traits in two line hybrids and found that TS 18 was the good general combiner for many traits.

Aananthi (2000) studied combining ability of TGMS lines and reported the predominance of additive gene action for the traits productive tillers per plant, panicle length, plant height, 100 grain weight and single plant yield. Among the TGMS lines TS 29 and TS 6 were reported as good combiners.

A study was undertaken by Bini Philip (2001) using 21 TGMS based rice hybrids obtained by crossing three TGMS lines with seven testers to evaluate gene action in a line x tester design. Predominance of non additive gene action was observed for most of the traits except for single plant yield, chlorophyll content and plant height. Harvest index showed both GCA and SCA variance in equal magnitude.

Radhidevi *et al.* (2002) performed combining ability analysis using two CMS, two TGMS lines, 12 testers and 48 hybrids in Line x Tester fashion. The greater role of non additive gene action was observed for most of the characters except days to 50 per cent flowering and panicle length which offers much scope for the exploitation of hybrid vigour through heterosis breeding.

Kalaiyarasi *et al.* (2002) reported the preponderance of non additive gene action in terms of yield components from a study

based on two line inter-subspecific rice hybrids.

Patil *et al.* (2003) studied combining ability by making 84 cross combinations from seven TGMS lines and 12 non-TGMS testers. The magnitude of specific combining ability variance was greater than the general combining ability for all the traits except days to 50 per cent flowering. Combining ability revealed the predominance of non additive gene action for productive tillers per plant, panicle length, grains per panicle and grain yield per plant and offered more scope for exploitation of hybrid vigour through heterosis breeding.

Gnanasekaran (2003) performed genetic analysis involving TGMS based rice hybrids. He reported that the traits days to 50 per cent flowering, plant height, number of productive tillers per plant, panicle exertion, 100 grain weight and kernel L/B ratio were governed by additive gene action while panicle length, number of grains per panicle, spikelet fertility, harvest index, single plant yield, kernel elongation and amylase content were controlled by non additive gene action.

Deepa Sankar *et al.* (2008) have reported that the traits days to 50 per cent flowering, productive tillers per plant, panicle exertion, panicle length, panicle weight, filled grains per panicle, spikelet fertility, 100 grain weight, Na<sup>+</sup> : K<sup>+</sup> ratio, harvest index and grain yield per plant are controlled by dominance gene action and plant height was controlled by additive gene action. The study was undertaken to get genetic information on salt tolerance in a set of rice genotypes comprising of five TGMS lines, eight salt tolerant testers and 40 hybrids obtained by crossing in line x tester design.

## Conclusion

Conventional breeding approaches *viz.*, recombination and heterosis breeding are the real problem solvers which have brought in food security. Gene action studies reveal good combiners for specific breeding

purposes. Though tissue culture, molecular and biotechnological methods employed aid in improving yield, quality and tolerance to stresses the main backbone of plant breeding has always been the conventional techniques employed. Efforts of breeders in creating such wonderful genotypes have led to the perpetuation of the human race.

## References

- Aananthi N. Genetic analysis in two line rice (*Oryza sativa* L.) hybrids [dissertation]. Coimbatore (TN): Tamil Nadu Agricultural University; 2000.
- Ahmed IV, Faiz SMA, Anwar AKM, Satter A. 1989. Relation of Na<sup>+</sup>: K<sup>+</sup> ratio of rice with salinity and yield. *Curr. Agric.* 13: 73-78.
- Akbar M. 1975. Water and chloride absorption in rice seedlings. *J. Agric. Res.* 13: 342-343.
- Akbar, M. 1986. Breeding for salinity resistance in rice. In : *Prospects for bio-saline research*, Eds. R. Ahmed and AS. Pietro, Pakistan: Department of Botany, University of Karachi, pp. 37-55.
- Akbar, M., and Ponnampuruma, FN. 1982. Saline soils of south and South East Asia as potential rice lands. In *Rice research strategies for the future*, Manila, Philippines: IRRI, pp. 265-281.
- Akbar M, Yabuno T. 1974. Breeding for saline resistant varieties of rice. Inheritance of delayed type panicle sterility induced by salinity. *Japan J. Breed.* 27: 237-240.
- Akbar, M., Khush, GS., and Hillerislambers, D. 1985. Genetics of salt tolerance in rice. In *Rice genetics*, Proceedings of the International Rice genetics symposium, Manila, Philippines: IRRI, pp. 399-410.
- Ansari, R., Khanzada, AN., and Naqvi, SSM. 1994. Studies on salt tolerance in rice. In *Current developments in salinity and drought tolerance in plants*, Eds. SSM. Naqvi, R. Ansari, TJ. Flowers and AR. Azmi, Tandojam, Pakistan: AEARC, pp 207-239.
- Ansari, R., Shereen, A., Alam, SM., Flowers, TJ., and Yeo, AR. 2003. Breeding for salt tolerance in rice. In *Advances in Rice Genetics. Supplement to rice Genetics*, Eds. GS. Khush, DS. Brar and B. Hardy, IV Proceedings of the Fourth International Rice Genetics Symposium, 22<sup>nd</sup> -27<sup>th</sup> October 2000, Los Banos, Philippines: IRRI, pp. 8-10.
- Ba Tao P, Truong DX, Buu BC. 1992. Assessment of breeding materials from salt tolerant deepwater rices in the Mekong Delta. *Nghien Cuu Lua. Omonrice.* 2: 56-63.
- Babu S. Studies on response of salt tolerant rice (*Oryza sativa* L.) hybrids over environments and *in vitro* conditions [dissertation]. Coimbatore (TN): Tamil Nadu Agricultural University; 2002.
- Baker RJ. 1978. Issue in diallel analysis. *Crop Sci.* 18: 533-536.
- Bhattacharya RK. 1978. Estimates of genetic parameter of some quantitative character in rice under saline soil conditions. *Oryza.* 15(2): 146-150.
- Bini Philip. Genetic analysis of Thermosensitive Genic Male Sterility (TGMS) based rice (*Oryza sativa* L.) hybrids [dissertation]. Coimbatore (TN): Tamil Nadu Agricultural University; 2001.
- Chen SH, Lu HR, Yang JB, Hu RY, Zhao MF. 1997. Heterosis, combining ability and correlation of agronomic traits in two line inter subspecific hybrid rice. *J. Fujian Agric. Univ.* 26(1): 1-7.
- Choung JI, Lee MS, Park SH, Yu SJ. 1991. Genetic analysis of major agronomic characters related to salt tolerance in rice. 1. Test of heterosis and combining ability of yield characters by diallel analysis. *Research reports of the rural development administration. Rice.* 33(3): 26-36.
- Deepa Sankar P. 2015. *In vitro* screening for salt tolerance in rice. *Research Journal of Pharmaceutical, Biological and Chemical Sciences.* 6(4):1694-1699.
- Deepa Sankar P, Subbaraman N, Lakshmi Narayanan S. 2008. Heterosis, combining

- ability and gene action studies in TGMS based rice hybrids under normal and salt affected environments. *Indian J. Agric. Res.* 42(3):177-182.
- Ghara AG, Nematzadeh G, Bagheri N, Ebrahimi A Oladi M. 2012. Evaluation of general and specific combining ability in parental lines of hybrid rice. *International Journal of Agriculture: Research and Review.* 2(4): 455-460.
- Gnanasekaran M. Genetic analysis involving TGMS based rice (*Oryza sativa* L.) hybrids [dissertation]. Coimbatore (TN): Tamil Nadu Agricultural University; 2003.
- Gregario GB Senadhira D. 1993. Genetic analysis of salinity tolerance in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* 86: 333-338.
- Jones, M.P. 1986. Genetic analysis of salt tolerance in mangrove swamp rice. In *Rice genetics*, Proceedings of the International Rice genetics symposium, 27<sup>th</sup>-31<sup>st</sup> May, 1985, Manila, Philippines: IRRI. pp. 411-422.
- Kalaiyarasi R, Palanisamy GA, Vaidyanathan P. 2002. The potentials and scope of utilizing TGMS lines in inter-subspecies crosses of rice (*Oryza sativa* L.). *J. Genet. Breed.* 56(2): 137-143.
- Karim NH, Haque MZ. 1986. Salinity tolerance at the reproductive stage of five rice varieties. *Bangladesh J. of Agric.* 11(4): 73-76.
- Krishnamurthy R, Shastry SVS, Freeman WH. 1971. Breeding rice for tolerance to drought and salinity. *Oryza.* 8: 47-56.
- Lakshmi Narayanan S. Genetic analysis and in vitro studies of salt tolerance in rice (*Oryza sativa* L.) [dissertation]. Coimbatore (TN): Tamil Nadu Agricultural University; 2000.
- Lavrichenko VG. 1985. Inheritance of salt resistance in rice hybrids for the first two generations. *Rice Abst.* 11: 31.
- Malini N, Sundaram T, Ramakrishnan S, Saravanan S. 2006. Prediction of hybrid vigour for yield attributes among synthesized hybrids in rice. *Research Journal of Agricultural and Biological Sciences.* 2: 166-170.
- Maruyama, K., Araki, H., and Kato, H. 1990. Thermosensitive genic male sterility induced irradiation. In *Rice Genetics II*, Manila, Philippines: IRRI, pp. 227-235.
- Mishra, B. 1995. Breeding for salt tolerance in crops. In *Annual Report of 1995*, Karnal, India: Central Soil Salinity Research Centre.
- Mishra B, Akbar M, Seshu, DV, Senadhira D. 1996. Genetics of salinity tolerance and ionic uptake in rice. *IRRN.* 21(1): 38-39.
- Mishra, B., Singh, RK., and D. Senadhira. 2003. Advances in breeding salt tolerant rice varieties. In *Advances in Rice Genetics. Supplement to rice Genetics IV*, Eds. G.S Khush, D.S. Brar and B. Hardy. Proceedings of the Fourth International Rice Genetics Symposium, 22-27<sup>th</sup> October 2000, Los Banos, Philippines: IRRI. pp. 5-7.
- Mishra, B., Singh, R.K., and Senadhira, D. 1997. Enhancing genetic resources and breeding for problem soils. In *Using Diversity: Enhancing and Maintaining Genetic Resources On-Farm*. Eds. Louise Sperling and Michael Loevinsohn. Proceedings of a workshop held on 19-21<sup>st</sup> June 1995, New Delhi, India: International Development Research Centre. pp. 216-229.
- Moeljopawiro S, Ikehashi H. 1981. Inheritance of salt tolerance in rice. *Euphytica.* 30: 291-300.
- Mosina SB. 1986. Breeding rice for salt resistance. *Truly.* 266/294: 5-15.
- Narayanan, KK., and Rangasamy, S.R. 1991. Genetic analysis for salt tolerance in rice. In *Rice genetics II*. Proceedings of the second international rice genetics symposium, 14-18<sup>th</sup> May 1990. Manila, Philippines: IRRI. pp. 167-193.
- Patil DV, Thiyagarajan K, Kamble, P Kamble P. 2003. Combining ability of parents for yield and yield contributing traits in two

- line hybrid rice (*Oryza sativa* L.) Crop Res. 25(3): 520-524.
- Pearson GA, Ayers AD, Ebehard DC. 1966. Relative salt tolerance of rice during germination and early seedling development. Soil Sci. 102: 151-156.
- Powar SL, Mehta VB. 1997. Salt tolerance of different rice varieties under coastal saline condition. Ann. Agric. Res. 18(4): 536-537.
- Radhidevi RP, Nagarajan P, Shanmugasundaram P, Babu RC, Jayanthi S, Subramani S. 2002. Combining ability analysis in three line and two line rice hybrids. Plant Archives. 2(1): 99-102.
- Rahimi M, Rabiei B, Samizadeh H, Ghasemi AH. 2010. Combining ability and heterosis in rice (*Oryza sativa* L.). J.Agric.Sci.Tech. 12: 223-231.
- Rangaswamy, M., Latha, R., Jayamani, P., Annadurai, A., and Thiyagarajan, K. 1998. Combining ability studies in two line rice hybrids. In *Breeding strategies for 21<sup>st</sup> century*, Proceedings of the First National Plant Breeding Congress, July 1<sup>st</sup>-3<sup>rd</sup>, 1998. Coimbatore, India: Indian Society of Plant Breeders, pp. 95-101.
- Rogbell, E.J. 1995. Studies on genetics of salt tolerance in rice (*Oryza sativa* L.) cultivars [dissertation]. Coimbatore (TN): Tamil Nadu Agricultural University; 1995.
- Rogbell EJ, Subbaraman N. 1997. Line x Tester analysis for combining ability in saline rice cultivars. Madras Agric. J. 84 (1): 22-25.
- Sampoornam, R., and Thiyagarajan, K. 1998. Combining ability for yield and yield related traits in two line hybrid rice (*Oryza sativa* L.). In *First Plant Breeding Congress*. Coimbatore: TNAU, pp-93.
- Shi MS, Deng JY. 1986. The discovery determination and utilization of the Hubei photosensitive genic male sterile rice (*Oryza sativa* L. subsp. *Japonica*). Acta Genetica Sinica. 13(2): 107-112.
- Tagawa T, Ishizaka N. 1963. Physiological studies on the tolerant rice plants to salinity. Proc. Crop Soc. 31: 249-252.
- Tan ZC, Li YY, Chen LB, Zhou GQ. 1990. Studies on ecological and adaptability of dual purpose line An-Nong S-1. Hybrid Rice. 1 : 1-3.
- Thirumeni S. Genetical studies on coastal saline rice (*Oryza sativa* L.) [dissertation]. Coimbatore (TN): Tamil Nadu Agricultural University; 1998.
- Thirumeni S, Subramanian M, Paramasivam K. 2000. Combining ability and gene action in rice under salinity. Tropical Agricultural Research. 12: 375-385.
- Thirumeni S, Subramanian M, Paramasivam K. 2003. Genetics of salt tolerance in rice (*Oryza sativa* L.). Indian J. Genet. 63(1): 75-76.
- Yeo, A.R., and Flowers, T.J. 1984. Mechanisms of salinity resistance in rice and their role as physiological criteria in plant breeding. In *Salinity tolerance in plant strategies for crop improvement*, Eds. R.C. Staples and G.H. Toenniessen. New York: John Wiley & Sons, pp. 151-170.
- Yoshida, S., Forno, D.A., Cock, S.H. and Gomez K.A. 1976. *Laboratory manual for physiological studies in Rice*. Manila, Philippines: IRRI.
- Yuan Longping. 2004. Hybrid rice for food security in the world. In *FAO Rice conference*, 12-13<sup>th</sup> Feb. 2004, Rome, Italy: FAO.