

EFFECT OF PLANTING PATTERN ON RADIATION USE EFFICIENCY, YIELD AND YIELD COMPONENTS OF SUNFLOWER

Beytollah Vahedi¹, Abdolghayoum Gholipouri², Mohammad Sedghi²

MSc student in Agriculture faculty of University of Mohaghegh Ardabili, Iran

Academic member in Agriculture faculty of University of Mohaghegh Ardabili, Iran

Abstract

In order to determinate suitable planting pattern of sunflower, a field experiment was done in research field of university of Mohaghegh Ardabili in 2009. Factorial experiment was done based on randomized completely block design with three replication and two factor includes cultivars with three levels (Armavirusky, Azargol and Progress) and planting pattern with three levels (75*16.5, 60*20.5 and 45*27.5) with constant density (8 plant.m⁻²). Results of study showed that planting pattern effect was significant on head diameter and radiation use efficiency but have no significantly effect on yield and yield components. Cultivar effect was significant on yield and yield components but has no significantly on head diameter and radiation use efficiency. Among the cultivars studied, the highest grain yield (3616 kg.ha⁻¹) belonging to Azargol cultivar and lowest grain yield (3131 kg.ha⁻¹) belong to Progress cultivar. Therefore, it is better that according to exist facilities and machinery suitable planting pattern shall be selected and in the study area Azargol cultivar being planted.

Key Words: planting pattern, sunflower, yield, radiation use efficiency.

Introduction

Sunflower (*Helianthus annus* L.) is annual crop from compositeae family which grows firm. During its growth depending on cultivar and environmental factors and between 90 to 150 days is variable [1].

Uniform plant distribution increased productivity of plants to environmental factors and it can be increased branch's productivity and yield [7, 8, 9 and 10]. Planting in rows thinner in comparison with larger row distance in optimal conditions, yield is increased [11, 12, 13 and 14]. Arashi and Jaafari [2] reported the most suitable planting pattern for mechanized cultivation in Mehr cultivar is 60×15. Abdolrahmani [3] in an experiment concluded that 80 thousand plant densities per hectar is recommended for Record cultivar in Isfahan plain. Khalifeh [4] reported that the number of seeds with increasing distance between rows increased. Chogan [5] in an experiment on different planting pattern on rainfed sunflower in Gorgan concluded that planting pattern 75×30 is suitable. Tae [6] also reported that the planting pattern 75×40 cm for dryland in Urmia for Armavirusky cultivar with mean yield 893 kg.h⁻¹ is the best density.

Leaf area index is one of the most important effective factors on grain yield. If the leaf area index in short time to reach desired level. The maximum yield is obtained. Slowly increased leaf cause weak crop cover and therefore less radiation is absorbed. Finally reduced plant growth rate will be followed [14]. Appropriate planting pattern can compensate for lower leaf area index.

Materials and Methods

This experiment was done in research field of university of Mohaghegh Ardabili in 2008. Factorial experiment was done based on randomized completely block design with three replication and two factor includes cultivars with three levels (Armavirusky, Azargol and Progress) and planting pattern with three levels (75×16.5, 60×20.5 and 45×27.5) with constant density (8 plant.m⁻²). Each plot experiment includes 5 rows of 4 m in length. After preparing the land value of 150 kg of phosphate and potassium fertilizer to the land was given as base fertilizer. 150 kg of N fertilizer two time was given the land. First 100 kg with base fertilizer and second time the rate of 50 kg was used at the time of flowering. Inside each hole at planting time three seeds were placed. Plants were thinned when the plants had 3 to 4 leaves. Irrigation during the growth was done based on plant need.

Intercepted radiation plant community was measured by using of Sunscan. Data collection was done at 12 pm. The radiation below canopy one in between rows and once on the rows were measured and their mean as the radiation passing was calculated. Accumulated radiation during the growth period was calculated by using of meteorological data.

For measuring yield and yield components when the behind of receptacle was brown, 8 plants of each plot were harvested with considering margins. Analyzing data, calculating correlation coefficient and drawing graphs was done by using of Mstat- c, SPSS and Excel respectively.

different (Table 2). This can be due to the larger receptacle and smaller seeds.

Cultivar effect on the total grain weight, filled and unfilled seed weight and unfilled percentage were significant (Table 1). Effect of cultivar on total seed weight, filled and unfilled seed weight and unfilled seed percentage was significant (Table 1). Maximum total seeds weight (45.2 gr) and filled seeds weight of receptacle belong to Azargol and however Armavirsky and Progress had more seed number but its seeds had less weight. Minimum unfilled seed weight and unfilled seed percentage belong to Azargol and were significant with two other cultivar (Table 2) but maximum unfilled seed weight belong to Progress cultivar and has no significant with Armavirsky cultivar.

Taghdiri et al [18] reported that lowest unfilled seed number belongs to Azargol cultivar in a study. Differences between cultivars in yield are related to genetic characteristics, percentage fertility flowers and insect pollinator activity. Therefore Azogol with having large flowers percentage of empty seed is less. Cultivar effect on seed thousand weight and harvest index was significant (Table 1). Maximum seed thousand weight (61.31 gr) belongs to Armavirsky cultivar that was not significant with Progress cultivar (47.19 gr). Taghdiri et al [18] reported that among cultivars studied the highest seed hundred weight (5.61 gr) belonging to Azargol. Harvest index of Azargol (36.72%) was superior of the two other cultivars. Harvest index with traits such as seed thousand weight ($r=0.71^{**}$), total grain weight ($r=0.58^{**}$) and seed yield ($r=0.58^{**}$) showed a significant positive correlation and with traits such as plant height ($r=-0.45^*$), unfilled seed percentage ($r=-0.56^{**}$), unfilled seed weight ($r=-0.58^{**}$) and shoot dry weight ($r=-0.71^{**}$) significant negative correlation (table 3).

Cultivar effect on shoot dry weight was significant (Table 1). Maximum dry weight (1165 g.m⁻²) belongs to Armavirsky cultivar. Seed yield was affected by cultivar and maximum and minimum seed yield belong to Azogol and Progress cultivars respectively (table 2).

Azargol cultivar had less dry weight and seed number in comparison with other two cultivars can be concluded that the superiority in grain yield is related to more seed thousand weight, harvest index and less unfilled seed.

Conclusion

The results of this study showed that Azogol cultivar with producing less foliage and higher seed yield than other cultivars studied is significant superior. Because between studied planting patterns was no significant differences in seed yield, then the planting pattern 16.5×75 is superior to other planting pattern, because of the availability of machinery for this type of

row spacing. Therefore Azogol cultivar with planting pattern 16.5×75 is recommended for this area.

References

1. Khajeh poor, M. R. 1994. Industrial crop production. Jahad Danshgahi Sanaati Esfahan. PP: 250.
2. Khalilismamani, M. R., Khajehpoor, M. R., Ghalavand, A. 1998. The effect of row spacing and plant density in row on growth and dry matter accumulation in cotton at Esfahan. Majaleh olom keshavarzi Iran, 29: 667-669.
3. Azari, A., Khajehpoor, M. R. 2005. Effect of planting pattern on development, growth, yield components and yields of seed and petal calycle in summer planting of Safflower, local mass of the kooseh Esfahan. Magalleh olum va fonon keshavarzi va manabe tabii, year 9, 3: 131-141.
4. Andrade, C. A. D., Constantin, J., Scapim, C. A., Ebraccini, A. D., Angelotti, F., 2001. Effect of weed competition in different spacing upon yield of three common bean (*Phaseolus vulgaris* L.) cultivars. Ciencia, Eagrotecnologica. 43: 561-568.
5. Board, J.E., Kamal, M., Harville, B. G., 1992. Temporal importance of greater light interception to increased yield in narrow-row soybean. Agron. J. 84: 575-579.
6. Fronza, V., Vieira, C., Cardoso, A. A., Cruz, C. D., Pereira, P. R. G., 1994. Response of erect bean (*Phaseolus Vulgaris* L.) cultivars to spacing and rates of mineral fertilizers. Revista Ceres. 41: 567-683.
7. Sharma, H. P., 1995. Response of Raj mash to row spacing on salt affected soil. Indian Journal of pulses Research. 8: 93-94.
8. Das, S. N., Mukherjee, A. K., Nanda, M. K., 1996. Effect of dates of sowing and row spacing on yield attributing factors of different varieties of French bean (*Phaseolus Vulgaris*). Agri. Digest Karnal. 16: 130-132.
9. Arshi, y., jafari, y. 1992. The studying of the most appropriate plant density in sunflower cv. Mehr at watery codition. Mjaleh tahghighat keshavarzi nahal va bazr, 8: 1-7.
10. Abdolrahmani, b. 2002. Effect of plant density on yield and agronomic traits of sunflower cv. Armavirsky under Dryland condition in Maragheh. Majaleh olom zerai Iran, 5: 216-224.
11. Khalifa, F. M., 1984. Effects of spacing on growth and yield of sunflower under two systems of dry farming in Sudan. J. Of Agri. Sci. 103: 213-222.
12. Jogan, R. 1992. The studying effect of plant density on sunflower yield at rainfed condition. Majaleh nahal va bazr. 8. Seed and plant improvement institute.

13. Tai, A. 1994. The study effect of plant density on sunflower at rainfed condition. Third congress agronomy and plant breeding sciences Iran, university of Tabriz.
14. Ganjali, A., Malekzadeh, S., Bagheri, A. 2000. The studying of plant density and planting pattern on changes process growth indices chickpea under watery condition in zone of Neishabour. *Majaleh olom va sanayeh gazai keshavarzi*, 14:33-41.
15. Atlasi pak, V., Mamghani, R., Meskarbashi, M., Nabipour, M. 2006. Effect of planting pattern on radiation use efficiency and dry matter accumulation in three cultivars canopy spring rapeseed. *majaleh elmi keshavarzi*, 29: 139-152.
16. Mazaherilaghab, H., Moradi, H., Sepehri, A. 2005. Evaluation effect of planting date and pattern on yield of sunflower cv. Armavirsky in zone of Marivan. *Majaleh pajohesh va sazanegi*, 18: 31-39.
17. Agele, S. O., Mraiyesa, I. O., Adeniji, I. A., 2007. Effects of variety and row spacing on radiation interception, Partitioning of dry matter and seed set efficiency in late season sunflower (*Helianthus annuus* L.) in a humid zone of Nigeria. *African Journal of Agriculture Research*. 3: 80-88.
18. Taghdiri, B., Ahmadvand, G., Mazaherilaghab, H. 2006. The effect of plant spacing on yield and yield components of four sunflower cultivars. *Majaleh pajohesh ;shavarzi, ab, khak va gyah dar keshavarzi*. 6: 26-35.
19. Pourdad, S. 2002. Study sunflower hybrids at rainfed condition. Seventh congress of the agronomy and plant breeding sciences in Iran, Karaj.