



BOTANY

ARTEMISININ CONTENT IN *ARTEMISIA SCOPARIA*

Aditi Singh* and Renu Sarin

Laboratory of Bioactive Compounds, Department of Botany, University of Rajasthan, Jaipur-302004 (INDIA)

Abstract

Artemisinin is considered as the natural, active and potent antimalarial drug *Artemisia annua* Linn. is the only known source for the production of artemisinin reported till date. An attempt was made on a perennial faintly odoratus herb, *A. scoparia* Waldst et Kit. to find out an alternative of *A. annua* for the production of artemisinin. Generation of hairy roots was also tried with an attempt to increase the concentration of artemisinin in the biofactories. In the present investigation *A. scoparia* was shown to contain artemisinin

Keywords: *A. scoparia*, *Agrobacterium rhizogenes*, antimalarial drug; artemisinin

Introduction

Malaria is one of the world's most important parasitic diseases. There are at least 300 million acute cases of malaria each year globally, resulting in more than a million deaths. Multi-drug resistance of the *Plasmodium* strains to the cheapest and most widely used antimalarials such as chloroquine, mefloquine and sulfadoxine-pyrimethamine is one of the biggest challenges in the fight against malaria.

Artemisia annua L. (sweet wormwood), a herb of the Asteraceae family has been used for centuries for the treatment of fever and malaria. This plant is the only known source for the production of artemisinin; a potent antimalarial. Artemisinin is the new and promising drug which is also active against resistant plasmodium malaria (15). As artemisinin cannot be synthesized chemically in an economically feasible way, *A. annua* is the only practical source of this valuable drug. Extensive work on the production of artemisinin from plant of *Artemisia annua* has been done (2,3,5,6,7,8,9,10,13,18,19). Artemisinin has also been reported from tissue culture of this plant (11,12,13,14,16,17) and biochemical as well as molecular approaches for enhanced production of artemisinin from *A. annua* has also been tried (1,17)

The present investigation has reported *A. scoparia* Waldst et Kit as a new source of artemisinin. Hairy root generation was also attempted in the present study with an objective to increase the content of artemisinin in the tissue culture of *A. scoparia*

Materials and Methods

Unorganized callus of *A. scoparia* was raised from the young nodal stem segments. Callus of the plant were obtained on MS medium supplemented with 2, 4-D (3mg/L), kinetin (0.25mg/L) and proline (100mg/L) Callus initiation started after 15-20 days of inoculation ;

4-6 weeks old callus as well as aerial plant parts were harvested and subjected to extraction procedures for artemisinin separately.

Aerial plant parts and harvested callus were dried and powdered and extracted by continuous percolation over a period of four to six hours using five to ten fold volume of the non-aqueous solvent ethanol. The extraction process was repeated three to five times to ensure maximum extraction of artemisinin from the herb. The resulting extract was concentrated to 1 to 5% of the original volume. The excess of water (four times of the reduced volume of ethanol extract) to be added to the concentrated to make it 80% aqueous followed by partitioning of the contents between water and hexane. For partitioning, the aqueous content and hexane may be used in a ratio of 1:1 or 2:1 v/v. Partitioning of aqueous content with hexane may be repeated three to five times using the same solvent ratio in order to ensure maximum transfer of artemisinin to hexane fraction. The combined hexane fractions were pooled together to obtain 1-5% of its original volume. The concentrated liquid was a light to dark green oily liquid. Ethyl acetate (10-20% v/v) is added to it. To remove the green pigmentation this liquid was treated with 1-3% w/v of activated charcoal. The yellowish liquid obtained after removal of activated charcoal (by filtration) was subjected to the evaporative crystallization yielding substantially pure artemisinin. The purified crystals thus obtained were further analyzed for artemisinin by infra-red spectroscopy, thin layer chromatography and nuclear magnetic resonance

Thin layer chromatography

Thin Layer Chromatography (TLC) was performed as follows: The extract so obtained were dissolved in ethanol and applied separately 1 cms above the edge

* Corresponding Author, Email: aditisingh_21@yahoo.co.in




Figure 5. Circular Ri plasmid of *A. rhizogene* strain A4 showing the different genetic regions.

The present investigation confirms that aerial plant parts and callus cultures of *A. scoparia* Waldst et Kit. has the potentiality to produce artemisinin. This study concluded that *A. scoparia* besides possessing antibacterial and insecticidal properties also contains artemisinin. Thus, the plants of *A. scoparia* hold a new promise to the field of medicinally and economically important bioactive compounds which should be explored further to open new vistas for mankind.

But being unable to induce hairy root formation in the calli it could be concluded that for hairy root induction some elicitor like acetosyringone is required or it could be attempted with a different strain of *A. rhizogenes*. Thus, with the help of advanced techniques there is a probability that the amount of artemisinin in *A. scoparia* could be increased and used in future for curing the deadly parasitic disease in humans.

Acknowledgements

The authors are grateful to Head, Department of Botany and Head, Department of Chemistry for providing the lab facilities to carry out this work and Dr. Shashi Pandey Rai, Banaras Hindu University, Varanasi, India for providing the *A. rhizogenes* A4 strain.

References

1. Abdin M.Z., Israr M. Rehman R.U. and Jain. S.K, Artemisinin, a novel antimalarial drug: Biochemical and molecular approaches for enhanced production. *Planta Med.* 69, 2003;289-293.
2. Acton N. and Klayman D.L.; Artemisinin, a new sesquiterpene lactone endoperoxide from *Artemisia annua*. *Planta Medica.* 47, 1985,442-445
3. Acton N., Klayman D.L. and Rollman I.J.; Reductive electrochemical HPLC assay for artemisinin (qinghaosu). *Planta Med.* 51, 1985, 445-446
4. de Vries P. Dien J.T.K., Clinical pharmacology and therapeutic potential of artemisinin and its derivatives in the treatment of malaria, *Drugs.* 52, 1996, 818-836.
5. El-Sohly, H.N. A large scale extraction technique of artemisinin from *Artemisia annua*. *J. Nat. Prod.*; 53,1990,1560-1564.
6. Ferreira J.F.S and Janick J.; Immunoquantitative analysis of artemisinin from *Artemisia annua* using polyclonal antibodies. *Phytochemistry* 41, 1996a, 97-104.
7. Ferreira J.F.S., Charles D.J , Wood K., Simon J.E and Janick J.; A comparison of gas chromatography and high performance liquid chromatography for artemisinin analyses. *Phytochem. Anal.* 5, 1994,116-120.
8. Ferreira J.F.S. and Janick J.; Roots as an enhancing factor for the production of artemisinin in shoot cultures of *Artemisia annua*. *Plant Cell Tissue Organ Cult.* 44, 1996 b ,211-217.
9. Ferreira J.F.S., Simon J.E. and Janick J, Developmental studies of *Artemisia annua*: Flowering and artemisinin production under greenhouse and field conditions. *Planta Med* 61, 1995 a;167-170.
10. Ferreira J.F.S., Simon J.E. and Janick J.; Relationship of artemisinin content of tissue-cultured, greenhouse-grown, and field-grown plants of *Artemisia annua*. *Planta Med.* 61, 1995 b ,351-355.
11. Fulzele D.P, Sipahimalani. A.T and Heble M.R , Tissue cultures of *Artemisia annua*: organogenesis and artemisinin production. *Phytother. Res.*5, 1991;149-153.
12. Jaziri M., Shimomura K., Yoshimatsu K., Fauconnier M.L., Marlier M. and Homes J., Establishment of normal and transformed root cultures of *Artemisia annua* L. for artemisinin production. *J. Plant Physiol.*145, 1995;175-177.
13. Jha J. , Jha T.B. and Mahato S.B.; Tissue culture of *Artemisia annua* L.: A potential source of an antimalarial drug . *Curr. Sci.*57,1988;344-346.
14. Martinez B.C. and Staba E.J.; The production of artemisinin in *Artemisia annua* L. tissue cultures. *Adv. Cell Cult.* 6, 1988;69-87.
15. McIntosh H.M and Olliaro P. Artemisinin derivatives for treating severe malaria (Cochrane Review). *The Cochrane Library* 2. Oxford: Update Software. 2001.

16. Nair, M.S.R. , Acton. N, Klayman D.L., Kendrick K., Basile D.V. and Mante S; Production of artemisinin in tissue cultures of *Artemisia annua*. J. Nat. Prod.49, 1986;504-507.
17. Nieuwerburgh F.C.W.V, Castele S.R.F.V., Maes L., Goossens A., Inze J.V, Bocxlaer D.. and Deforce D.L.D.; Quantitation of artemisinin and its biosynthetic precursors in *Artemisia annua* L. by High Performance Liquid Chromatography-electrospray quadrupole time of flight tandem mass spectrometry. Journal of Chromatography 1118, 2006;180-187.
18. Theoharides A.D. Smyth M.H., Ashmore R.W., Halverson J.M., Zhou Z.M., Ridder W.E.and Lin. A.J, Anal. Chem. 60, 1988;115.
19. Titulaer, H.A. , Zuidema J. , Kager P.A., Wetsteyn J.C, Lugt C.B and Merkus F.W. , The pharmacokinetics of artemisinin after oral, intramuscular and rectal administration to volunteers. J Pharm Pharmacol 42, 1990; 810–813.